


London Mathematical Society
Lecture Note Series 302

Discrete and Continuous Nonlinear Schrödinger Systems

M. J. Ablowitz, B. Prinari and A. D. Trubatch



**London
Mathematical
Society**

BRIDGE
UNIVERSITY PRESS

London Mathematical Society Lecture Note Series. 302

Discrete and Continuous Nonlinear Schrödinger Systems

M. J. Ablowitz

University of Colorado, Boulder

B. Prinari

Università di Lecce

A. D. Trubatch

United States Military Academy, West Point



CAMBRIDGE
UNIVERSITY PRESS

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS

The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York, NY 10011-4211, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa
<http://www.cambridge.org>

© Mark J Ablowitz Barbara Prinari A. David Trubatch 2004

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2004

Printed in the United Kingdom at the University Press, Cambridge

Typeset Times 10/13 pt. and Helvetica *System* L^AT_EX 2_ε [TB]

A catalog record for this book is available from the British Library.

Library of Congress Cataloging in Publication Data

Ablowitz, Mark J.

Discrete and continuous nonlinear Schrödinger systems / M.J. Ablowitz, B. Prinari,
A.D. Trubatch.

p. cm. – (London Mathematical Society lecture note series ; 302)

Includes bibliographical references and index.

ISBN 0-521-53437-2 (pbk.)

I. Schrödinger equation. 2. Nonlinear theories. 3. Inverse scattering transform.

I. Prinari, B., 1972– II. Trubatch, A. D., 1968– III. Title. IV. Series.

QC174.26.W28A26 2003

530.12'4–dc21 2003048555

ISBN 0 521 53437 2 paperback

Managing Editor: Professor N.J. Hitchin, Mathematical Institute,
University of Oxford, 24–29 St Giles, Oxford OX1 3LB, United Kingdom

The titles below are available from booksellers, or from Cambridge University Press at www.cambridge.org

- 100 Stopping time techniques for analysts and probabilists. L. EGGHE
- 105 A local spectral theory for closed operators. I. ERDELYI & WANG SHENGWANG
- 107 Compactification of Siegel moduli schemes. C.-L. CHAI
- 109 Diophantine analysis. J. LOXTON & A. VAN DER POORTEN (eds)
- 113 Lectures on the asymptotic theory of ideals. D. REES
- 116 Representations of algebras. P.J. WEBB (ed)
- 119 Triangulated categories in the representation theory of finite-dimensional algebras. D. HAPPEL
- 121 Proceedings of *Groups – St Andrews 1985*. E. ROBERTSON & C. CAMPBELL (eds)
- 128 Descriptive set theory and the structure of sets of uniqueness. A.S. KECHRIS & A. LOUVEAU
- 130 Model theory and modules. M. PREST
- 131 Algebraic, extremal & metric combinatorics. M.-M. DEZA, P. FRANKL & I.G. ROSENBERG (eds)
- 141 Surveys in combinatorics 1989. J. SIEMONS (ed)
- 144 Introduction to uniform spaces. I.M. JAMES
- 146 Cohen-Macaulay modules over Cohen-Macaulay rings. Y. YOSHINO
- 148 Helices and vector bundles. A.N. RUDAKOV *et al*
- 149 Solitons, nonlinear evolution equations and inverse scattering. M. ABLowitz & P. CLARKSON
- 150 Geometry of low-dimensional manifolds 1. S. DONALDSON & C.B. THOMAS (eds)
- 151 Geometry of low-dimensional manifolds 2. S. DONALDSON & C.B. THOMAS (eds)
- 152 Oligomorphic permutation groups. P. CAMERON
- 153 L-functions and arithmetic. J. COATES & M.J. TAYLOR (eds)
- 155 Classification theories of polarized varieties. TAKAO FUJITA
- 158 Geometry of Banach spaces. P.F.X. MÜLLER & W. SCHACHERMAYER (eds)
- 159 Groups St Andrews 1989 volume 1. C.M. CAMPBELL & E.F. ROBERTSON (eds)
- 160 Groups St Andrews 1989 volume 2. C.M. CAMPBELL & E.F. ROBERTSON (eds)
- 161 Lectures on block theory. BURKHARD KÜLSHAMMER
- 163 Topics in varieties of group representations. S.M. VOVSİ
- 164 Quasi-symmetric designs. M.S. SHRIKANDÉ & S.S. SANE
- 166 Surveys in combinatorics, 1991. A.D. KEEDWELL (ed)
- 168 Representations of algebras. H. TACHIKAWA & S. BRENNER (eds)
- 169 Boolean function complexity. M.S. PATERSON (ed)
- 170 Manifolds with singularities and the Adams-Novikov spectral sequence. B. BOTVINNIK
- 171 Squares. A.R. RAJWADE
- 172 Algebraic varieties. GEORGE R. KEMPF
- 173 Discrete groups and geometry. W.J. HARVEY & C. MACLACHLAN (eds)
- 174 Lectures on mechanics. J.E. MARSDEN
- 175 Adams memorial symposium on algebraic topology 1. N. RAY & G. WALKER (eds)
- 176 Adams memorial symposium on algebraic topology 2. N. RAY & G. WALKER (eds)
- 177 Applications of categories in computer science. M. FOURMAN, P. JOHNSTONE & A. PITTS (eds)
- 178 Lower K- and L-theory. A. RANICKI
- 179 Complex projective geometry. G. ELLINGSRUD *et al*
- 180 Lectures on ergodic theory and Pesin theory on compact manifolds. M. POLİ ICOTT
- 181 Geometric group theory I. G.A. NIBLO & M.A. ROLLER (eds)
- 182 Geometric group theory II. G.A. NIBLO & M.A. ROLLER (eds)
- 183 Shintani zeta functions. A. YUKIE
- 184 Arithmetical functions. W. SCHWARZ & J. SPILKER
- 185 Representations of solvable groups. O. MANZ & T.R. WOLF
- 186 Complexity: knots, colourings and counting. D.J.A. WELSH
- 187 Surveys in combinatorics, 1993. K. WALKER (ed)
- 188 Local analysis for the odd order theorem. H. BENDER & G. GLAUBERMAN
- 189 Locally presentable and accessible categories. J. ADAMEK & J. ROSICKY
- 190 Polynomial invariants of finite groups. D.J. BENSON
- 191 Finite geometry and combinatorics. F. DE CLERCK *et al*
- 192 Symplectic geometry. D. SALAMON (ed)
- 194 Independent random variables and rearrangement invariant spaces. M. BRAVERMAN
- 195 Arithmetic of blowup algebras. WOLMER VASCONCELOS
- 196 Microlocal analysis for differential operators. A. GRIGIS & J. SJÖSTRAND
- 197 Two-dimensional homotopy and combinatorial group theory. C. HOG-ANGELONI *et al*
- 198 The algebraic characterization of geometric 4-manifolds. J.A. HILLMAN
- 199 Invariant potential theory in the unit ball of C^n . MANFRED STOLL
- 200 The Grothendieck theory of dessins d'enfant. L. SCHNEPS (ed)
- 201 Singularities. JEAN-PAUL BRASSELET (ed)
- 202 The technique of pseudodifferential operators. H.O. CORDES
- 203 Hochschild cohomology of von Neumann algebras. A. SINCLAIR & R. SMITH
- 204 Combinatorial and geometric group theory. A.J. DUNCAN, N.D. GILBERT & J. HOWIE (eds)
- 205 Ergodic theory and its connections with harmonic analysis. K. PETERSEN & I. SALAMA (eds)
- 207 Groups of Lie type and their geometries. W.M. KANTOR & L. DI MARTINO (eds)
- 208 Vector bundles in algebraic geometry. N.J. HITCHIN, P. NEWSTEAD & W.M. OXBURY (eds)
- 209 Arithmetic of diagonal hypersurfaces over finite fields. F.Q. GOUVEA & N. YUI
- 210 Hilbert C^* -modules. E.C. LANCE
- 211 Groups 93 Galway / St Andrews I. C.M. CAMPBELL *et al* (eds)
- 212 Groups 93 Galway / St Andrews II. C.M. CAMPBELL *et al* (eds)

- 214 Generalised Euler-Jacobi inversion formula and asymptotics beyond all orders. V. KOWALENKO *et al*
- 215 Number theory 1992-93. S. DAVID (ed)
- 216 Stochastic partial differential equations. A. ETHERIDGE (ed)
- 217 Quadratic forms with applications to algebraic geometry and topology. A. PFISTER
- 218 Surveys in combinatorics, 1995. PETER ROWLINSON (ed)
- 220 Algebraic set theory. A. JOYAL & I. MOERDIJK
- 221 Harmonic approximation. S.J. GARDINER
- 222 Advances in linear logic. J.-Y. GIRARD, Y. LAFONT & L. REGNIER (eds)
- 223 Analytic semigroups and semilinear initial boundary value problems. KAZUAKI TAIRA
- 224 Computability, enumerability, unsolvability. S.B. COOPER, T.A. SLAMAN & S.S. WAINER (eds)
- 225 A mathematical introduction to string theory. S. ALBEVERIO *et al*
- 226 Novikov conjectures, index theorems and rigidity I. S. FERRY, A. RANICKI & J. ROSENBERG (eds)
- 227 Novikov conjectures, index theorems and rigidity II. S. FERRY, A. RANICKI & J. ROSENBERG (eds)
- 228 Ergodic theory of Z^d actions. M. POLLICOTT & K. SCHMIDT (eds)
- 229 Ergodicity for infinite dimensional systems. G. DA PRATO & J. ZABCZYK
- 230 Prolegomena to a middlebrow arithmetic of curves of genus 2. J.W.S. CASSELS & E.V. FLYNN
- 231 Semigroup theory and its applications. K.H. HOFMANN & M.W. MISLOVE (eds)
- 232 The descriptive set theory of Polish group actions. H. BECKER & A.S. KECHRIS
- 233 Finite fields and applications. S. COHEN & H. NIEDERREITER (eds)
- 234 Introduction to subfactors. V. JONES & V.S. SUNDER
- 235 Number theory 1993-94. S. DAVID (ed)
- 236 The James forest. H. FETTER & B. GAMBOA DE BUEN
- 237 Sieve methods, exponential sums, and their applications in number theory. G.R.H. GREAVES *et al*
- 238 Representation theory and algebraic geometry. A. MARTSINKOVSKY & G. TODOROV (eds)
- 240 Stable groups. FRANK O. WAGNER
- 241 Surveys in combinatorics, 1997. R.A. BAILEY (ed)
- 242 Geometric Galois actions I. L. SCHNEPS & P. LOCHAK (eds)
- 243 Geometric Galois actions II. L. SCHNEPS & P. LOCHAK (eds)
- 244 Model theory of groups and automorphism groups. D. EVANS (ed)
- 245 Geometry, combinatorial designs and related structures. J.W.P. HIRSCHFELD *et al*
- 246 p -Automorphisms of finite p -groups. E.I. KHUKHRO
- 247 Analytic number theory. Y. MOTOHASHI (ed)
- 248 Tame topology and o-minimal structures. LOU VAN DEN DRIES
- 249 The atlas of finite groups: ten years on. ROBERT CURTIS & ROBERT WILSON (eds)
- 250 Characters and blocks of finite groups. G. NAVARRO
- 251 Gröbner bases and applications. B. BUCHBERGER & F. WINKLER (eds)
- 252 Geometry and cohomology in group theory. P. KROPHOLLER, G. NIBLO, R. STÖHR (eds)
- 253 The q -Schur algebra. S. DONKIN
- 254 Galois representations in arithmetic algebraic geometry. A.J. SCHOLL & R.L. TAYLOR (eds)
- 255 Symmetries and integrability of difference equations. P.A. CLARKSON & F.W. NIJHOFF (eds)
- 256 Aspects of Galois theory. HELMUT VÖLKLEIN *et al*
- 257 An introduction to noncommutative differential geometry and its physical applications 2ed. J. MADORE
- 258 Sets and proofs. S.B. COOPER & J. TRUSS (eds)
- 259 Models and computability. S.B. COOPER & J. TRUSS (eds)
- 260 Groups St Andrews 1997 in Bath, I. C.M. CAMPBELL *et al*
- 261 Groups St Andrews 1997 in Bath, II. C.M. CAMPBELL *et al*
- 262 Analysis and logic. C.W. HENSON, J. IOVINO, A.S. KECHRIS & E. ODELL
- 263 Singularity theory. BILL BRUCE & DAVID MOND (eds)
- 264 New trends in algebraic geometry. K. HULEK, F. CATANESE, C. PETERS & M. REID (eds)
- 265 Elliptic curves in cryptography. I. BI-KE, G. SEROUSSI & N. SMART
- 267 Surveys in combinatorics, 1999. J.D. LAMB & D.A. PREECE (eds)
- 268 Spectral asymptotics in the semi-classical limit. M. DIMASSI & J. SJÖSTRAND
- 269 Ergodic theory and topological dynamics. M.B. BEKKA & M. MAYER
- 270 Analysis on Lie groups. N.T. VAROPOULOS & S. MUSTAPHA
- 271 Singular perturbations of differential operators. S. ALBEVERIO & P. KURASOV
- 272 Character theory for the odd order theorem. T. PETERFALVI
- 273 Spectral theory and geometry. E.B. DAVIES & Y. SAFAROV (eds)
- 274 The Mandelbrot set, theme and variations. TAN LEI (ed)
- 275 Descriptive set theory and dynamical systems. M. FOREMAN *et al*
- 276 Singularities of plane curves. E. CASAS-ALVERO
- 277 Computational and geometric aspects of modern algebra. M.D. ATKINSON *et al*
- 278 Global attractors in abstract parabolic problems. J.W. CHOLEWA & T. DŁOTKO
- 279 Topics in symbolic dynamics and applications. F. BLANCHARD, A. MAASS & A. NOGUEIRA (eds)
- 280 Characters and automorphism groups of compact Riemann surfaces. THOMAS BREUER
- 281 Explicit birational geometry of 3-folds. ALESSIO CORTI & MILES REID (eds)
- 282 Auslander-Buchwitz approximations of equivariant modules. M. HASHIMOTO
- 283 Nonlinear elasticity. Y. FU & R.W. OGDEN (eds)
- 284 Foundations of computational mathematics. R. DEVORE, A. ISERLES & E. SÜLI (eds)
- 285 Rational points on curves over finite fields. H. NIEDERREITER & C. XING
- 286 Clifford algebras and spinors 2ed. P. LOUNESTO
- 287 Topics on Riemann surfaces and Fuchsian groups. E. BUJALANCE, A.F. COSTA & E. MARTINEZ (eds)
- 288 Surveys in combinatorics, 2001. J. HIRSCHFELD (ed)
- 289 Aspects of Sobolev-type inequalities. L. SALOFF-COSTE
- 290 Quantum groups and Lie Theory. A. PRESSLEY (ed)
- 291 Tits buildings and the model theory of groups. K. TENT (ed)
- 292 A quantum groups primer. S. MAJID
- 293 Second order partial differential equations in Hilbert spaces. G. DA PRATO & J. ZABCZYK
- 294 Introduction to the theory of operator spaces. G. PISIER

Preface

Nonlinear systems are generic in the mathematical representation of physical phenomena. It is unusual for one to be able to find solutions to most nonlinear equations. However, a certain physically significant subclass of problems admits deep mathematical structure that further allows one to find classes of exact solutions. Solitons are a particularly important subclass of such solutions. Solitons are localized waves that, in an appropriate sense, interact elastically with each other. They have proved to be extremely interesting to physicists and engineers due, in part, to their localized and stable nature.

This broad field of study is sometimes called “soliton theory” or “integrable systems.” This field has witnessed numerous important developments, which have been studied intensively worldwide over the past 30 years. Some of the directions that researchers have pursued include the following: direct methods to find solutions; studies of the underlying analytic structure of the equations; associated Painlevé-type solutions and relevant generalizations; tests to locate integrable systems; studies of the underlying geometric structures inherent in integrable systems; Bäcklund and Darboux transformations, which can be used to produce new classes of solutions; and so on.

In principle, one would like to be able to solve the general initial-value problem associated with these special nonlinear soliton systems. Depending on the boundary conditions under consideration, sometimes this is feasible. Moreover, in some cases, the mathematical representation is constructive, useful, and indeed elegant. The case for which the Cauchy problem admits a complete solution and the qualitative properties of the solution are well understood corresponds to initial data decaying sufficiently rapidly at infinity. Because of the constructive and relatively explicit nature of the solution, this case has received considerable attention. The method of solution is usually referred to as the inverse scattering transform (IST). The IST applies to many interesting nonlinear soliton systems, including nonlinear partial differential equations in $1 + 1$

(one space–one time) and $2 + 1$ (two space–one time) dimensions, nonlinear discrete (difference) evolution equations, and singular integro-differential equations. A discussion of the IST as it applies to many of these cases can be found in various monographs (cf. [6]).

Nevertheless, there are particular problems that, because of their wide applicability, deserve special attention. It is the purpose of this book to investigate one such important set of equations: nonlinear Schrödinger (NLS) systems. NLS systems in continuous media have been studied heavily since the mid-1960s. The continuous scalar NLS equation arises in the prototypical situation governing slowly varying waves of small amplitude (cf. [30]). In the early 1970s, it was shown [91] that the NLS equation governs the long-distance pulse propagation in optical fibers. In the late 1980s and 1990s, it was discovered [131], [122] that vector NLS systems govern the propagation of polarized waves in optical fibers. Today, these NLS equations, with suitable modifications and additional terms, are used routinely to make predictions about the transmission of information in fibers. Without doubt, wave transmission in optical fibers, used for communication purposes, is an application of critical importance.

In recent years there has developed significant interest in the study of nonlinear waves in media that are governed by nonlinear semi-discrete evolution systems (discrete in space–continuous time). Once again there is a class of physically interesting nonlinear systems that includes the so-called discrete NLS equations. These discrete NLS equations reduce to the continuous NLS equation when the discretization parameter vanishes.

For certain scalar- and vector-continuous and discrete NLS systems, the IST method can be applied in an effective and complete way. The initial-value problem, for given data decaying sufficiently rapidly at infinity, can be linearized in terms of integral equations. Multisoliton solutions can be obtained in all cases. This is described in detail in this book. In an appendix we also describe the IST associated with another class of important discrete equations: the Toda lattice [165] and the so-called nonlinear ladder network [121], [98]. While the research literature has many (but not all) of the results obtained here, it requires researchers to access numerous papers. Early fundamental research on NLS systems appears in the papers of Zakharov and Shabat [192], who first analyzed the scalar-continuous NLS equation; Manakov [120], regarding the continuous-vector NLS equation; and by Ablowitz and Ladik [11, 12], who introduced the scalar integrable discrete NLS equation. The vector extension of the discrete NLS system is more recent (cf. [18]). The unified description of these physically interesting integrable discrete and continuous NLS systems within the context of the IST methodology does not appear anywhere else. One of our motivations in writing this book was to develop the direct and inverse scattering

formalism based on the Riemann–Hilbert approach. From a pedagogical point of view, readers will find that the structure follows the one laid out for the Korteweg–de Vries equation in the monograph of Ablowitz and Clarkson [6]. Here we have also attempted to include many of the mathematical details, to make the book suitable for students as well as researchers who wish to study this topic.

We gratefully acknowledge support for these studies by the National Science Foundation, the Air Force Office of Scientific Research, and the Colorado Commission on Higher Education.

Contents

<i>Preface</i>	<i>page vii</i>
1 Introduction	1
1.1 Solitons and soliton equations	1
1.2 The inverse scattering transform – Overview	3
1.3 Nonlinear Schrödinger systems	5
1.4 Physical applications	9
1.5 Outline of the work	16
2 Nonlinear Schrödinger equation (NLS)	18
2.1 Overview	18
2.2 The inverse scattering transform for NLS	19
2.3 Soliton solutions	39
2.4 Conserved quantities and Hamiltonian structure	42
3 Integrable discrete nonlinear Schrödinger equation (IDNLS)	46
3.1 Overview	46
3.2 The inverse scattering transform for IDNLS	48
3.3 Soliton solutions	83
3.4 Conserved quantities and Hamiltonian structure	87
4 Matrix nonlinear Schrödinger equation (MNLS)	90
4.1 Overview	90
4.2 The inverse scattering transform for MNLS	91
4.3 Soliton solutions	113
4.4 Conserved quantities and Hamiltonian structure	128
5 Integrable discrete matrix NLS equation (IDMNLS)	130
5.1 Overview	130
5.2 The inverse scattering transform for IDMNLS	133
5.3 Soliton solutions	185
5.4 Conserved quantities	201

Appendices	
A Summation by parts formula	204
B Transmission of the Jost function through a localized potential	206
C Scattering theory for the discrete Schrödinger equation	208
C.1 Introduction	208
C.2 Direct scattering problem	210
C.3 Inverse scattering problem	220
C.4 Time evolution and solitons for the Toda lattice and nonlinear ladder network	226
D Nonlinear Schrödinger systems with a potential term	229
D.1 Continuous NLS systems with a potential term	229
D.2 Discrete NLS systems with a potential term	233
E NLS systems in the limit of large amplitudes	239
<i>Bibliography</i>	243
<i>Index</i>	255

Chapter 1

Introduction

1.1 Solitons and soliton equations

Ever since the observation of the “great wave of translation” in water waves, by J. Scott Russell in 1834 [146, 147] while he rode on horseback near a narrow canal in Edinburgh, localized (nonoscillatory) solitary waves have been known to researchers studying wave dynamics. Despite Russell’s detailed observations, it was many years before mathematicians formulated the relevant equation, now known as the Korteweg–de Vries (KdV) equation that governs those waves (cf. [38, 39, 112]). From the period 1895–1960, the study of water waves was essentially the only application in which solitary waves were found. However, in the 1960s it was discovered that the KdV equation is a relevant model in many other physical contexts, such as plasma physics, internal waves, lattice dynamics, and others. Critically, in their study of the Fermi–Pasta–Ulam lattice equation [75] Zabusky and Kruskal (1965) found that the KdV equation was the governing equation (cf. [189]). Moreover, in a wholly new discovery, Zabusky and Kruskal observed that the solitary waves of KdV are “elastic” in their interaction. That is, the solitary waves pass through one another and subsequently retain their characteristic form and velocity. Zabusky and Kruskal called these elastically interacting solitary waves *solitons*. The work of Gardner, Green, Kruskal, and Miura [82] showed how direct and inverse scattering techniques could be used to linearize the initial-value problem of KdV. The solitons also were shown to correspond to eigenvalues of the time-independent Schrödinger equation. The remarkable discovery of the soliton was the first in a chain of events that culminated in a mathematical theory of solitons in KdV. This work opened a rich vein of research that continues today, more than 35 years later. Further discussion of both the historical background and subsequent development of soliton theory can be found in [21].

Subsequent to the development of a mathematical theory of the solitons of KdV, further research revealed that solitons, with their distinctive elastic interactions, arise in numerous important physical systems. These systems are governed, respectively, by a diverse collection of evolution equations that are characterized mainly by the fact that they admit soliton solutions. For example, soliton solutions have been found in a number of nonlinear partial differential equations in $1 + 1$ dimensions (i.e., one space and one time dimension) and $2 + 1$ (i.e., two spatial dimensions and one time dimension). In addition, soliton solutions have been found in semi-discrete (discrete in space, continuous in time) and doubly discrete (discrete in space and time) nonlinear evolution equations and in nonlinear singular integro-differential equations, among others. A survey of some of these can be found in [6]. It should also be noted that there is a four-dimensional system, referred to as the self-dual Yang–Mills (SDYM) equations, that plays an important role in the study of soliton theory or integrable systems. Indeed, the SDYM equations can be viewed as a “master” integrable system from which virtually all other systems can be obtained as special reductions (cf. Atiyah and Ward [25]; Ward [181, 182, 183]; Belavin and Zakharov [29]; Mason et al. [124, 125]; Chakravarty et al. [50, 51]; Maszczyk et al. [126]; Ablowitz et al. [5]).

Researchers in physics and engineering have understood that stable localized solitary waves, even those that do not have the special property of elastic interaction, have many important applications, and their study has led to substantial research in specialized fields (e.g., nonlinear optics) all by itself. Nevertheless, the systems in which the solitary waves interact elastically, the “true” soliton systems, are important special cases. Moreover, there is an intrinsic richness in the mathematical theory of these soliton systems. Accordingly, the field of research associated with integrable systems has grown, developed, and expanded in many directions. One centrally important issue is the method of solution, sometimes referred to as the inverse scattering transform (IST), for these soliton equations. For a number of physically significant equations, the IST can be carried out in an explicit, effective, and illuminative manner. In particular, the IST is a fruitful approach, and it is the basis for the study of the nonlinear Schrödinger systems described in this book.

There are numerous books, review articles, and edited collections (see, for instance, [6, 21, 47, 65, 74, 140]) that delve widely and deeply into the theory of integrable systems. In this book, we give a detailed description of both the IST and the soliton solutions of integrable nonlinear Schrödinger systems, which are mathematically and physically important soliton equations. The collection of systems examined in this book comprises both continuous and semi-discrete systems of equations. As will be described in Section 1.4, these

particular systems arise in the modeling of a wide array of physical wave phenomena. In this book, we present most of the known results for these nonlinear Schrödinger systems, as well as some new ones, in a comprehensive, unified framework built with the mathematical machinery of the inverse scattering transform.

1.2 The inverse scattering transform – Overview

The IST is a method that allows one to linearize a class of nonlinear evolution equations. In doing so one can obtain global information about the structure of the solution. In many respects, one can view the IST as a nonlinear version of the Fourier transform.

The solution of the initial-value problem of a nonlinear evolution equation by IST proceeds in three steps, as follows:

1. the forward problem – the transformation of the initial data from the original “physical” variables to the transformed “scattering” variables;
2. time-dependence – the evolution of the transformed data according to simple, explicitly solvable evolution equations;
3. the inverse problem – the recovery of the evolved solution in the original variables from the evolved solution in the transformed variables.

In fact, with the IST machinery one can do more than solve the initial-value problem; one also can construct special solutions of the evolution equation by positing an elementary solution in the transformed variables and then applying the inverse transformation to obtain the corresponding solution in the original variables. In general, the soliton and multisoliton solutions of soliton equations can be constructed in this way. In particular, in the subsequent chapters of this book, we explicitly construct the soliton solutions of four different nonlinear Schrödinger (NLS) systems. Moreover, one can in principle obtain the long-time asymptotics of solutions. In this book, we obtain formulas for the collision-induced phase shifts of solitons in NLS systems, including the polarization shift of solitons in the vector systems, from the asymptotics of the associated scattering data.

An essential prerequisite of the IST method is the association of the nonlinear evolution equation with a pair of linear problems, a linear eigenvalue problem and a second associated linear problem, such that the given evolution equation results as the compatibility condition between them. The pair of linear operators used to construct the associated linear problems is sometimes referred to as a “Lax pair,” due to a formulation by Lax [115]. The solution of the nonlinear evolution equation appears as a coefficient in the associated linear eigenvalue

problem. For example, in the work of Gardner et al., the solution of the KdV equation is associated with the potential in the linear Schrödinger equation. The eigenvalues and continuous spectrum of this linear eigenvalue problem constitute the transformed variables. The second associated linear problem determines the evolution of the transformed variables.

The associated eigenvalue problem introduces an intermediate stage in both the forward and inverse problems of the IST. In the forward problem, the first step is to construct eigenfunction solutions of the associated linear problem. These eigenfunctions depend on both the original spatial variables and the spectral parameter (eigenvalue). Second, with these eigenfunctions, one determines *scattering data* that are independent of the original spatial variables. In the inverse problem, the first step is the recovery of the eigenfunctions from the (evolved) scattering data. Finally, one recovers the solution in the original variables from these (evolved) eigenfunctions. As noted previously, the evolution of the scattering data is determined by the second associated linear operator and can be computed explicitly.

The properties of the eigenfunctions are key to the formulation of the inverse problem. In general, the solutions of the associated eigenvalue problem also satisfy linear integral equations. For the NLS systems discussed here (as well as other soliton equations in $1 + 1$ dimensions), by using such integral equations one can show that the eigenfunctions are sectionally analytic functions of the spectral parameter. By taking into account the analyticity properties of the eigenfunctions, one can formulate the inverse problem (in particular, the recovery of the eigenfunctions from the scattering data) as a generalized Riemann–Hilbert problem. The Riemann–Hilbert problem is then transformed into a system of linear algebraic–integral equations. Typically in the formulation of the IST, and in particular for the nonlinear Schrödinger systems considered in this book, the scattering data satisfy symmetry relations that are independent of the evolution. These symmetries in the scattering data are essential and must be taken into account in the solution of the inverse problem.

As explained previously, to apply the IST to a nonlinear evolution equation, one must first find a pair of linear operators that can be associated with the nonlinear equation. However, the general method of construction of the Lax pair for a given evolution equation remains an open problem. Nevertheless, for several equations of physical and mathematical interest, such a pair has been found and the IST developed.

In a major step forward, following the works of Gardner et al. [82] and Lax [115], in 1972 Zakharov and Shabat [192] showed that the method also could be applied to another physically significant nonlinear evolution equation, namely,

the nonlinear Schrödinger equation (NLS). Subsequently, Manakov extended this approach to the solution of a pair of coupled NLS equations [120]. In fact, Manakov's work applies equally well to a system of N coupled NLS equations, a system that we refer to as vector NLS (VNLS). Using these ideas, Ablowitz, Kaup, Newell, and Segur developed a method to find a rather wide class of nonlinear evolution equations solvable by this technique [10]. In their work they named the technique the inverse scattering transform (IST). Later, Beals and Coifman analyzed the direct and inverse scattering associated with higher order systems of linear operators [28].

The IST has been extended to semi-discrete nonlinear evolution equations (discrete in space and continuous in time) as well as doubly discrete (discrete in both space and time) systems. Flaschka adapted the IST to solve the Toda lattice equation [79], and Manakov used a similar formulation to solve a nonlinear ladder network [121]. Subsequently, Ablowitz and Ladik developed a method to construct families of semi-discrete and doubly discrete nonlinear systems along with their respective linear operator pairs, as required for the solution of the nonlinear systems via the IST [11, 12] (see also [21]). Included in the formulation of Ablowitz and Ladik are an integrable semi-discretization of NLS (which we refer to as integrable discrete NLS, or IDNLS) as well as a doubly discrete integrable NLS. In Chapter 5 we will further extend the IST method to an integrable semi-discretization of the VNLS that was introduced in [18].

While the work mentioned in the preceding paragraphs consists of applications of the IST method to $1 + 1$ -dimensional evolution equations, since the early 1980s significant progress has also been made in the extension of the IST approach to $2 + 1$ -dimensional systems. For example, the Kadomtsev–Petviashvili equation [100], which is a $2 + 1$ -dimensional generalization of the KdV equation, and the Davey–Stewartson equation [59], which is a natural $2 + 1$ -dimensional integrable extension of the NLS equation, can be solved via the IST. However, the extension of the IST to such systems is beyond the scope of this book. A review of some developments in the application of the IST to $2 + 1$ -dimensional systems can be found in [6].

It should also be noted that IST for periodic and other boundary conditions has been considered, but we will not discuss this here. Additional references can be found in the Bibliography.

1.3 Nonlinear Schrödinger systems

The scalar nonlinear Schrödinger (NLS) equation

$$iq_t = q_{xx} \pm 2|q|^2 q \quad (1.3.1)$$

is a physically and mathematically significant nonlinear evolution equation. It results from the coupled pair of nonlinear evolution equations

$$iq_t = q_{xx} - 2rq^2 \quad (1.3.2a)$$

$$-ir_t = r_{xx} - 2qr^2 \quad (1.3.2b)$$

if we let $r = \mp q^*$.

The NLS equation (1.3.1) arises in a generic situation. It describes the evolution of small amplitude, slowly varying wave packets in nonlinear media [30]. Indeed, it has been derived in such diverse fields as deep water waves [190, 31]; plasma physics [191]; nonlinear optical fibers [91, 92]; magneto-static spin waves [194]; and so on. Mathematically, it attains broad significance because it is integrable by the IST [192], it admits soliton solutions, it has an infinite number of conserved quantities, and so on.

We also note that the form of the NLS equation (1.3.1) with a minus sign in front of the nonlinear term is sometimes referred to as the “defocusing” case. The defocusing NLS equation does not admit soliton solutions that vanish at infinity. However, it does admit soliton solutions that have a nontrivial background intensity (called dark solitons) [92, 193]. We will only discuss the IST for functions decaying sufficiently rapidly at infinity.

The vector nonlinear Schrödinger equation,

$$iq_t^{(1)} = q_{xx}^{(1)} + 2(|q^{(1)}|^2 + |q^{(2)}|^2)q^{(1)} \quad (1.3.3a)$$

$$iq_t^{(2)} = q_{xx}^{(2)} + 2(|q^{(1)}|^2 + |q^{(2)}|^2)q^{(2)}, \quad (1.3.3b)$$

arises, physically, under conditions similar to those described by the NLS when there are two wavetrains moving with nearly the same group velocities [144, 185]. Moreover, VNLS models physical systems in which the field has more than one component; for example, in optical fibers and waveguides, the propagating electric field has two components that are transverse to the direction of propagation. Manakov [120] first examined equation (1.3.3) as an asymptotic model for the propagation of the electric field in a waveguide. Subsequently, this system was derived as a key model for lightwave propagation in optical fibers (cf. [72], [122], [131], [179]).

In the literature, the system (1.3.3) is sometimes referred to as the coupled NLS equation. This system admits vector-soliton solutions, and the soliton collision is elastic. Moreover, the dynamics of soliton interactions can be explicitly computed [120]. (In [142] a different point of view is discussed.) Vector-soliton collisions are analyzed in Sections 4.3 (continuous) and 5.3 (discrete). In these sections, the elasticity of vector-soliton interactions and the order-dependence of these interactions are described in detail.

Both the VNLS equation (1.3.3) and its generalization,

$$i\mathbf{q}_t = \mathbf{q}_{xx} \pm 2 \|\mathbf{q}\|^2 \mathbf{q}, \quad (1.3.4)$$

where \mathbf{q} is an N -component vector and $\|\cdot\|$ is the Euclidean norm, are integrable by the IST. In [120] only the case $N = 2$ is studied, but the extension to more components is straightforward. The N -component equation can be derived, with some additional conditions, as an asymptotic model of the interaction of N wavetrains in a weakly nonlinear, conservative medium (cf. [144]).

In optical fibers and waveguides, depending on the physics of the particular system, the propagation of the electromagnetic waves may be described by variations of equation (1.3.3). Note that the VNLS equation is the ideal (exactly integrable) case. For example, a model with physical significance is [131, 132, 150, 184, 187]

$$iq_t^{(1)} = q_{xx}^{(1)} + 2(|q^{(1)}|^2 + B|q^{(2)}|^2)q^{(1)} \quad (1.3.5a)$$

$$iq_t^{(2)} = q_{xx}^{(2)} + 2(B|q^{(1)}|^2 + |q^{(2)}|^2)q^{(2)}, \quad (1.3.5b)$$

which is equivalent to equation (1.3.3) when $B = 1$. However, based on the properties of equations (1.3.5), apparently it is not integrable when $B \neq 1$ (see the discussion in [18]).

The VNLS (1.3.4) has a natural matrix generalization in the system

$$i\mathbf{Q}_t = \mathbf{Q}_{xx} - 2\mathbf{Q}\mathbf{R}\mathbf{Q} \quad (1.3.6a)$$

$$-i\mathbf{R}_t = \mathbf{R}_{xx} - 2\mathbf{R}\mathbf{Q}\mathbf{R}, \quad (1.3.6b)$$

where \mathbf{Q} and \mathbf{R} are $N \times M$ and $M \times N$ matrices, respectively. When $\mathbf{R} = \mp \mathbf{Q}^H$ (here and in the following, the superscript H denotes the Hermitian, i.e., conjugate transpose), the system (1.3.6a)–(1.3.6b) reduces to the single matrix equation

$$i\mathbf{Q}_t = \mathbf{Q}_{xx} \pm 2\mathbf{Q}\mathbf{Q}^H\mathbf{Q}, \quad (1.3.7)$$

which we refer to as matrix NLS or MNLS. The VNLS corresponds to the special case when \mathbf{Q} is an N -component row vector and \mathbf{R} is an N -component column vector, or vice versa. In particular, we obtain the system (1.3.3) when $M = 1$ and $N = 2$.

Both the NLS and the VNLS equations admit integrable discretizations that, besides being used as the basis for constructing numerical schemes for the continuous counterparts, also have physical applications as discrete systems (see, e.g., Aceves et al. [22, 23]; Braun and Kivshar [40]; Christodoulides and Joseph [54]; Claude et al. [55]; Darmanyan et al. [57, 58]; Davydov [60, 61, 62]; Eilbeck et al. [69]; Eisenberg et al. [70, 71]; Flach et al. [76, 77];