




Six-Membered
Transition States in

ORGANIC SYNTHESIS

Jaemoon Yang

SIX-MEMBERED TRANSITION STATES IN ORGANIC SYNTHESIS

Jaemoon Yang

Montana State University

Department of Chemistry

 **WILEY-
INTERSCIENCE**

A JOHN WILEY & SONS, INC., PUBLICATION

Copyright © 2008 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permission>.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Yang, Jaemoon

Six-membered transition states in organic synthesis / by Jaemoon Yang.

p. cm.

Includes index.

ISBN 978-0-470-17883-6 (cloth)

1. Reaction mechanisms (Chemistry) 2. Stereochemistry 3. Organic compounds—Synthesis. I. Title.

QD502.5.Y36 2008

547'.2—dc22

2007019896

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

**SIX-MEMBERED
TRANSITION STATES IN
ORGANIC SYNTHESIS**

*To my sons,
Walt and Larry*

PREFACE

When I was a graduate student in the Department of Chemistry at the University of Pittsburgh, I took organic chemistry courses taught by Professors Craig S. Wilcox and Dennis P. Curran. One of the most amazing topics that I learned about from their lectures was stereoselective synthesis in organic chemistry.

Stereochemistry is a concept of paramount importance in chemistry. Stereoselective reactions, be they diastereoselective or enantioselective, are therefore a valuable tool in producing compounds of the desired stereochemistry. Every stereoselective reaction has an energetically preferred transition state that can explain the formation of the major stereoisomer. A reasonable transition state is very important not only in rationalizing the experimental results, but also in further advancing the chemical system that one is studying.

Since the seminal proposal in 1957 by Howard E. Zimmerman and Marjorie D. Traxler regarding the stereoselective Ivanov reaction, six-membered chairlike transition states have been recognized as one of the most convincing methods used in organic chemistry to describe the course of reactions that have a well-organized molecular ensemble geometry. In this book I describe organic reactions that go through well-defined six-membered transition states. The reactions are classified into four categories: [3,3]-sigmatropic rearrangements, aldol reactions, metal allylation reactions, and stereoselective reductions. Each chapter begins with a section on general considerations in which I gather all the computational studies known to me that support the proposal of a six-membered transition state. Each reaction has a brief introduction, a description of the six-membered chairlike transition state, and applications selected from natural product synthesis. In presenting reactions and transition states, I have tried to deliver the arguments and conclusions exactly the way they are outlined in the original references. When questions arise or further information on a transition state is sought, readers are strongly encouraged to study the references listed at the end of each section.

This book will serve as a starting point in learning the amazing features of six-membered chairlike transition states in stereoselective organic reactions. With this book, I hope that students and practitioners alike will be able to propose reasonable transition states for the description of newly discovered stereoselective reactions.

Comments and suggestions from readers are always welcome. I can be reached by email at bizibeaver@yahoo.com.

Acknowledgments

I would like to thank the American Chemical Society and Elsevier for their generous permission to use materials for this book. A list of copyrighted materials is included.

Without the many people who have helped me, this book could not have appeared. I would like to thank Professor Tom Livinghouse of Montana State University for his critical reading of the entire manuscript. Drs. Rohan Beckwith, Xing Dai, Tim Peelen, and Janelle Thompson each read portions of the manuscript and provided a number of helpful comments. Thanks are also due to graduate students Elisa Leonardo and Bryce Sundahl at the Livinghouse Laboratory for their invaluable editorial assistance. Thanks also to the anonymous reviewers who read sample chapters and gave me invaluable suggestions. I have one very special person to whom I would like to express my sincere gratitude: Walt Harris, head football coach at the University of Pittsburgh between 1996 and 2005. Not only has coach Harris been a constant source of inspiration to me, he has also been a big part of my family's life, and I truly thank him for that.

Finally, I would like to thank my wife, Wenjing Xu, for her criticisms, encouragement, and suggestions over the course of writing the book.

JAEMOON YANG

Bozeman, Montana

CONTENTS


Preface	ix
Introduction	1
1 [3,3]-Sigmatropic Rearrangements	5
General Considerations, 5	
Reactions, 13	
1.1 Claisen Rearrangement, 13	
1.2 Johnson–Claisen Rearrangement, 20	
1.3 Ireland–Claisen Rearrangement, 27	
1.4 Cope Rearrangement, 32	
1.5 Anionic Oxy-Cope Rearrangement, 36	
1.6 Aza-Cope–Mannich Reaction, 43	
2 Aldol Reactions	49
General Considerations, 49	
Reactions, 57	
2.1 Asymmetric <i>Syn</i> -Aldol Reaction, 57	
2.2 Asymmetric <i>Anti</i> -Aldol Reaction, 78	
2.3 Proline-Catalyzed Asymmetric Aldol Reaction, 91	
3 Metal Allylation Reactions	97
General Considerations, 97	
Reactions, 102	
3.1 Boron Allylation Reaction, 102	
3.2 Silicon Allylation Reaction, 127	
4 Stereoselective Reductions	147
General Considerations, 147	
Reactions, 151	
4.1 Diastereoselective <i>Syn</i> -Reduction of β -Hydroxy Ketones, 151	

- 4.2 Diastereoselective *Anti*-Reduction of β -Hydroxy Ketones, 161
- 4.3 Asymmetric Reduction, 173


List of Copyrighted Materials	197
Abbreviations	199
Subject Index	201
Scheme Index of Natural Products	209

INTRODUCTION

In 1957, Zimmerman and Traxler published their study on the reaction of benzaldehyde with the magnesium enolate of phenylacetic acid: namely, the *Ivanov reaction*¹ (Scheme I). The major product from the reaction is an *anti*- or *threo*-isomer of 2,3-diphenyl-3-hydroxypropionic acid, and the minor product is a *syn*- or *erythro*-isomer.² Although in 1957 the Ivanov reaction had been known for quite some time, no reasonable proposal had been put forward to explain the stereochemical outcome observed for the reaction. In explaining the ratio of the two stereoisomers, the authors made a seminal proposal that the condensation reaction would go through a six-membered transition state (Scheme II). The coordination of benzaldehyde carbonyl group with magnesium brings the two reactants in close contact in both chairlike transition state **A** and boatlike transition state **B**. The authors speculated that the particular spatial arrangement of the four substituents in the transition state could determine the stereochemistry of the products. For the Ivanov reaction of benzaldehyde, transition state **A** would be favored over **B** because transition state **A** involves a lower-energy approach to bonding than that of the alternative transition state **B**, which experiences an energetically unfavorable gauche interaction between the two phenyl substituents¹ (Scheme III).




Scheme I



TS A: chairlike transition structure

TS B: boatlike transition structure


Scheme II



Scheme III

Due to its simplicity and outstanding prediction power, the *Zimmerman–Traxler transition state* has frequently been used in explaining the stereochemical outcome of certain stereoselective reactions. The characteristics of the *Zimmerman–Traxler transition state* can be summarized as follows:

1. The transition state is for a six-atom system and thus is six-membered.
2. The transition state involves six electrons and thus exhibits the aromatic character of benzene.³
3. A chairlike transition state is favored over a boatlike transition state. There are, however, exceptions.



Scheme IV

TABLE 1 Conformational Energies of Monosubstituted Cyclohexanes

R	$-\Delta G$ (kcal/mol)	EQ/AX
Me	1.74	19:1
Et	1.79	19:1
<i>i</i> -Pr	2.21	42:1
<i>t</i> -Bu	4.7	>99:1
C ₆ H ₅	2.8	>99:1
OMe	0.55	2.5:1

4. When two chairlike transition states compete, the transition state in which a bulky substituent occupies an equatorial position is favored over the state that has the same substituent in an axial position. The free-energy difference between the two potential transition states can be approximated by using the ΔG or A-values of the monosubstituted cyclohexanes⁴ (Table 1).

In the following four chapters, readers will find some of the most frequently cited and most synthetically relevant examples of the Zimmerman–Traxler or six-membered transition state. In presenting reactions that go through a six-membered chairlike transition state, I pay special attention to including computational studies, in an effort to prove the existence of a six-membered chairlike transition state. Although not all six-membered transition states have been studied computationally, recent interest in using computers in studies of stereoselective reactions would certainly confirm the legitimacy of Zimmerman–Traxler transition states for many more reactions.⁵

Before we embark on our journey into the world of six-membered transition states, I would like to speak briefly about one reaction, to illustrate how a transition state is drawn throughout the book. The enzyme-catalyzed transformation of chorsimate (**2**) to prephenate (**3**) is a classic example of a [3,3]-sigmatropic *Claisen rearrangement*⁶ (Scheme IV). As an old bond is being broken and at the same time a new bond is formed in the transition state, the transition state for the Claisen rearrangement of chorismate to prephenate would look more like transition state **A** than like **B**. Still, for the convenience of following the bond connection event clearly, I prefer to draw the transition state like **B**.

REFERENCES

1. Zimmerman, H. E.; Traxler, M. D. *J. Am. Chem. Soc.* **1957**, 79, 1920.
2. For definitions of *syn* and *anti*, see Masamune, S.; Ali, S. A.; Snitman, D. L.; Garvey, D. S. *Angew. Chem. Int. Ed.* **1980**, 19, 557.
3. (a) Day, A. C. *J. Am. Chem. Soc.* **1975**, 97, 2431; (b) Carey, F. A.; Sundberg, R. J., *Advanced Organic Chemistry*, Part A, 3rd ed.; Plenum Press: New York, **1990**; Chap. 11.
4. Eliel, E. L.; Wilen, S. H.; Mander, L. N. *Stereochemistry of Organic Compounds*; Wiley: New York, **1994**; Chap. 11.
5. Lipkowitz, K. B.; Kozlowski, M. C. *Synlett* **2003**, 1547.
6. (a) Andrews, P. R.; Haddon, R. C. *Aust. J. Chem.* **1979**, 32, 1921; (b) Copley, S. D.; Knowles, J. R. *J. Am. Chem. Soc.* **1985**, 107, 5306.

1 [3,3]-Sigmatropic Rearrangements


GENERAL CONSIDERATIONS

The Claisen and Cope rearrangements are two of the best known sigmatropic rearrangements in organic chemistry¹ (Scheme 1.I). As the rearrangement involves six electrons in a six-atom system, these two reactions serve as excellent examples of the ubiquitous existence of a six-membered transition state in organic chemistry.


In 1912, Ludwig Claisen discovered that the allyl ether **1** of ethyl acetoacetate underwent a reaction to afford **2** upon heating in the presence of ammonium chloride² (Scheme 1.II). Similarly, the allyl naphthyl ether **3** transformed into 1-allyl-2-naphthol (**4**) in 82% yield at 210 °C. The reaction, now known as the *Claisen rearrangement*, is general for a variety of aliphatic and aromatic ethers and is recognized as one of the most synthetically useful reactions in organic chemistry.³

The Claisen rearrangement is a thermally induced [3,3]-sigmatropic rearrangement of allyl vinyl ethers to form γ,δ -unsaturated carbonyl compounds.⁴ Due to the concerted nature and synthetic utilities of the Claisen rearrangement, much effort has been devoted to understanding the mechanism of the reaction.⁵ Although the extent of delocalization of the six electrons involved in the transition state may depend on the nature of the substrates, it is believed that the rearrangement goes through a six-membered aromatic transition state⁶ (Scheme 1.III).


To uncover the transition-state structures for Claisen rearrangement of the parent allyl vinyl ether,⁷ Vance et al. performed *ab initio* quantum mechanical calculations⁸ (Scheme 1.IV). When the transition structures were calculated using the 6-31G* basis set, the partially formed C₁–C₆ bond length is 2.26 Å and the partially broken C₄–O bond length is 1.92 Å in chairlike transition structure **A**. These two bond lengths were confirmed by Meyer et al. in a later study employing different-level calculations.⁹ Another important finding in Vance et al.'s study is that chairlike transition structure **A** is more stable than boatlike structure **B**, by 6.6 kcal/mol. The conclusion thus supports the proposals of chairlike transition structures for the stereoselectivities observed for the Claisen rearrangement reactions of substituted molecules.




Scheme 1.I



Scheme 1.II




Scheme 1.III




Scheme 1.IV

One classic example that confirms the preference of Claisen rearrangement for a chairlike transition state was provided by Hansen and others. In 1968, they investigated the Claisen rearrangement of the crotyl propenyl ethers **5a** and **5b** to examine the stereochemistry of the rearrangement in the gas phase at 160°C¹⁰ (Scheme 1.V). Both the *E,E*- and *Z,Z*-isomers rearrange to afford a *syn*-isomer as the major product. The stereochemical outcome of the reaction can be explained




Scheme 1.V




Scheme 1.VI

in terms of a six-membered transition state¹⁰ (Scheme 1.VI). Between the two transition states for **5a**, chairlike transition state **A** is favored over boatlike state **B** to afford a *syn*-isomer as the major product. Chairlike transition state **C** can explain the formation of the *syn*-isomer that is enantiomeric to **6-syn**. Other indirect evidence for the existence of a chairlike transition state is the fact that the *E,E*-isomer **5a** reacts nine times faster than the *Z,Z*-isomer **5b**. This difference in the reaction rate can be understood by examining transition states **A** and **C**: Transition state **C** for **5b** is of higher energy than transition state **A** for **5a**, due presumably to the 1,3-diaxial interactions arising from the axial methyl groups in transition state **C**.



<i>7E/7Z</i>	8a/8b	Yield, %
83:17	84:16	79
4:96	72:28	91

Scheme 1.VII



Scheme 1.VIII