RATERAL PRODUCTS

PT.1



natural products isolation

separation methods for antimicrobials, antivirals and enzyme inhibitors

edited by

Gerald H. Wagman and Raymond Cooper

Schering-Plough Research, 60 Orange Street, Bloomfield, NJ 07003, U.S.A.



ELSEVIER SCIENCE PUBLISHERS B.V. Sara Burgerhartstraat 25 P.O. Box 211, 1000 AE Amsterdam, The Netherlands

Distributors for the United States and Canada:

ELSEVIER SCIENCE PUBLISHING COMPANY INC. 655 Avenue of the Americas New York, NY 10010, U.S.A.

ISBN 0-444-87147-0 (Vol. 43) ISBN 0-444-41616-1 (Series)

© Elsevier Science Publishers B.V., 1989

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V./ Physical Sciences & Engineering Division, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Special regulations for readers in the USA – This publication has been registered with the Copyright Clearance Center Inc. (CCC), Salem, Massachusetts. Information can be obtained from the CCC about conditions under which photocopies of parts of this publication may be made in the USA. All other copyright questions, including photocopying outside of the USA, should be referred to the publisher.

No responsibility is assumed by the Publisher for any injury and/or damage to persons of property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Printed in The Netherlands

LIST OF CONTRIBUTORS

- TADASHI ARAI Biotherapy Research Association, Tokyo, Japan
- JOHN H. CARDELLINA II Natural Products Laboratory, Department of Chemistry, Montana State University, Bozeman, Montana, USA
- RAYMOND COOPER Microbial Products Research, Schering Corporation, Bloomfield, New Jersey, USA
- LOUISE W. CRANDALL Fermentation Products Research, Lilly Research Laboratories, Indianapolis, Indiana, USA
- LINDA E. FELLOWS Jodrell Laboratory, Royal Botanic Gardens, Kew, England, UK
- HANS-PETER FIEDLER Institute of Biology II, Department of Microbiology I, University of Tubingen, Tubingen, Federal Republic of Germany
- GEORGE W. J. FLEET Dyson Perrins Laboratory, University of Oxford, Oxford, England, UK
- VINCENT P. GULLO Microbial Products Research, Schering Corporation, Bloomfield, New Jersey, USA
- SETSUO HARADA Central Research Division, Takeda Chemical Industries, Ltd., Osaka, Japan
- JILL E. HOCHLOWSKI, Antiinfective Research, Abbott Laboratories, Abbott Park, Illinois, USA
- DAVID G. LYNN Department of Chemistry, University of Chicago, Chicago, Illinois, USA
- JOSEPH A. MARQUEZ Microbial Products Research, Schering Corporation, Bloomfield, New Jersey, USA
- JAMES B. MCALPINE Antiinfective Research, Abbott Laboratories, Abbott Park, Illinois, USA
- RONALD MIERZWA Microbial Products Research, Schering Corporation, Bloomfield, New Jersey, USA
- THOMAS MILLER Merck Sharp and Dohme Research Laboratories, Rahway, New Jersey, USA
- JON S. MYNDERSE Fermentation Products Research, Lilly Research Laboratories, Indianapolis, Indiana, USA
- JOHN D. ORR Department of Chemistry, University of Chicago, Chicago, Illinois, USA
- WILLIAM L. PARKER The Squibb Institute for Medical Research, Princeton, New Jersey, USA

- MAHESH PATEL Microbial Products Research, Schering Corporation, Bloomfield, New Jersey, USA
- SIDNEY PESTKA Department of Molecular Genetics and Microbiology, University of Medicine and Dentistry of New Jersey, Robert Wood Johnson Medical School, Piscataway, New Jersey, USA
- ROBERT D. SITRIN Merck Sharp and Dohme Research Laboratories, West Point, Pennsylvania, USA
- CHRISTOPHER E. SMITH Department of Chemistry, University of Chicago, Chicago, Illinois, USA
- HAMAO UMEZAWA (dec.) Institute of Microbial Chemistry, Tokyo, Japan
- GAIL FOLENA WASSERMAN Smith Kline and French Laboratories, Philadelphia, Pennsylvania, USA
- KENNETH E. WILSON Merck Sharp and Dohme Research Laboratories, Rahway, New Jersey, USA

FOREWORD

A text that provides a compilation of an active area of research should be authored by individuals who themselves have made significant contributions to that field. It is within this aegis that the present authors have edited this latest addition to the library of useful natural products publications.

In earlier publications in this series, two volumes have appeared which were the efforts of investigators who had made important contributions to antibiotic discoveries. These were "Chromatography of Antibiotics" in two editions, 1973 and 1978, and "Antibiotics; Isolation, Separation, and Purification", 1978. One of the authors, Gerald Wagman, has joined with Raymond Cooper to carry on this tradition of erudition.

Microbiology has continued its march into other prolific fields of exploration. Microorganisms display an impressive biosynthetic ingenuity, and have become a useful source of enzyme inhibitors, active peptides, and immunomodulators. This present volume has addressed these new developments. The organisms have not changed, but the techniques for the identification of their products have become more sensitive and selective. This volume, which compiles the experience of the experts in their fields, has allowed for the dissemination of these changing techniques. Since important bioactive compounds are also being found from terrestrial and marine sources, it is appropriate to expand the scope of this book to include them.

My perspective of the field is now viewed through a retired eye, but my vision is clear that this volume will be a valued addition to the continuing generations of microbial screeners and natural products chemists.

Marvin J. Weinstein, Ph.D.
Vice-President, Microbiology
Schering Corporation (Retired)

PREFACE

Progress in the discovery of natural products requires a multidisciplinary approach involving the expertise of scientists in such fields - as biology, botany, molecular biology, chemistry and pharmacology. Although any given project may be interdisciplinary in nature, the contributions of the natural products chemist to the isolation of a particular compound are unique. Using modern technology it has often been possible to purify and identify a new and potentially useful compound on a scale so small that several years ago was considered impossible. The advances in the biological, medicinal and pharmaceutical sciences have been maintained in no small part due to improvements in chromatographic methods and separation techniques. In the cases of isolation of small quantities of bioactive material, the challenge of structure determination has been complemented by the ever impressive sophistication of spectroscopic methods and instrumentation. The scope of this book is to encompass, in as much detail as possible, the most recent progress made in the isolation and separation of compounds of natural origin. This book is an outgrowth of the previous book in this field (Antibiotics, Eds. M. Weinstein and G. Wagman) but is expanded to cover entirely new material and other natural products. We have attempted this project by bringing together 14 diverse chapters aimed at presenting modern methods of isolating antibiotics, marine and plant derived substances, enzyme inhibitors and interferons. The chosen topics are organized to present modern methodology for separation of antimicrobials, antivirals and enzyme inhibitors from a variety of sources. These chapters cover families or groups of compounds that have gained prominence in the last ten years, e.g. avermectins, interferons.

The first two chapters are devoted to newer methodologies applied to purification of a variety of compounds. There is an extensive review on applications of counter-current chromatography. We are witnessing a renaissance in liquid chromatography due to some innovative technology, and a review of several different techniques is presented. The applications of the emerging HPLC - photodiode array technology are reviewed in Chapters 2 and 4. It should be noted in the previous volume that aminoglycosides, tetracyclines and macrolides were reviewed; in this volume HPLC methods development, detection and screening of these compounds are presented

reflecting the advancement and the present interest in these families of compounds within the pharmaceutical industry. Although affinity chromatography is not a new approach, a review of the first application to the separation of antibiotics directly from fermentation broth is given in Chapter 3. Using this approach, a large family of glycopeptides has been isolated.

There are several selected topics indicating the isolation of antimicrobials. These include the family of antitumor compounds known as the saframycins and a comprehensive review of a very important new class of antiparasitic agent, the avermectins. An important source for antimicrobials and other bioactive natural products has been from the sea and an update of isolation and purification of a variety of marine—derived compounds is given in Chapter 10. The prominence and importance of β —lactams in the field of antimicrobial chemistry cannot be underestimated. For this reason we have included three chapters indicating the major advances over the past 10 years. The monobactams and thienamycins are extremely unstable natural products produced in culture broth. The successful discovery and isolation of small quantities have led to their subsequent development as new β -lactam antibiotics in the market place.

Today, directed screens for enzyme inhibitors is an integral part of a natural products program. Isolation and purification of these natural products from fermentation and plant sources are included. The chapter on enzyme inhibitors was written by Professor Hamao Umezawa shortly before his untimely death. Dr. Umezawa's son, Dr. Kazuo Umezawa, has kindly given of his time to put the manuscript in final form suitable for publication. We wish to express our sincere appreciation for his efforts in completing this work.

The chapter which follows shows that a relationship exists between plant-derived enzyme inhibitors and those derived from microorganisms. In discussing alkaloidal glycosidase inhibitors, this work reflects the interest of the plant chemist in the isolation of water soluble compounds. The final chapter is a futuristic essay indicating the challenges of the plant chemist to isolate minute amounts of natural products. By discovering these bioactive principles it may open the door to studying the fascinating biological properties which they possess.

We have chosen to include in this volume a chapter devoted to the isolation of interferons in recognition of the impact of molecular biology in the field of natural products and also in part as a reflection of the direction microbial products may take in the years to come.

We wish to thank Drs. Vincent Gullo, George Miller and Claude H. Nash of Schering Research for their encouragement and help in making this book possible. We also wish to acknowledge the excellent cooperation of Dawn Foster, for her assistance in the typing. Lastly, and most important, this volume would not have been possible without the superb cooperation we received from all of the chapter authors. It was a real pleasure to have worked with them and we appreciate their patience with us!

Gerald H. Wagman Raymond Cooper

CONTENTS

List of contributorsvii
Forewordix
Preface xi
Countercurrent Chromatographyl James B. McAlpine and Jill E. Hochlowski
HPLC Detection Methods for Microbial Products from Fermentation Broth
Affinity and HPLC Purification of Glycopeptide Antibiotics111 Robert D. Sitrin and Gail Folena Wasserman
Nikkomycins and Polyoxins153 Hans-Peter Fiedler
Saframycins and Isoquinolines191 Tadashi Arai
New Cepnalosporins233 Setsuo Harada
Monocyclic Beta-Lactam Antibiotics259 William L. Parker
Isolation of Carbapenems293 Kenneth E. Wilson
Avermectins and Related Compounds
Bioactive Compounds from Marine Organisms and Cultivated Blue- Green Algae
The Interferons
Enzyme Inhibitors Produced by Microorganisms481 Hamao Umezawa
Alkaloidal Glycosidase Inhibitors from Plants
Chemical Communication and Control of Development

Abbrevia	ations	
Subject	Index	603
List of	Numbered Compounds	615
Journal	of Chromatography Library (o	ther volumes in the series).617

Chapter 1

COUNTERCURRENT CHROMATOGRAPHY

J.B.McALPINE and J.E.HOCHLOWSKI

Antiinfective Research, Abbott Laboratories, Abbott Park, Illinois 60064

1	INTRODU	JCTION	3
	1.1 His	storical Development	3
	1.2 Bas	sic Theory	4
	1.3 Scc	ope and Limitations	5
2	INSTRUM	IENTATION	7
	2.1 Dr	oplet Countercurrent Chromatography (DCCC)	7
	2.2 Ro	tary Locular Countercurrent Chromatography (RLCC)	8
	2.3 Sp	inning Column Countercurrent Chromatography (SCCC)	9
	2.4 Ce	ntrifugal Countercurrent Chromatography with a	
	Ro	otating Seal Joint (CCCC-RJ)	10
	2.5 Co	il Planet Centrifuge Countercurrent Chromatography (CPC)	11
	2.5.1	The Horizontal Synchronized Multilayered Coil	1
	2.5.2	Other Configurations	15
3	PRACTIC	CAL CONSIDERATIONS	17
	3.1 Ch	noosing a Solvent System	17
	3.2 Pu	imping Systems and Sample Application	19
		uent Monitoring Systems	
4	SPECIFIC	C EXAMPLES	21
	4.1 Dt	roplet Countercurrent Chromatography Examples	21
	4.1.1	Resolution of Enantiomers	21
	4.1.2	Terpenoids from Matricaria chamomilla	22
	4.1.3	Separation of Iridoid Glycosides	23
	4.1.4	Isolation of a Phenolic Glycoside	24
	4.2 Ro	otary Locular Countercurrent Chromatography Examples	24
	4.2.1	Resolution of Norephedrine	24
	4.2.2	Isolation of a Phenylpropanoid Plant Growth Regulator	24
(6	4.2.3	Isolation of Flavanoid C-glycosides	25
	4.2.4	Isolation of Raucaffricine	26
	4.3 Sp	pinning Column Countercurrent Chromatography Examples	26
	4.3.1	Separation of Glucose Derivatives	26
	4.3.2	Separation of Neutral Macrolide Antibiotics	27

	4.3.	3 Purification of Mocimycin	7
	4.3.	4 Purification of Chartreusin	3
	4,4	Examples of the Use of Centrifugal Countercurrent Chromatography	
		with Rotating Seal Joint	3
	4.4.	1 Separation of Retinals	3
	4.4.	2 Separation of Sugars	9
	4.4.	3 Separation of Fatty Acids	9
	4.5	Examples of Use of the Coil Planet Centrifuge)
	4.5.	1 Sesquiterpenes from Artemisa annua	C
	4.5.	2 Sesquiterpene Antibiotics - Pentalenolactone	C
	4.5.	3 Phenoxazinepeptolides - The Actinomycins	1
	4.5.	4 Tetramic Acids - The Tirandamycins	2
	4.5.	5 Macrolide Antibiotics - 2-Norerythromycins	2
	4.5.	6 Macrolide Antibiotics - Niddamycins	2
	4.5.	7 Quinoxaline Antibiotics	4
	4.5.	8 Pyrroles	5
	4.5.	9 Use as an Extraction Device	5
5	TABLE	ES	7
	5.1	Droplet Countercurrent Systems	7
	5.2	Rotary Locular Systems4	
	5.3	Spinning Column Systems	1
	5.4	Systems Used With the Centrifugal Countercurrent with Rotating Seal Joint 4	2
	5.5	Coil Planet Centrifuge Systems 4	3
6	VEND	ORS4	8
7	REFER	RENCES4	9

Chapter 1

COUNTERCURRENT CHROMATOGRAPHY

J.B.McALPINE and J.E.HOCHLOWSKI

Antiinfective Research, Abbott Laboratories, Abbott Park, Illinois 60064

1	INTROD	UCTION	3
	1.1 Hi	storical Development	3
	1.2 Ba	sic Theory	4
	1.3 Sc	ope and Limitations	5
2	INSTRUM	MENTATION	7
	2.1 Dr	roplet Countercurrent Chromatography (DCCC)	7
	2.2 Ro	otary Locular Countercurrent Chromatography (RLCC)	8
	2.3 Sp	ninning Column Countercurrent Chromatography (SCCC)	9
	2.4 Ce	entrifugal Countercurrent Chromatography with a	
	R	otating Seal Joint (CCCC-RJ)	10
	2.5 Co	oil Planet Centrifuge Countercurrent Chromatography (CPC)	1
	2.5.1	The Horizontal Synchronized Multilayered Coil	11
	2.5.2	Other Configurations	15
3	PRACTIC	CAL CONSIDERATIONS	17
	3.1 Ch	hoosing a Solvent System	17
	3.2 Pu	imping Systems and Sample Application	19
	3.3 El	uent Monitoring Systems	19
4	SPECIFIC	C EXAMPLES	21
	4.1 Dr	roplet Countercurrent Chromatography Examples	21
	4.1.1	Resolution of Enantiomers	21
	4.1.2	Terpenoids from Matricaria chamomilla.	22
	4.1.3	Separation of Iridoid Glycosides	23
	4.1.4	Isolation of a Phenolic Glycoside	24
	4.2 R	otary Locular Countercurrent Chromatography Examples	24
	4.2.1	Resolution of Norephedrine	24
	4.2.2	Isolation of a Phenylpropanoid Plant Growth Regulator	24
	4.2.3	Isolation of Flavanoid C-glycosides	25
	4.2.4	Isolation of Raucaffricine	26
	4.3 S ₁	pinning Column Countercurrent Chromatography Examples	26
	4.3.1	Separation of Glucose Derivatives	26
	4.3.2	Separation of Neutral Macrolide Antibiotics	27

	4.3.3	Purification of Mocimycin	27
	4.3.4	Purification of Chartreusin	
	4.4 E	xamples of the Use of Centrifugal Countercurrent Chromatography	
		with Rotating Seal Joint	
	4.4.1	Separation of Retinals	28
	4.4.2	Separation of Sugars	29
	4.4.3	Separation of Fatty Acids	29
	4.5 E	xamples of Use of the Coil Planet Centrifuge	30
	4.5.1	Sesquiterpenes from Artemisa annua	30
	4.5.2	Sesquiterpene Antibiotics - Pentalenolactone	30
	4.5.3	Phenoxazinepeptolides - The Actinomycins	31
	4.5.4	Tetramic Acids - The Tirandamycins	32
	4.5.5	Macrolide Antibiotics - 2-Norerythromycins	32
	4.5.6	Macrolide Antibiotics - Niddamycins	32
	4.5.7	Quinoxaline Antibiotics	34
	4.5.8	Рупоеѕ	35
	4.5.9	Use as an Extraction Device	35
5	TABLES		37
	5.1 D	roplet Countercurrent Systems	37
	5.2 R	otary Locular Systems	40
	5.3 S	pinning Column Systems	41
	5.4 S	ystems Used With the Centrifugal Countercurrent with Rotating Seal Joint	42
	5.5 C	oil Planet Centrifuge Systems	43
6	VENDO	RS	48
7	REFERE	NCES	49

1. INTRODUCTION.

Countercurrent chromatography is a liquid-liquid partition chromatography technique employing two immiscible liquid phases. One phase is designated as the stationary phase and is retained as such by either gravitational or centrifugal forces while the second phase, the mobile phase, is passed through the stationary phase. Separation of the components of a mixture is achieved by introduction of the sample at one end of the stationary phase with the mobile phase. Each component is then retained by the stationary phase to a degree dependent on its partition coefficient for the two-phase system. Individual components are eluted from the other end of the stationary phase by the mobile phase. That component with the partition coefficient most in favor of the mobile phase will be eluted first.

As a separation technique, countercurrent chromatography has several inherent advantages. The number of two-phase systems available is virtually limitless and in order to achieve a separation between any two compounds all that is required is that a two-phase system be found in which the two compounds have significantly different partition coefficients. Loss of sample by irreversible absorption or chemical degradation, a common problem in chromatographic methods involving a solid support, is seldom experienced during countercurrent chromatography. Several of the more modern instruments give rapid chromatographies and, with the appropriate solvent systems, can provide the mildest possible conditions for a chromatographic separation.

Varieties of new instruments have recently been developed or are currently enjoying rapid advances and these offer the chemist high resolution separation techniques which may be tailored to any particular purification problem. Ito and Conway have written several detailed chapters on the design and theory behind the many different instruments designed by the former. ¹⁻⁹ The reader is directed to these writings for greater appreciation of the historical development of current instruments and a more thorough understanding of the theoretical aspects. The latter is dealt with here only to that extent required for optimal utilization of the technique. Hostettmann and Hostettmann have provided many practical reviews with emphasis on the separation of plant products mainly with Droplet Countercurrent systems. ¹⁰⁻¹⁵ Elsewhere in this volume Mynderse *et al.* give a detailed description of the use of the Centrifugal Countercurrent with Rotating Joint Seal for the separation and purification of tunichromes.

Although most of the examples discussed herein have been drawn from the antibiotic area, we have included examples from other areas where these have involved (i) the use of unusual solvent systems or (ii) compounds of uncommon polarity or (iii) where the number of published examples of use of a specific instrument was particularly limited. Anticipating that the chemist will find the tables most useful we have endeavored to include a wide a variety of solvent systems and to cover as many different classes of natural products as possible.

1.1 Historical Development.

Countercurrent chromatography was originally developed in the early 1950's with much of the pioneering work done by Lyman Craig. This work is well documented 16 and culminated in the

design and commercialization of multi-tube glass countercurrent distribution instruments which were commonly referred to as "Craig machines". The high resolution capabilities inherent in the method surpassed anything available from other separation techniques at the time. The instruments, however, were large, cumbersome, and fragile and required dedicated venting systems to remove organic solvent vapors. Moreover a typical separation, including set-up and instrument cleaning would encompass a week's work. None-the-less, the technique was used to achieve difficult, or otherwise impossible separations. Gross ¹⁷ for example, separated individual components of the Gramicidin complex, a family of pentadecapeptide antibiotics the individual members of which in some instances differed by a single residue change of valine to isoleucine.

With the advent of high resolution solid-phase support chromatography systems, especially HPLC, that allowed for rapid and convenient separations, countercurrent distribution fell into relative disuse. The design and commercialization, within the last ten years, however, of several new continuous-flow countercurrent chromatography instruments has provided a resurgence of interest in this separation methodology.

1.2 Basic Theory.

For the Craig machine, where each cell is a discrete unit and equilibrium distribution is achieved between each transfer, the mathematics of distribution of a component of a given partition coefficient for a set number of transfers is relatively simple. Some examples of these are shown in Figure 1-1. Given the volumes of upper and lower phase in each cell and the partition coefficient of a particular component of a mixture, the number of transfers required to elute that component of an be readily calculated. These calculations assume that the partition coefficient of one




Figure 1-1 Theoretical distribution curves of solutes of various Partition Coefficients from 0.1 to 10 in the tubes of a Craig machine after 500 transfers with equal volumes of each phase