

**David Rubenstein
David Wayne
Lecture Notes on
Clinical Medicine**

Third edition

04183
R 5
E1703 R895 No. 3

Lecture Notes on Clinical Medicine

DAVID RUBENSTEIN

M.A. M.D. F.R.C.P. (Lond.)

*Physician, Addenbrooke's Hospital,
Cambridge*

DAVID WAYNE

M.A. B.M. F.R.C.P. (Lond. & Edin.)

*Physician, James Paget Hospital,
Gorleston, Great Yarmouth*

THIRD EDITION



BLACKWELL SCIENTIFIC PUBLICATIONS

OXFORD LONDON

EDINBURGH BOSTON MELBOURNE

© 1976, 1980, 1985 by
Blackwell Scientific Publications
Editorial offices:
Osney Mead, Oxford OX2 OEL
8 John Street, London WC1N 2ES
9 Forrest Road, Edinburgh EH1 2QH
52 Beacon Street, Boston,
Massachusetts 02108, USA
706 Cowper Street, Palo Alto
California 94301, USA
99 Barry Street, Carlton
Victoria 3053, Australia

All rights reserved. No part of this
publication may be reproduced,
stored in a retrieval system, or transmitted,
in any form or by any means,
electronic, mechanical, photocopying,
recording or otherwise
without the prior permission of
the copyright owner

First published 1976
Reprinted 1977, 1978, 1979
Second edition 1980
Reprinted, 1981, 1982, 1983, 1984
Third edition 1985

Portuguese translation 1980

Set in Lasercomp Photina

Printed and bound in Great Britain
at the Alden Press, Oxford

DISTRIBUTORS

USA

Blackwell Mosby Book Distributors
11830 Westline Industrial Drive
St Louis, Missouri 63141

Canada

Blackwell Mosby Book Distributors
120 Melford Drive
Scarborough, Ontario M1B 2X4

Australia

Blackwell Scientific Book Distributors
214 Berkeley Street, Carlton
Victoria 3053

British Library

Cataloguing in Publication Data

Rubenstein, David

Lecture notes on clinical medicine.—3rd ed.

1. Pathology 2. Medicine

I. Title II. Wayne, David

616 RB111

ISBN 0-632-01254-4

Contents

ix	Preface to the third edition
xi	Preface to the first edition
PART 1	THE CLINICAL APPROACH
Nervous system	Cranial nerves: field defects, blindness, eye movements, squint, pupillary reflexes, Horner's syndrome, hearing, dizziness and giddiness, vertigo, Menière's disease, nystagmus, facial palsy, aetiology of cranial nerve palsies, speech disorders, bulbar and pseudobulbar palsies. Tests of higher cerebral functions. Arms: motor system, reflexes, sensory system, coordination. Legs: motor system, knee and ankle reflexes, plantar response, sensory system, coordination. Abnormality of gait. Notes: sensory testing, patterns of sensory loss in limbs, motor testing, patterns of motor loss in limbs, isolated peripheral nerve lesions, incoordination, tremor
3	
Eyes	Optic fundus: myelinated nerve fibres, diabetic fundus, hypertensive fundus, papilloedema, optic atrophy, exudates and haemorrhages. Uveitis, iritis and choroiditis
29	
Limbs	Legs. Arms: rheumatoid arthritis, osteoarthritis, gout
32	
Head and neck	Head. Neck: cervical lymphadenopathy, enlarged thyroid
34	
Abdomen	Examination. Notes: splenomegaly, hepatomegaly, hepatosplenomegaly, palpable kidneys, mass in right subcostal region, ascites. Jaundice: haematology, urine analysis, liver function tests, mitochondrial antibodies X-ray and ultrasound of abdomen, needle liver biopsy. Congenital non-haemolytic hyperbilirubinaemias. Uraemia. Dysphagia. Diarrhoea
37	
Respiratory system	Examination. Notes: clubbing, haemoptysis, pleural effusion, cyanosis. Blood gases: interpretation. Ventilatory function tests
50	
Cardiovascular system	Arterial pulse: special pulses. Neck veins: raised jugular venous pressure. Heart: difficult systolic murmurs, continuous murmurs, heart failure. Hypertension. Myocardial infarct. Cardiac arrhythmias: sick sinus syndrome, torsade de pointes. Electrocardiograms: interpretation, sample patterns, mean frontal QRS axis, hemiblocks, fascicular blocks.
57	

Echocardiography: mitral valve disease, aortic valve disease, cardiomyopathy. Cardiac pressure studies. Cardiac radionuclide studies

Haematology 84 Anaemia. Blood: notes. Skin haemorrhage. Coagulation defects. Lymphadenopathy

Diabetes 94 Management: diet, drugs, assessment of diabetic control insulin resistance, symptoms of poor control. Examination. Diabetes and pregnancy. Diabetes and surgery

Skin History, examination

99

PART 2 ESSENTIAL BACKGROUND INFORMATION

Neurology 103 Headache. Unilateral facial pain: migraine, cluster headaches, neuralgias, temporal arteritis, atypical facial pain, unilateral pain. Cerebrovascular disease: 'strokes', transient ischaemic attacks, lateral medullary syndrome, extracerebral haemorrhage, brain death. Epilepsy. Meningitis. Intracranial abscess. Multiple sclerosis. Parkinson's disease and extrapyramidal disorders; Parkinsonism, hepatolenticular degeneration, chorea, Cerebral tumours: symptoms and signs, acoustic neuroma, benign intracranial hypertension. Motor neurone disease. Peripheral neuropathy. Neuralgic amyotrophy. Mononeuritis multiplex. Hereditary ataxias. Syphilis of the nervous system. Acute post-infective polyneuropathy (Guillain-Barré syndrome). Poliomyelitis. Disorders of spinal cord: syringomyelia, syringobulbia, subacute combined degeneration of the cord, peroneal muscular atrophy, cord compression, cervical spondylosis. Disorders of muscles. Polymyositis and allied disorders. Myasthenia gravis. Myotonia. Muscular dystrophies. Psychiatry. Depression. Mania. Anxiety. Hysteria. Acute confusional states. Dementia. Schizophrenia

Connective tissue and rheumatic diseases 146 Systemic lupus erythematosus. Polyarteritis nodosa. Dermatomyositis. Polymyositis. Scleroderma. Mixed connective tissue disease. Polymyalgia rheumatica. Temporal arteritis. Organ-specific autoimmune disease. Rheumatoid arthritis. Diseases resembling rheumatoid arthritis: Still's disease, Sjögren's syndrome, Felty's syndrome, psoriatic arthritis. Ankylosing spondylitis. Reiter's disease. Osteoarthritis

Endocrine disease 162 Thyroid: non-toxic goitre, thyrotoxicosis, hypothyroidism, autoimmune thyroiditis, subacute thyroiditis, thyroid cancer. Pituitary: hypopituitarism, acromegaly, diabetes insipidus, prolactinoma. Adrenal: glands, Cushing's syndrome, Conn's syndrome, adrenogenital syndrome, adrenal insufficiency,

	phaeochromocytoma, multiple endocrine adenomatosis. Shortness of stature. Tests of endocrine function: Cushing's syndrome, adrenocortical insufficiency, thyroid function
Metabolic disease 182	Diabetes mellitus. Carcinoid syndrome. Porphyrria. Lipid disorders. Metabolic bone disease: osteoporosis, osteomalacia and rickets, uraemic bone disease, Paget's disease. Endocrine bone disease: hyperparathyroidism, hypoparathyroidism, thyroid bone disease. Hypercalcaemia. Gout
Renal disease 207	Proteinuria. Nephrotic syndrome. Haematuria. Acute renal failure. Chronic renal failure. Glomerulonephritis. Urinary tract infection. Chronic interstitial nephritis. Polycystic kidney disease. Hepatorenal syndrome. Haemolytic uraemic syndrome. Fluids and electrolytes: salt and water, water excess, inappropriate ADH syndrome, hypokalaemia, hyperkalaemia, metabolic acidosis, hyperaldosteronism
Liver disease 226	Acute viral hepatitis: hepatitis A, hepatitis B, non-A non-B hepatitis. Chronic hepatitis: chronic persistent hepatitis, chronic lobular hepatitis, chronic active hepatitis. Alcoholic hepatitis. Cirrhosis: classification, hepatocellular failure, portal hypertension, management of ascites in cirrhosis. Budd–Chiari syndrome. Rare cirrhoses: idiopathic haemochromatosis, hepatolenticular degeneration (Wilson's disease). Drug jaundice: hypersensitivity reactions, direct hepatotoxicity, haemolytic jaundice
Gastroenterology 238	Gastric and duodenal ulceration. Endocrine tumours of the gut. Gastrointestinal haemorrhage. Hiatus hernia. Ulcerative colitis. Crohn's disease. Steatorrhoea and malabsorption: primary idiopathic steatorrhoea, other causes of malabsorption. Diverticular disease. Irritable bowel syndrome. Ischaemic cholangitis. Pancreas: carcinoma, acute pancreatitis, chronic pancreatitis. Gall bladder: acute cholecystitis, chronic cholecystitis
Respiratory disease 259	Chronic bronchitis and emphysema. Bronchiectasis. Cystic fibrosis. Asthma: severe asthma. Respiratory failure: acute on chronic respiratory failure. Acute anaphylaxis. Pneumonia: bronchopneumonia, lobar pneumonia, other bacterial pneumonias, legionnaire's disease, recurrent bacterial pneumonia, viral pneumonia, mycoplasma pneumoniae, opportunistic infection of the lungs, aspiration pneumonia. Lung abscess. Carcinoma of the bronchus. Bronchial adenoma. Sarcoidosis. Tuberculosis. Occupational lung diseases. Extrinsic allergic alveolitis. Pulmonary embolism. Pneumothorax. Hyperventilation syndrome. Fibrosing alveolitis. Adult respiratory distress syndrome

Cardiovascular disease 285	Ischaemic heart disease: angina, myocardial infarction. Heart failure. Hypertension. Acute rheumatic fever. Valvular heart disease: mitral stenosis, mitral regurgitation, aortic stenosis, aortic regurgitation, tricuspid regurgitation, pulmonary stenosis, atrial myxoma. Congenital heart disease: working classification, atrial septal defect, patent ductus arteriosus, ventricular septal defect, Fallot's tetralogy, pulmonary stenosis, coarctation of the aorta, Eisenmenger's syndrome, Marfan's syndrome. Infective endocarditis: acute, subacute, 'culture-negative endocarditis'. Acute pericarditis. Constrictive pericarditis. Syphilitic aortitis and carditis. Cardiomyopathy: hypertrophic cardiomyopathy, congestive cardiomyopathy. Peripheral arterial disease: intermittent claudication, acute obstruction, ischaemic foot. Raynaud's phenomenon
Dermatology 319	Primary skin disorders: psoriasis, lichen planus, pityriasis rosea, eczema, acne vulgaris, rosacea. Fungus infections: candidiasis, ringworm, treatment of dermatophytes in general. Drug eruptions: urticarial reactions, purpura, other disorders. Skin manifestations of systemic disease: erythema nodosum, haemolytic streptococcal infection, malignancy, xanthomatosis, other rare manifestations. Mucosal ulceration: Behçet's disease. Bullous lesions: dermatitis herpetiformis, pemphigus vulgaris, pemphigoid, erythema multiforme, skin pigmentation
Haematology 330	Anaemia: classification, iron deficiency anaemia, B ₁₂ and folic acid deficiency, pernicious anaemia, haemolytic anaemia, hereditary spherocytosis, hereditary elliptocytosis, glucose-6-phosphate dehydrogenase deficiency, haemoglobinopathies, sickle cell disease, thalassaemia, paroxysmal haemoglobinuria, marrow aplasia. Leukaemias: acute lymphatic leukaemia, chronic lymphatic leukaemia; acute myeloid leukaemia, chronic myeloid leukaemia. Glandular fever. Multiple myelomatosis, Hodgkin's disease. Non-Hodgkin's lymphoma. Infectious mononucleosis.
Drug overdoses 341	Clinical presentation, management
Imported diseases 346	Malaria, Typhoid. Dysentery. Giardiasis. Worms
Recommended reading 351	Journals General texts Specialist texts
354	SI units conversion table
355	Index

Preface to the third edition

There have been great changes in medicine since the second edition was published, and we have had to revise to some extent nearly every page and often altered whole sections completely. We are indebted to the following doctors who have read our revised chapters at an early stage and made further suggestions: their help has been invaluable. We cannot, however, blame them for any errors which we ourselves might have missed. Please write and tell us if you find any.

Peter Carey	Haematology
Aiden Challen	Liver disease
Peter Critchley	Neurology
Dick Dickinson	Gastroenterology
David Ellis	Respiratory diseases
Valerie Kyle	Rheumatic diseases
Neil Lewis	Cardiovascular and renal diseases
Neil Walker	Dermatology
Trevor Wheatley	Endocrine and metabolic diseases

We are also indebted to the many readers who have written to us, including those who have suggested improvements or pointed out inaccuracies. Once again, Sir Edward Wayne has read the entire text and made many useful suggestions.

Some of the ECGs first appeared in Schamroth's *An Introduction to Electrocardiography* (1977, 5th edition, Blackwell Scientific Publications, Oxford) and Table 15 appeared in *Medicine* published by Medical Education International Ltd.

David Rubenstein, David Wayne
January, 1985.

Preface to first edition

This book is intended primarily for the junior hospital doctor in the period between qualification and the examination for Membership of the Royal Colleges of Physicians. We think that it will also be helpful to final year medical students and to clinicians reading for higher specialist qualifications in surgery and anaesthetics.

The hospital doctor must not only acquire a large amount of factual information but also use it effectively in the clinical situation. The experienced physician has acquired some clinical perspective through practice; we hope that this book imparts some of this to the relatively inexperienced. The format and contents are designed for the examination candidate but the same approach to problems should help the hospital doctor in his everyday work.

The book as a whole is not suitable as a first reader for the undergraduate because it assumes much basic knowledge and considerable detailed information has had to be omitted. It is not intended to be a complete textbook of medicine and the information it contains must be supplemented by further reading. The contents are intended only as lecture notes and the margins of the pages can be used by the reader to add additional material of his own.

The book is divided into two parts: *the clinical approach* and *essential background information*. In the first part we have considered the situation which a candidate meets in the clinical part of an examination or a physician in the clinic. This part of the book thus resembles a manual on techniques of physical examination, though it is more specifically intended to help the candidate carry out an examiner's request to perform a specific examination. It has been our experience in listening to candidates' performances in examinations and hearing the examiner's subsequent assessment, that it is the failure of a candidate to examine cases systematically and his failure to behave as if he were used to doing this every day of his clinical life that leads to adverse comments.

In the second part of the book a summary of basic clinical facts is given in the conventional way. We have included most common diseases but not all, and we have tried to emphasise points which are understressed in many textbooks. Accounts are given of many conditions which are relatively rare. It is necessary for the clinician to know about these and to be on the lookout for them both in the clinic and in examinations. Supplementary reading is essential to understand their basic pathology but the information we give is probably all that need be remembered by

the non-specialist reader and will provide adequate working knowledge in a clinical situation. It should not be forgotten that some rare diseases are of great importance in practice because they are treatable or preventable, e.g. infective endocarditis, hepatolenticular degeneration, attacks of acute porphyria. Some conditions are important to examination candidates because patients are ambulant and appear commonly in examinations, e.g. neurosyphilis, syringomyelia, atrial and ventricular septal defects.

We have not attempted to cover the whole of medicine but by cross-referencing between the two sections of the book and giving information in summary form we have completely omitted few subjects. Some highly specialised fields such as the treatment of leukaemia were thought unsuitable for inclusion.

A short account of psychiatry is given in the section on neurology since many patients with mental illness attend general clinics and it is hoped that readers may be warned of gaps in their knowledge of this important field. The section on dermatology is incomplete but should serve for quick revision of common skin disorders.

Wherever possible we have tried to indicate the relative frequency with which various conditions are likely to be seen in hospital practice in this country and have selected those clinical features which in our view are most commonly seen and where possible have listed them in order of importance. The frequency with which a disease is encountered by any individual physician will depend upon its prevalence in the district from which his cases are drawn and also on his known special interests. Nevertheless rare conditions are rarely seen; at least in the clinic. Examinations, however, are a 'special case'.

We have used many generally accepted abbreviations, e.g. ECG, ESR, and have included them in the index instead of supplying a glossary.

Despite our best efforts, some errors of fact may have been included. As with every book and authority, question and check everything—and please write to us if you wish.

We should like to thank all those who helped us with producing this book and in particular Sir Edward Wayne and Sir Graham Bull who have kindly allowed us to benefit from their extensive experience both in medicine and in examining for the Colleges of Physicians.

David Rubenstein, David Wayne
November 1975

原 书 缺 页

原
书
缺
页

Part 1 The clinical approach

Nervous system

The candidate is usually asked to examine a specific area, e.g. 'Examine the cranial nerves', 'Examine the lower limbs', 'Examine the arms' or 'Examine the eyes'. By far the commonest neurological disorders suitable for a clinical examination are multiple sclerosis and the results of cerebrovascular disease. Diabetes is a common disorder which fairly frequently gives rise to neurological manifestations. Carcinomatous neuropathy should always be considered when the signs are difficult to synthesise. Neurosyphilis is becoming progressively rarer—the patient will belong to the prepenicillin era, i.e. be over 50 years old. Parkinsonism is relatively common. Motor neurone disease, myopathies, myasthenia gravis and the neurological manifestations of vitamin B₁₂ deficiency are all rare in practice but more frequently seen in examinations.

In terms of examination technique the practising physician must examine case after case, both normal and abnormal, until he has developed a system which is rapid, accurate and second nature to him. An appearance of professionalism in your neurological examination may encourage the examiner to take a less unfavourable view of minor errors than he might if you appear hesitant, clumsy or imprecise. Analysis of clinical signs is improved if you can remember some anatomical diagrams (pages 4, 5, 6, 21, 107 and 110).

'Examine the cranial nerves'

Many abnormalities of the cranial nerves are the results of chronic disease and patients with them are commonly seen in examinations.

The commonest disorders in examinations are multiple sclerosis (optic atrophy, nystagmus [often ataxic], cerebellar dysarthria), stroke, and Bell's palsy. The manifestation of cerebral tumour, aneurysm, syphilis, dystrophia myotonica and myasthenia gravis are seen much less frequently. It is useful to memorise diagrams of cross-sections of the brain-stem and one of the floor of the fourth ventricle (pages 5 and 6) since these may greatly improve analysis of a cranial nerve lesion. Do not spend long on the first or second cranial nerves unless there is good reason to suspect an abnormality. If the optic fundus is abnormal, the examiner is likely to ask you to look at it specifically. Eye movements must be carefully examined. Do not confuse ptosis (third nerve or sympathetic) with paresis of the orbicularis oculi (seventh nerve). Make sure you can explain clearly and concisely the difference between an upper and lower motor neurone lesion

of the seventh nerve. The corneal reflex is an essential part of the complete examination of the cranial nerves.

The following approach is recommended.

Smell *'Has there been any recent change in your sense of smell?' If so, test formally with 'smell bottles'.*

Eyes Observe, and test when necessary, for:

- *ptosis*

Third nerve lesion (complete or partial ptosis)

Sympathetic lesion (partial ptosis) as part of Horner's syndrome

Muscle weakness. Myasthenia gravis (and rarely, dystrophia myotonica, facio-scapulo-humeral dystrophy, congenital and tabo-paresis)

NB Ptosis is not due to a seventh nerve lesion.

- *visual fields* to confrontation (second nerve) (page 5)

- *external ocular movements* (third, fourth and sixth nerves) (page 7) and *nystagmus* (page 11)

- *the fundi* (second nerve) (page 29)

- *visual acuity* either formally with Snellen's charts or quickly with literature from the bedside locker.

Face (seventh nerve)

- *'Screw up your eyes very tightly.'* Compare how deeply the eyelashes are buried on the two sides. Unilateral weakness is invariably due to a lower motor neurone lesion.

- *'Grin.'* Compare the nasolabial grooves.

Mouth

- *'Clench your teeth'* (fifth nerve, motor). Feel the masseters and test the jaw jerk if indicated. The jaw jerk is obtained from placing one finger horizontally across the front of the jaw and tapping the finger with a tendon hammer with the jaw relaxed and the mouth just open. An increased jaw jerk occurs in upper motor neurone lesions of the fifth cranial nerve (pseudobulbar palsy, page 13).

- *'Open your mouth and keep it open'* (fifth nerve, motor: ptery-




Figure 1. Cross-section through spinal cord.

goids). You should not be able to force it closed. With a unilateral lesion, the jaw deviates towards the weaker side (the weak muscle cannot keep it open).

- 'Say aaah' (ninth and tenth nerves). Normally the uvula and soft palate move upwards and remain central and the posterior pharyngeal wall moves a little. With a unilateral lesion the soft palate is pulled away from the weaker side (there may also be 'curtain movement' of the posterior pharyngeal wall away from the weaker side).

- 'Put your tongue out' (twelfth nerve). Look for wasting, fasciculation and whether it protrudes to one side (towards the weaker side since the weaker muscle cannot push it out).

Neck (eleventh nerve)

- 'Lift your head off the pillows' or 'Put your chin on your right (or left) shoulder' while you resist the movement. Look at, and palpate the sternomastoids. 'Shrug your shoulders' while you push them down. Look at and palpate the bulk of trapezius.

Ears (eighth nerve)

Test hearing with a wrist watch held at various distances from the ears (compare with your own) and perform Weber and Rinne tests. You should ask for an auriscope if indicated. The commonest cause of conductive (air conduction) deafness is wax.

Facial sensation (fifth nerve)

Test the three divisions on both sides with cotton wool. Check the corneal reflexes (often the first clinical deficit in fifth nerve lesions). Ask the patient if the sensation is equally unpleasant on the two sides.

Notes

It is frequently helpful to be able to draw cross-sections of the spinal cord and brain-stem when considering lesions in those areas (figures 1-3).

Field defects

You are in principle comparing the patient's visual fields with your own. When testing his right eye, he should look straight into




Figure 2. Cross-section through open medulla.




Figure 3. Floor of the fourth ventricle showing cranial nerve nuclei.

your left eye with his head held at arm's length. 'Keep looking at my eye and tell me when you first see my finger out of the corner of your eye.' Then bring your finger towards the centre of the field of vision from the four main directions (right, left, up, down). It is preferable to use a white-headed hat pin if you have one. The nasal and superior fields are limited by the nose and eyebrow respectively but this is not often of clinical importance. Field defects are described by the side of the visual field which is lost, i.e. temporal field loss indicates loss of the temporal field of vision and denotes damage to the nasal retina or its connections back to the visual cortex. Perimetry will accurately define defects.

- **Temporal hemianopia** in one eye alone or in both eyes (bitemporal hemianopia) suggests a chiasmal compression usually from a pituitary tumour.
- **Homonymous hemianopia**, i.e. loss of nasal field in one eye and temporal field in the other. This may occur with any postchiasmal lesion, commonly following a posterior cerebral vascular lesion (usually with macular sparing). The side of the field loss is opposite to the side of the damaged cortex (i.e. a right-sided cerebral lesion produces a left homonymous hemianopia).
- **Upper quadrantic field loss** suggests a temporal lesion of the opposite cortex or optic radiation. It is homonymous.

- *Central scotoma.* Loss of vision in the centre of the visual field is detected by passing a white or red tipped pin across the front of the eyes which are held looking forward. This occurs in acute retrobulbar neuritis most commonly due to multiple sclerosis.

Blindness A history of transient blindness, total or partial (with specific field defects usually in one eye) is not uncommon in migraine. It may also follow vertebral arteriography and carotid transient ischaemic attacks.

Sudden blindness also occurs with

- retinal detachment
- acute glaucoma
- vitreous haemorrhage in diabetes
- temporal arteritis and retinal artery or vein obstruction
- fractures of the skull
- raised intracranial pressure

NB The light reflex is absent except in cortical blindness.

Senile changes and glaucoma account for about two-thirds of blindness in this country. Diabetes is the major non-ocular (systemic) cause (7–10%) due chiefly to vitreous haemorrhage and cataract. Trachoma is a common cause on a world-wide basis. *Hysterical blindness* is uncommon and should never be confidently assumed.

The blindness of *temporal arteritis* is preventable if steroid therapy is started in time.

Eye movements

These are controlled by the third, fourth and sixth nerves and conjugate movement integrated by the medial longitudinal bundle. This connects the above nuclei together and to the cerebellum and the vestibular nuclei.

Squint

Congenital concomitant squints are present from childhood and are due to a defect of one eye. The angle between the longitudinal axes of the eyes remains constant on testing extraocular movements, and there is no diplopia.

Paralytic squint is acquired and results from paralysis of one or more of the muscles which move the eye, or paralysis from proptosis. On testing external ocular movements, the angle between the eye axes varies and there is no diplopia.

Rule 1. Diplopia is maximum when looking in the direction of action of the paralysed muscle.

Rule 2. The image furthest from the midline arises from the 'paralysed' eye. This may be determined by covering up each eye in turn and asking which image has disappeared.

NB It is sometimes easier to test movements in each eye separately.