

American
DRUG INDEX

1986

30th Edition

NORMAN F. BILLUPS, R.Ph., M.S., Ph.D.

Dean and Professor of Pharmacy
College of Pharmacy
The University of Toledo

Associate Editor

SHIRLEY M. BILLUPS, R.N., B.Ed.

Oncology Nurse
The Toledo Hospital



J. B. LIPPINCOTT COMPANY
Philadelphia

London New York São Paulo
Mexico City St. Louis Sydney

Copyright © 1986, by J. B. Lippincott Company

Copyright © 1956, 1958, 1959, 1960, 1961, 1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985 by J. B. Lippincott Company

This book is fully protected by copyright and, with the exception of brief excerpts for review, no part of it may be reproduced in any form by print, photoprint, microfilm or any other means, without the written permission of the publishers.

ISBN 0-397-50736-4

Library of Congress Catalog Card Number 55-6286

Printed in the United States of America

Preface

The 30th Edition of the *American Drug Index* (ADI) has been prepared for the identification, explanation, and correlation of the many pharmaceuticals available to the medical, pharmaceutical, and allied health professions. The need for this index has become acute as the variety and number of drugs and drug products have continued to multiply. For this reason the ADI should be most useful to pharmacists, nurses, physicians, medical transcriptionists, dentists, sales personnel, students, and teachers in the fields incorporating pharmaceuticals. *Special note to medical transcriptionists:* Although all names in the ADI appear in bold capitalized type, trade products are easily identifiable by the manufacturer's name in parentheses immediately following the trade name. The names for official products are preceded by a bullet (•) and should appear in lower case in transcription. The therapeutic and pharmaceutical classes are denoted by a box (■) preceding the name and should be a helpful cross-reference guide.

The organization of the ADI falls into the following ten major sections: Monographs of Drug Products, Common Abbreviations Used in Medical Orders, Approximate Practical Equivalents, Common Systems of Weight and Measure, Glossary, Title Changes For U.S.P. XXI and N.F. XVI Monographs, Container Requirements for U.S.P. XXI Drugs, Container and Storage Requirements for Sterile U.S.P. XXI Drugs, Pharmaceutical Company Labeler Code Index Numbers, and Pharmaceutical Manufacturer and/or Drug Distributor Addresses. The U.S.P. XXI and N.F. XVI Title Changes section has been added to make the ADI an even more complete and valuable reference tool.

MONOGRAPHS: The organization of the monograph section of the ADI is fundamentally alphabetical with extensive cross-indexing. Names listed are generic (also called non-proprietary, public name, or common name); brand (also called trade-mark, proprietary, or specialty); and chemical. Synonyms that are in general use also are included. All names used for a pharmaceutical appear in alphabetical order with the pertinent data given under the brand name by which it is made available.

The monograph for a typical brand name product consists of the manufacturer, generic and/or chemical names, composition and strength, pharmaceutical forms available, package size, dosage, and use.

Generic names appear in alphabetical order, followed by the corresponding recognition of the drug to the U.S.P. (United States Pharmacopeia), N.F. (National Formulary), and USAN (United States Adopted Names). Each of these officially recognized generic names is preceded by a bullet (•) at the beginning of each entry. The information is in accord with the U.S.P. XXI and N.F. XVI and through Supplement 2, which became official on July 1, 1985; and the USAN-1986.

British Approved Names, indicated by (B.A.N.), are included in this edition with the permission of the Controller of Her Britannic Majesty's Stationery Office.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to tetracycline HCl. Here under the generic name are listed the various brand names. Following are combinations of tetracycline HCl organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage, and use.

The multiplicity of generic names for the same therapeutic agent has complicated the nomenclature of these agents. Examples of multiple generic names for the same chemical substance are (1) parabromdylamine, brompheniramine; (2) acetaminophen, p-hydroxy acetanilid, N-acetyl-p-aminophenol; (3) guaifenesin, glyceryl guaiacolate, glyceryl guaiacol ether, guaianesin, guaifylline, guaiphenesin, guayanesin, methphenoxydiol; (4) pyrilamine, pyranisamine, pyranilamine, pyraminyl, anisopyradamine.

The cross-indexing feature of the *ADI* permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given. A second form of cross-indexing involves the listing of drugs under various therapeutic and pharmaceutical classes (i.e., antacids, antihistamines, diuretics, laxatives, etc.). Each of these classes is preceded by a box (■) at the beginning of each entry.

ABBREVIATIONS: A listing of Common Abbreviations Used in Medical Orders is included as an aid in interpreting medical orders. The Latin or Greek word and abbreviation are both given with the meaning.

CONVERSION FACTORS: The listing of Approximate Practical Equivalents is added as an aid in calculating and converting dosages among the metric, apothecary, and avoirdupois systems.

WEIGHT AND MEASURE: Tables containing the Common Systems of Weight and Measure are included to aid the practitioner in calculating dosages in the metric, apothecary, and avoirdupois systems.

GLOSSARY: A listing of commonly used terms, with brief definitions, is included as an aid in interpreting the use given for drug monographs listed in the *ADI*.

NEW TITLES FOR U.S.P. XXI AND N.F. XVI MONOGRAPHS: 119 new official monographs appeared through Supplement 2 to the 1985 U.S.P. XXI and N.F. XVI, and an additional 535 new monographs have

appeared since the 1980 U.S.P. XX and N.F. XV. Since many of these new monographs apply to commonly used drug products, this section has been added to aid the practitioner in identifying drug products.

CONTAINER AND STORAGE REQUIREMENTS FOR U.S.P. XXI COMPENDIAL DRUGS: These sections have been added on container and storage requirements specified by the U.S.P. XXI for compendial drugs, to aid the practitioner in storing and dispensing.

ORAL DOSAGE FORMS THAT SHOULD NOT BE CRUSHED: For the first time this year, a section has been added to alert the health care practitioner about oral dosage forms that should not be crushed, and to serve as an aid in consulting with patients. Examples of products falling into the "non-crush" category are extended-release, enteric-coated, encapsulated beads, wax matrix, sublingual, and encapsulated liquid formulations.

LABELER CODE INDEX: The Pharmaceutical Labeler Code Index is presented to aid the practitioner in the identification of drug products. In this section, the codes are listed in numerical order followed by the name of the manufacturer.

MANUFACTURER ADDRESSES: The name, address, and zip code of virtually every American Pharmaceutical Manufacturer and/or Drug Distributor are listed in alphabetical order in this section. Additionally, a pharmaceutical labeler code number appears before the address of each company as a further aid in the identification of drug products.

A special word of appreciation and acknowledgment is given to my wife, Shirley, who served again this year as my Associate Editor—and helped to compile much of the information that appears in the monograph section of this volume. Special thanks are also extended to the manufacturers who supplied product information and made helpful comments and suggestions for this index, and to Dr. Kenneth S. Alexander, who organized the information on Container and Storage Requirements for U.S.P. XXI Drugs.

Correspondence or communication with reference to a drug or drug products listed in the *American Drug Index* should be directed to Norman F. Billups, Ph.D., Dean, College of Pharmacy; The University of Toledo; Toledo, Ohio 43606, or call 419-537-4235.

Norman F. Billups, R.Ph., Ph.D.

Contents

1. Monographs	1-679
2. Common Abbreviations Used In Medical Orders	680-681
3. Approximate Practical Equivalents	682
4. Common Systems of Weight and Measure	683
5. Glossary	684-692
6. New U.S.P. XXI and the N.F. XVI Monographs	693-694
7. Container Requirements for U.S.P. XXI Drugs	695-706
8. Container and Storage Requirements for Sterile U.S.P. XXI Drugs	707-715
9. Oral Dosage Forms That Should Not Be Crushed	716-719
10. Pharmaceutical Company Labeler Code Index	720-725
11. Pharmaceutical Manufacturers and/or Drug Distributors	726-740

[•] Denotes official name: Generic name or chemical name approved by F.D.A. or recognized by the U.S.P., N.F., or USAN.

[■] Denotes therapeutic and pharmaceutical classes.

A

A-200 PYRINATE. (Norcliff Thayer) **Liquid:** Pyrethrins 0.17%, piperonyl butoxide technical 2.00%, petroleum distillate 5.00%. Bot. 2 fl. oz., 4 fl. oz.
Gel: Pyrethrins 0.33%, piperonyl butoxide technical 4.00%, petroleum distillate 5.33%. Tube 1 oz.
 Use: Topically, pediculosis.

A.A.A. OINTMENT. (Jenkins) Ammoniated mercury 2%, acid salicylic 1.25%, acid boric 1%, zinc oxide 15%. Jar 1 oz., 1 lb.
 Use: Impetigo, nonspecific eczema & minor skin irritations.

A AND D HAND CREAM. (Schering) Emollients isopropyl myristate & squalane. Tube 2 oz.
 Use: Dry skin, sunburn, irritations.

A AND D OINTMENT. (Kenyon) Vits. A & D in lanolin-petrolatum base. Jar 1 lb.

A AND D OINTMENT. (Schering) Petrolatum-lanolin. Tube 1.5 oz., 4 oz., Jar 1 lb.

A & D TABLETS. (Barth's) Vit. A 10,000 I.U., D 400 I.U./Tab. Bot. 100s, 500s.

A AND D VITAMIN CAPSULES. (Lannett) Vit. A 5000 I.U., D 400 I.U./Cap. Bot. 500s, 1000s.
 Use: Vit. supplement.

A AND D VITAMIN CREAM. (Eckerd) Bot. 1.5 oz., 4 oz.

A & D VITAMIN OINTMENT. (Quality Generics) Vit. A & D in lanolin-petroleum base Tube 2 oz., Jar 1 lb.
 Use: Emollient.

A AND D VITAMIN OINTMENT. (Lannett) In a petrolatum-lanolin base. Jar 1, 5 lb.

A AND D VITAMIN OINTMENT. (Ulmer) Jar 1 lb., 2 oz.

A-A PLUS. (Galen) Al hydroxide gel 400 mg., Mg oxide 175 mg., Na lauryl sulfate 25 mg., Me cellulose 100 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscine HBr 0.0065 mg./Tab. Bot. 100s, 1000s. Susp. 1 pt.

AAS INFANTIL W/VITAMIN C. (Winthrop Products) Acetylsalicylic acid, Vit. C.
 Use: Analgesic, antipyretic.

AAS TABLETS. (Winthrop Products) Acetylsalicylic acid.
 Use: Analgesic, antipyretic.

ABBOCIN. Oxytetracycline, B.A.N.

ABBOKINASE. (Abbott) Urokinase 250,000 I.U./5 ml. Lyophilized powder. Vial 5 ml.
 Use: Plasminogen activator.

ABBOKINASE OPEN-CATH. (Abbott) Urokinase for catheter clearance 5,000 I.U./ml. Unival.
 Use: Clearance of fibrin occluded catheters.

ABBOTT AFP-EIA. (Abbott) Test Kits 100s.
 Use: Enzyme immunoassay for the quantitative measurement of alpha-fetoprotein in human serum to aid in the management of testicular cancer.

ABBOTT CMV TOTAL AB EIA. (Abbott) Test kits 100s.
 Use: Enzyme immunoassay for the detection of antibody to cytomegalovirus in human serum, plasma, and whole blood.

ABBOTT-HB_e TEST. (Abbott) Test kits 100s.

Use: Radioimmunoassay or enzyme immunoassay for detection of hepatitis B_e antigen and/or antibody to hepatitis B_e antigen.

ABBOTT IgE EIA. (Abbott) Test kits 100s.

Use: Enzyme immunoassay for quantitative determination of IgE in human serum and plasma.

ABC COMPOUND w/CODEINE #3. (Zenith) Aspirin 325 mg., caffeine 40 mg., butalbital 50 mg., codeine phosphate 30 mg./Cap.
 Use: Analgesic.

ABCDG VITAMIN CAPSULES. (Lannett) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 2 mg., C 30 mg./Cap. Bot. 1000s.

Use: Vitamin supplement.

ABDEC BABY VITAMIN DROPS. (Parke-Davis) Vit. A 1500 I.U., D 400 I.U., B-1 0.5 mg., B-2 0.6 mg., B-6 0.4 mg., pantothenic acid 3 mg., niacin 8 mg., C 35 mg./ml. Bot. 1½ oz. w/dropper.
 Use: Vitamin deficiencies.

ABDEC WITH FLUORIDE BABY VITAMIN DROPS.

(Parke-Davis) Vit. A 1500 I.U., D 400 I.U., C 35 mg., B-1 0.5 mg., B-2 0.6 mg., B-6 0.4 mg., niacin 8 mg., pantothenic acid 3 mg./ml. w/fluoride. Bot. 50 ml.
 Use: Vitamin therapy with fluoride.

ABDO. (Eastwood) Cap. Bot. 100s.

ABDOL W/MINERALS. (Parke-Davis) Vit. C 50 mg., B-2 2.5 mg., B-12 1 mcg., B-6 0.5 mg., B-1 2.5 mg., niacinamide 20 mg., Vit. A 5000 I.U., D 400 I.U., Ca pantothenate 2.5 mg., folic acid 0.1 mg., iodine 0.15 mg., Mn 1 mg., K 5.0 mg., Fe 15 mg., Cu 1 mg., Zn 0.5 mg., Mg 1 mg., Ca 44 mg., P 34 mg./Cap. Bot. 100s.
 Use: Vitamin-mineral deficiency states.

ABSORBABLE CELLULOSE COTTON OR GAUZE.

See: Oxidized Cellulose (Various Mfr.)

• **ABSORBABLE DUSTING POWDER, U.S.P. XXI.**

• **ABSORBABLE GELATIN FILM, U.S.P. XXI.** Sterile nonantigenic, absorbable, water-insoluble, gelatin film.
 See: Gelfilm (Upjohn)

• **ABSORBABLE GELATIN SPONGE, U.S.P. XXI.**

Gelatin sponge.

Use: Surgical aid.

See: Gelfoam, Preps. (Upjohn)

• **ABSORBABLE SURGICAL SUTURE, U.S.P. XXI.**

(Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

Use: Surgical aid.

• **ABSORBENT GAUZE, U.S.P. XXI.**

Use: Surgical aid.

ABSORBINE ARTHRITIC PAIN LOTION. (W.F.

Young) Bot. 2 oz., 4 oz.

ABSORBINE FOOT POWDER. (W.F. Young) Zinc stearate, parachloroxylenol, aluminum chlorohydrate, allantoinate, benzethonium chloride, menthol. Plastic bot. 3 oz. w/shaker top.
 Use: Athlete's foot treatment.

ABSORBINE, JR. (W.F. Young) Wormwood, thymol, chloroxylenol, menthol, acetone, zinc stearate, parachloroxylenol, aluminum chlorohydrate, allantoinate, benzethonium chloride, menthol. Liq. Bot. 1 oz., 2 oz., 4 oz., 12 oz. w/applicator.

Use: Muscle soreness, stiffness, treatment of athlete's foot.

ABUSCREEN. (Roche Diagnostics) An immunological and radiochemical assay for morphine and morphine glucuronide in nanogram levels. Utilizes ^3H labeled morphine assayed with liquid scintillation counters using conventional beta counting procedures and ^{125}I labeled morphine requiring gamma scintillation equipment. 100 tests.

ACABEL. Bevonium Methylsulphate, B.A.N.

• **ACACIA.** N.F. XVI. Syrup N.F. XVI. Acacia senegal, gum arabic. (Perick) pow., 0.25 lb., 1 lb.; tears, 0.25-1 lb.; whole, 0.25-1 lb.

Use: Demulcent, emulsifier, suspending agent.

A-CAINE. (A.V.P.) Rectal ointment. Tube 1.25 oz.

Use: Rectal care, anti-hemorrhoidal.

A CAPSULES. (Stanlabs) Natural Vit. A 5,000 I.U., 10,000 I.U., 25,000 I.U., 50,000 I.U./Cap. Bot. 100s. Synthetic or soluble, 10,000 I.U., 25,000 I.U., 50,000 I.U./Cap. Bot. 100s.

A-CAPS. (Drug Industries) Vit. A 50,000 I.U./Cap. Bot. 100s, 500s.

Use: Vitamin A therapy.

• **ACARBOSE.** USAN.

Use: Inhibitor.

ACCELERASE. (Organon) Pancrelipase: Lipase 4000 U., amylase 15,000 U., protease 15,000 U., other pancreatic enzymes mixed with bile salts 65 mg., cellulase 2 mg., calcium carbonate 20 mg./Cap. Bot. 60s.

Use: Functional digestive disorders.

ACCURBRON. (Merrell Dow) Theophylline, anhydrous 10 mg./ml. Bot. pt.

Use: Bronchial asthma treatment.

ACCUSENS T TASTE FUNCTION KIT. (Westport)

Test for ability to distinguish among salty, sweet, sour and bitter tastants. Kit contains 15 bottles (60 ml.) tastants and 30 taste record forms.

Use: Diagnostic taste function kit.

ACCUTANE. (Roche) Isotretinoin 10 mg., or 40 mg./Cap. Bot. 100s.

Use: Treatment of severe recalcitrant cystic acne.

A-C-D SOLUTION, U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution.

Baxter-600 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol.

Cutter-500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol.

Diamond-(Abbo-Vac) 250 cc., 500 cc.

Use: Anticoagulant for preparation of plasma or whole blood.

A-C-D SOLUTION MODIFIED. (Squibb) Acid citrate dextrose anticoagulant solution modified.

Use: Anticoagulant for use in radiolabeling red blood cells.

ACDRILE. Methyl Cysteine, B.A.N.

• **ACEBUTOLOL.** USAN. (+)-1-(2-Acetyl-4-butyramido-phenoxy)-3-isopropylaminopropan-2-ol. (Ives) (+)-N-[3-Acetyl-4-[2-hydroxy-3-[(1-methylethyl)amino]propoxy]-phenyl] butamide.

Use: Beta adrenergic receptor blocking agent.

See: Sectral, Cap. (Ives)

• **ACECAINIDE HYDROCHLORIDE.** USAN.

Use: Cardiac depressant.

• **ACECLIDINE.** USAN. 3-Quinuclidinol acetate (ester). Glauco-stat. Under study.

Use: Parasympathomimetic.

• **ACEDAPSONE.** USAN. 4',4''-Sulfonylbis (acetanilide). Use: Antimalarial; antibacterial (leprostatic).

ACEDOVAL. (Vale) Dover's powder 15 mg., ipecac 1.5 mg., aspirin 162 mg., caffeine anhydrous 8.1 mg./Tab. Bot. 100s, 1000s, 5000s.

ACEDYNE. (Vale) Aspirin 64.8 mg., caffeine anhydrous 3.2 mg., atropine sulfate 0.032 mg./Tab. Bot. 100s, 1000s, 5000s.

ACEFYLLINE PIPERAZINE. Acepifylline, B.A.N.

• **ACEGLUTAMIDE ALUMINUM.** USAN.

Use: Antulcerative.

ACELAX. (Marcen) Danthron 100 mg., prune concentrate 64 mg., Chondrus drispus 64 mg./Tab. Bot. 100s, 1000s.

Use: Laxative.

ACENOCOUMAROL. 3-(α -Acetonyl-4-nitrobenzyl)-4-hydroxy-coumarin. Nicoumalone, B.A.N.

A'CENOL. (Kay) Acetaminophen 5 gr./Tab. Bot. 1000s.

Use: Analgesic.

ACEPHEN. (G&W) Adult: Acetaminophen 650

mg./Supp. Box 12s, 100s. **Pediatric:**

Acetaminophen 120 mg./Supp. Box 12s, 100s.

Use: Analgesic, antipyretic.

ACEPIFYLLINE. B.A.N. Piperazine theophyllin-7-ylate. Acefylline Piperazine (I.N.N.). Etophyllate.

Use: Spasmolytic.

ACEPROMAZINE. 10-(3-Dimethylaminopropyl)-3-acetyl-phenothiazine maleate. Plegicil. (Ayerst) 1-(10-(3-(Dimethylamino)propyl)-10H-phenothiazin-2-yl)ethanone e(Z)-2-butenedioate.

Use: Tranquillizer.

ACEPROMAZINE. B.A.N. 2-Acetyl-10-(3-dimethylaminopropyl)phenothiazine. Notensil maleate.

Use: Tranquillizer.

• **ACEPROMAZINE MALEATE.** USAN.

Use: Sedative.

ACEROLA C. (Kirkman) Vit. C 100 mg./Wafer. Bot. 100s.

ACEROLA-C. (Barth's) Vit. C 300 mg./Wafer. Bot. 30s, 90s, 180s, 360s.

ACEROLA-PLEX. (Barth's) Vit. C 100 mg., bioflavonoids 50 mg./Tab. Bot. 100s, 500s.

ACETA. (Century) Acetaminophen. **Elix.** (120 mg./5 cc) Bot. 4 oz., pint, gal. **Tab.** (5 gr.) Bot. 100s, 1000s, 3000s, 5000s.

Use: Analgesic.

ACETA W/CODEINE. (Century) Acetaminophen 325 mg., codeine phosphate 0.5 gr./Tab. Bot. 100s.

Use: Analgesic.

ACETAMIDE. N-[5-(Aminosulfonyl)-3-methyl-1,3,4-thiadiazol-2(3H)-ylidene]-. Methazolamide.

ACETAMIDE, 2,2-DICHLORO-N-[2-HYDROXY-1-(HYDROXYMETHYL)-2-(4-NITROPHENYL)ETHYL].

Chloramphenicol, U.S.P. XXI.

ACETAMIDE, 2-(DIETHYLAMINO)-N-(2,6-DIMETHYLPHENYL)-. Lidocaine, U.S.P. XXI.

ACETAMIDE, N-[5-(AMINOSULFONYL)-1,3,4-THIADIAZOL-2-YL]-. Acetazolamide, U.S.P. XXI.

ACETAMIDE, N-(4-HYDROXYPHENYL)-. Acetaminophen, U.S.P. XXI.

ACETAMIDE, N-(5,6,7,9-TETRAHYDRO-1,2,3,10-TETRAMETHOXY-9-OXOBENZO[a]-HEPTALEN-7-YL)-. Colchicine, U.S.P. XXI.

p-ACETAMIDOBENZOIC ACID SALT of 2-dimethylaminoethanol.

See: Deaner, Tab. (Riker).

5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALAMIC ACID. Iothalamic Acid, U.S.P. XXI.

5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALAMIC ACID, SODIUM SALT. Iothalamate Sodium Injection, U.S.P. XXI.

• **ACETAMINOPHEN, U.S.P. XXI.** Elix., for Effervescent Oral Soln., Tabs, Cap. U.S.P. XXI. N-Acetyl-p-amino-phenol. Acetamide, N-(4-hydroxyphenyl)-. 4'-Hydroxyacetanilide.

Use: Analgesic, antipyretic.

See: Acephen, Supp. (G&W)

Aceta, Tab. Elix., Supp. (Century)

Acetagesic (Kay)

Actamin, Tab. (Buffington)

Acetaminophen Uniserts, Suppos. (Upsher-Smith)

Aminodyne, Elix. (Bowman)

Anapap, Tab. (Fellows)

Anelix, Elix. (Kirkman)

Anuphen, Supp. (Comatic)

Apap, Cap., Tab. (Various Mfr.)

Capital, Tab. (Carnrick)

Chlor-A-Tyl. (Jenkins)

Dapa, Tab. (Ferndale)

Datriil 500, Tab. (Bristol-Myers)

Dimindol, Tab., Elix. (Blue Line)

Dolanex, Elix. (Lannett)

Dularin, Syr. (Donner)

Febridol, Elix., Tab. (Amid)

Fendon, Tab. (APC)

G-1 (Hauck)

Janupap, Tab. (Reid-Provident)

Lestemp, Elix. (Reid-Provident)

Liquiprin, Solu. (Norcliff Thayer)

Lyteca, Syr., Tab. (Westerfield)

Medigesic (Medics)

Nebs, Tab., Liq. (Eaton)

Nilprin 7.5, Tab. (AVP)

Parten, Tab. (Parmed)

Phenaphen, Cap., Tab. (Robins)

Proval, Cap., Elix., Drop., Tab. (Reid-Provident)

SK-Apap, Tab., Elix. (SmithKline)

Tapar, Tab. (Parke-Davis)

Temetan, Elix., Tab. (Nevin)

Tempra, Drop, Syr., (Mead Johnson)

Tenlap, Elix. (Merrell Dow)

Tussapap, Tab. (Sutliff & Case)

Tylenol, Drops, Elix., Liq., Tab., Chew. Tab. (McNeil)

Tylenol Extra-Strength, Tab., Cap. (McNeil)

Valadol, Tab., Liq. (Squibb)

Valorin Tab. (Otis Clapp)

ACETAMINOPHEN W/COMBINATIONS

Aceta w/Codeine, Tab. (Century)

Akes-N-Pain, Cap. (Edw. J. Moore)

Al-Ay Modified, Tab. (Bowman)

Algecol, Tab. (Winston)

Allerest Headache Strength, Tab. (Pharmcraft)

Allygesic W/Ergotamine, Cap. (Elder)

Allylvon, Cap. (Elder)

Alumadrine, Tab. (Fleming)

Amaphen, Cap. (Trimen)

Aminodyne, Elix. (Bowman)

Anaphen, Cap. (Mallard)

Anodynos-DHC, Tab. (Berlex)

Anodynos Forte, Tab. (Buffington)

Apap w/Codeine, Tab. (Central)

Apapalone, Tab. (Amfre-Grant)

Arthol Tabs. (Towne)

Arthralgen, Tab. (Robins)

Bancaps (Westerfield)

Bancaps-C (Westerfield)

Banesin-Forte, Tab. (Westerfield)

Bexahist, Cap. (Bexar)

Biophens-S, Tab. (La Crosse)

Blanex, Cap. (Edwards)

Bowman Cold Tab. (Bowman)

B-Pap., Liq. (Wren)

BQ Cold Tabs (Bristol-Myers)

Bromo-Seltzer, Gran. (Warner-Lambert)

Butigetic, Tab. (McNeil)

Capital w/Codeine, Tab., Susp. (Carnrick)

Cardui, Tab. (Chattem Labs.)

Cerose Comp., Cap. (Ives)

Chexit, Tab. (Dorsey)

Christodyne-DHC, Tab. (Paddock)

Clistin-D, Tab. (McNeil)

Coastaldyne, Tab. (Coastal)

Coastalgic, Tab. (Coastal)

Codalan, Tab. (Lannett)

Codap (Reid-Provident)

Codimal, Tab. (Central)

Colrex Compound, Elix. (Rowell)

Comtrex, Cap., Liq., Tab. (Bristol-Myers)

Conar-A, Tab., Susp. (Beecham Labs)

Conex, Preps. (Westerfield)

Congesprin, Liq., Tab. (Bristol-Myers)

Contact Jr., Liq. (Menley & James)

Coricidin Sinus Headache Tab. (Schering)

Coryban-D, Syr. (Roerig)

Corzans, Tab. (Moore Kirk)

Co-Tylenol, Liq., Tab. (McNeil)

Dapase, Tab. (Ferndale)

Darvocet-N, Tab. (Lilly)

Darvocet-N 100, Tab. (Lilly)

Daycare, Liq. (Vicks)

Demerol APAP, Tab. (Winthrop-Breon)

Dengesic, Tab. (Scott-Alison)

Desa-Hist AT, Tab. (Pharmics)

Dolene AP-65, Tab. (Lederle)

Dolopar, Liq. (O'Neal)

Drinacet Cap. (Phillips Roxane)

Drinophen, Cap. (Lannett)

D-Sinus, Cap. (Laser)

Dularin-TH, Tab. (Dooner)

Duoprin, Tab. (Dunhall)

Duo-3X, Tab. (Med. Spec.)

Dynosal, Tab. (Mallard)
 Empracet with Codeine, Tab. (Burroughs Wellcome)
 Enz-Cold, Cap. (Edw. J. Moore)
 Esemgesic, Tab. (Spencer-Mead)
 Esgic, Tab. (Gilbert)
 Excedrin, Cap., Tab. (Bristol-Myers)
 F.C.A.H. Caps. (Scherer)
 Femicin, Tab. (Norcliff Thayer)
 Fendol, Tab. (Buffington)
 Gemnisyn, Tab. (Rorer)
 Guaiamine, Cap. (Sutliff & Case)
 Histalets, Tab. (Edw. J. Moore)
 Histogesic, Tab. (Century)
 Histosal #2, Tab. (Ferndale)
 Hi-Temp, Tab. (Saron)
 Hycomine Comp., Tab. (Du Pont)
 Isomel, Cap. (Reid-Provident)
 Kiddies Sialco, Tab. (Foy)
 Kleer, Syr. (Scrip)
 Koryza Tab. (Fellows-Testagar)
 Maxigesic, Cap. (W.F. Merchant)
 Midran, Tab. (Quality Generics)
 Midrin, Cap. (Carnrick)
 Myocalm, Tab. (Parmed)
 Myolate, Tab. (Panray)
 Naldetuss, Syr. (Bristol)
 N-D Gesic, Tab. (Hyrex)
 Neo-Pyranistan, Tab. (Standex Labs.)
 Nokane, Tab. (Wren)
 Novahistine Singlet, Tab. (Merrell Dow)
 Novahistine Sinus Tablets, Tab. (Merrell Dow)
 Nyloac PB, Tab. (Beard)
 Nyquil, Liq. (Vicks)
 Opasal, Cap. (Elder)
 Ornex, Cap. (Menley & James)
 Pamprin, Tab. (Chatter Labs.)
 Panitol H.M.B., Tab. (Wesley)
 Panritis, Tab. (Pan Amer.)
 Parafon Forte, Tab. (McNeil)
 Partuss-A, Tab. (Parmed)
 Partuss T.D., Tab. (Parmed)
 Pavadon Elix. (Coastal)
 Pedituss, Liq. (Sutliff & Case)
 Pedric, Elix., Wafers (Vale)
 Percogesic, Tab. (Du Pont)
 Phenahist, Cap., Elix. (Amid)
 Phenaphen #2, #3, #4 (Robins)
 Phenaphen-650, Tab. (Robins)
 Phrenilin, Tab. (Carnrick)
 Presalin, Tab. (Mallard)
 Prodolor, Tab. (Amfre-Grant)
 Proval No. 3, Tab. (Reid-Provident)
 Renpap, Tab. (Wren)
 Rentuss, Cap., Syr. (Wren)
 Repan, Tab. (Everett)
 Rhinex, D-Lay, Tab. (Lemmon)
 Rhinidrin, Tab. (Central)
 Rhinogesic, Tab. (Vale)
 Rhinspec, Tab. (Lemmon)
 S.A.C. Sinus, Tabs. (Towne)
 Salatin w/Codeine, Tab. (Ferndale)
 Saleto, Tab. (Mallard)
 Saleto-D, Tab. (Mallard)

Salphenyl, Cap., Liq. (Mallard)
 Santussin, Prep. (Sandia)
 Scotgesic, Cap., Elix. (Scott/Cord)
 Scotuss Pediatric Cough Syr. (Scott/Cord)
 Sedacane, Cap. (Edw. J. Moore)
 Sedalgesic, Tab. (Table Rock)
 Sedragesic, Tab. (Lannett)
 Sialco, Tab. (Foy)
 Sinarest, Tab. (Pharmacraft)
 Sine-Aid, Tab. (McNeil)
 Sinoze, Tab. (Bowman)
 Sinubid, Tab. (Warner Chilcott)
 Sinulin, Tab. (Carnrick)
 Sinumal, Tab., Liq. (Cook)
 Sinus Tab., Tab. (Zenith)
 Sinutab, Prods. (Warner-Lambert)
 SK-APAP W/codeine, Tab. (SmithKline)
 SK-65 APAP, Tab. (SmithKline)
 Spantuss, Liq. (Arco)
 Spendrisin, Tab. (Spencer-Mead)
 Stopain, Tab. (Quality Generics)
 Sunril, Cap. (Emko)
 Super-Anahist Tablets (Warner-Lambert)
 Symptomax, Liq. (Reid-Provident)
 Talacen, Cap. (Winthrop-Breon)
 T-Caps, Cap. (Burlington)
 Tega-Code, Cap. (Ortega)
 Tegapap, Liq., Tab. (Ortega)
 Teragen, Tab. (O'Neal)
 T-Gesic, Cap. (Tennessee)
 Triaminicin, Tab. (Dorsey)
 Triaprin, Cap. (Dunhall)
 Trind, Syr. (Mead Johnson)
 Trind-DM, Syr. (Mead Johnson)
 Tussagesic, Tab., Susp. (Dorsey)
 Two-Dyne, Tab. (Hyrex)
 Tylenol Preps. (McNeil)
 Tylenol with Codeine, Elixir, Tab. (McNeil)
 Tylox, Cap. (McNeil)
 Valihist, Cap. (Clapp)
 Vicks Daycare, Liq. (Vicks)
 Windolor, Tab. (Winston)
 Wygesic, Tab. (Wyeth)
 Zenex Comp., Cap. (Zenith)

• **ACETAMINOPHEN AND ASPIRIN TABLETS,**
 U.S.P. XXI.

Use: Analgesic.

• **ACETAMINOPHEN AND DIPHENHYDRAMINE**
CITRATE TABLETS, U.S.P. XXI.

• **ACETAMINOPHEN ORAL SUSPENSION, U.S.P.**
XXI.

Use: Analgesic.

ACETAMINOPHENOL

See: Acetaminophen.

ACETAMINOPHEN UNISERTS. (Upsher-Smith)

Acetaminophen 120 mg./Supp. Ctn. 12s; 325
 mg./Supp. Ctn. 12s, 50s; 650 mg./Supp. Ctn. 12s,
 50s, 500s.

Use: Analgesic, antipyretic.

ACETANILID. (Various Mfr.) (Acetylamino benzene,
 acetylaniline, antifebrin) N-phenylacetamide cry.

Use: Antipyretic, analgesic.

ACETARSOL.

See: Acetarstone, Tabs.

ACETARSTONE. 3-Acetamido-4-hydroxy-phenylarsonic acid. Acetarsol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Osvarsan, Paroxyl, Stovarsol.

ACETARSTONE SALT OF ARECOLINE.

See: Drocabil.

ACETAZIDE. Acetazolamide, B.A.N.

• **ACETAZOLAMIDE**, U.S.P. XXI. Tab. U.S.P. XXI. N-(5-Sulfamoyl-1,3,4-thiadiazol-2-yl) acetamide. Acetamide, N-[5-(aminosulfonyl)-1,3,4-thiadiazol-2-yl]-.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Tab., Sequels (Lederle)

Hydrazol, Tab. (Softcon)

ACETAZOLAMIDE. B.A.N. 5-Acetamido-1,3,4-thiadiazole-2-sulfonamide.

Use: Treatment of glaucoma.

See: Acetazide

Diamox

• **ACETAZOLAMIDE SODIUM, STERILE**, U.S.P. XXI.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Inj. (Lederle)

ACET-DIA-MER-SULFONAMIDE. Sulfacetamide, sulfadiazine & sulfamerazine.

See: Buffonamide, Susp., Tab. (Reid-Provident)

Buf-Sul, Tabs. & Susp. (Sheryl)

Cetazine, Tab. (Bowman)

Coco-Diazine, Susp. (Lilly)

Sulfadiazine, sulfamerazine & sulfacetamide, Susp.

ACETEST REAGENT TABLETS. (Ames) Sod.

nitroprusside, disodium phosphate, aminoacetic acid, lactose. Tab. Bot. 100s, 250s.

Use: Test for acetone or diacetic acid in urine.

• **ACETIC ACID**, U.S.P. XXI, Glacial, Otic soln., U.S.P. XXI. Diluted, N.F. XVI.

Use: Pharmaceutical aid (acidifying agent).

• **ACETIC ACID IRRIGATION**, U.S.P. XXI. 0.25% solution. (Abbott) 250 ml glass cont.; 250 ml.

Use: Irrigating solution

ACETIC ACID, POTASSIUM SALT. Potassium

Acetate, U.S.P. XXI.

ACETICYL.

See: Acetylsalicylic Acid.

ACETILUM ACIDULATUM.

See: Acetylsalicylic Acid. (Various Mfr.)

• **ACETOHEXAMIDE**, U.S.P. XXI. Tab. U.S.P. XXI. N-(p-acetylphenylsulfonyl)-N'-cyclo-hexylurea. 1-((p-acetylphenylsulfonyl)-3-cyclohexylurea. Benzenesulfonamide, 4-acetyl-N-[[cyclohexylamino]carbonyl].

Use: Blood sugar lowering compound, antidiabetic.

See: Dymelor, Tab. (Lilly)

• **ACETOHYDROXAMIC ACID**, U.S.P. XXI. Tab. U.S.P. XXI.

Use: Enzyme inhibitor.

See: Lithosten (Mission)

ACETOJEN, JR. (Jenkins) Acetylsalicylic acid 1.5 gr., phenacetin 1 gr., hyoscyamus ext. 0.0125 gr., (total alkaloids 0.0002 gr.) gelsemium ext. 0.0125 gr./Tab. Bot. 1000s.

ACETOLAX. (Mills) Acetphenylisatin 5 mg., Vit. B-1 1 mg., sod. carboxymethyl cellulose 500 mg./Tab. Bulk Pkg. 1000s.

Use: Laxative.

ACETOL.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOMEROCTOL. Acetato [2-hydroxy-5-(1,1,3,3-tetramethylbutyl) phenyl] mercury. Under study.

Use: Topical antiseptic.

• **ACETONE**, N.F. XVI.

ACETONE or DIACETIC ACID TEST.

See: Acetest, Tab. (Ames)

3-(alpha-ACETONYLBENZYL)-4-HYDROXYCOUMARIN SODIUM SALT. Warfarin Sodium.

ACETOPHEN CAPSULE. (Jenkins) Aspirin 3.5 gr., phenacetin 2.5 gr., caffeine 0.5 gr./Cap. Bot. 1000s.

ACETOPHEN C.T. GREEN. (Jenkins) Aspirin 3.5 gr., phenacetin 2.5 gr., caffeine 0.25 gr./Tab. Bot. 1000s.

• **ACETOPHENAZINE MALEATE**, U.S.P. XXI. Tab. U.S.P. XXI. 2-Acetyl-10- [3-[4-(β-hydroxyethyl)-piperazinyl] propyl] phenothiazine dimaleate. 10-[3-[4-(2-Hydroxyethyl)-1-piperazinyl]-propyl]-phenothiazine-2-yl methyl ketone maleate (1:2).

Use: Tranquilizer.

See: Tindal Maleate, Tab. (Schering)

ACETOPHENETIDIN. Phenacetin, U.S.P. XXI.

Ethoxyacetanilide.

Use: Antipyretic, analgesic.

ACETORPHINE. B.A.N. O²-Acetyl-7,8-dihydro-7α[1(R)-hydroxy-1-methylbutyl]-O⁶-methyl-6,-14-endoethenormorphine.

Use: Narcotic analgesic.

ACETOSAL.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIC ACID.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIN.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSULFONE. 4,4'-Diaminodiphenylsulfone-2-N-acetylsulfonamide sodium. (N¹-acetyl-6-sulfanilyl-metanilamido) sodium.

• **ACETOSULFONE SODIUM.** USAN.

Use: Antibacterial.

ACETOXYPHENYLMERCURY.

See: Phenylmercuric acetate.

ACETPHENARSINE.

See: Acetarstone, Tab. (City Chemical)

ACETPHENILISATIN.

See: Oxyphenisatin Acetate Prep.

W/D-Calcium pantothenate, sodium sulfosuccinate.

See: Peri-Pantyl, Tab. (McGregor)

W/Docusate sodium, sodium carboxymethylcellulose.

See: Scrip-Lax, Tab. (Scrip)

ACETRIZOATE SODIUM. 3-Acetamido-2,4,6-triiodobenzoate sodium.

W/Polyvinylpyrrolidone.

See: Salpix, Vial (Ortho)

ACETRIZOIC ACID. 3-Acetyl-amino-2,4,6-triiodobenzoic acid.

See: Acetrizoate, Sod.

ACET-THEOCIN SODIUM.

See: Theophylline Sod. acetate (Various Mfr.)

ACETYL ADALIN.

See: Acetylcarbromal. (Various Mfr.)

p-ACETYLAMINOENZALDEHYDE THIOSEMICARBAZONE. (Amithiozone, Antib, Berculon A, Benzothiozon, Conteben, Myuizone, Neustab, Tebethion, Thiomicid, Thioparamizone, Thiacetazone).

Use: With streptomycin in the treatment of tuberculosis.

ACETYLAMINOENZENE.

See: Acetanilid. (Various Mfr.)

N-ACETYL-p-AMINOPHENOL. Acetaminophen, U.S.P. XXI.**ACETYLANILINE.**

See: Acetanilid. (Various Mfr.)

ACETYL-BETA-METHYLCHOLINE CHLORIDE.

See: Mecholyl Chloride, Amp. (Baker, J.T.)

ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE.

See: Acetylcarbromal. (Various Mfr.)

ACETYLCARBROMAL. (Meyer) Acetyl adalin, acetylbromodiethylacetyl-carbamide. Pow. for mfg.

Use: Sedative.

See: Paxarel, Tab. (Circle)

W/Bromisovalum, scopolamine aminoxide HBr.

See: Tranquinal, Tab. (Barnes-Hind)

• ACETYLCHOLINE CHLORIDE, U.S.P. XXI. For Ophth. Soln., U.S.P. XXI. Ethanaminium,2-(acetyloxy)-N, N, N-trimethyl-, chloride. Choline chloride acetate.

Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator. Paroxysmal tachycardia.

See: Miochol Ophthalmic (CooperVision)

ACETYLCHOLINE-LIKE THERAPEUTIC AGENTS.

See: Cholinergic agents.

• ACETYLCYSTEINE, U.S.P. XXI. Sol. U.S.P. XXI. N-Acetyl-L-cysteine.

Use: Mucolytic agent.

See: Mucomyst, Sol. (Mead Johnson)

• ACETYLCYSTEINE AND ISOPROTERENOL HYDROCHLORIDE INHALATION SOLUTION, U.S.P. XXI.

Use: Mucolytic agent.

See: Mucomyst with Isoproterenol, Sol. (Mead Johnson)

ACETYLCYSTEINE. B.A.N. N-Acetyl-L-cysteine. Air-bron

Use: Liquefaction of bronchial secretion.

ACETYLDIGITOXIN. α -Digitoxin monoacetate. Acetyldigitoxose. The natural-occurring cardiac glycoside defined as the alpha acetyl ester of digitoxin.

Use: Digitoxin therapy. Cardiotonic.

ACETYLDIHYDROCODEINONE. Thebacon, B.A.N.**ACETYLIN.**

See: Acetylsalicylic Acid. (Various Mfr.)

ACETYLMETHADOL. I.N.N. Methadyl acetate, B.A.N.**ACETYLPHENYLISATIN.**

See: Oxyphenisatin Acetate.

1-(p-ACETYLPHENYL)SULFONYL-3-CYCLO-HEXYLUREA. Acetohexamide.**ACETYLRESORCINOL.**

See: Resorcinol Monoacetate. (Various Mfr.)

ACETYLSAL.

See: Acetylsalicylic Acid

ACETYLSALICYLATE ALUMINUM.

See: Aluminum Aspirin

ACETYLSALICYLIC ACID. Aspirin, U.S.P. XXI.

Acidum acetylsalicylicum, 2-acetoxybenzoic acid, Acetillum Acidulatum, Acetophen, Acetui, Acetosali, Acetosalic Acid, Acetosalin, Aceticyl, Acetylsal, Acylpyrin, Aspro, Helicon, Rhodine, Salacetyl, Salcetogen, Saletin.

See: Aspirin Preps. (Various Mfr.)

ACETYLSALICYLIC ACID, ACETOPHENETIDIN AND CAFFEINE.

See: A.P.C., Prep. (Various Mfr.)

ACETYL SULFAMETHOXYPYRIDAZINE. 3-(N-Acetyl-sulfanilamido)-6-methoxypyridazine.**N¹-ACETYLSULFANILAMIDE.** (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulfacetamide, N-Sulfanilylacetamide.)

Use: Sulfonamide therapy.

ACETYL SULFISOXAZOLE. Sulfisoxazole Acetyl, U.S.P. XXI.

See: Gantrisin Acetyl, Syr., Susp. (Roche)

Lipo-Gantrisin Acetyl, Liq. (Roche)

ACETYLTANNIC ACID. Tannic acid acetate.

Use: Antiperistaltic.

AC EYE DROPS. (Walgreen) Tetrahydrozoline HCl 0.05%, zinc sulfate 0.25%. Bot. 0.75 oz.

Use: Allergy, colds, eye relief.

ACHLORHYDRIA DETERMINATION.

See: Diagnex Blue, Prep. (Squibb)

ACHLORHYDRIA THERAPY.

See: Acidol. (Various Mfr.)

Glutamic Acid HCl (Various Mfr.)

Muripsin, Tab. (Norgine)

Normacid, Tab. (Stuart)

ACHOL. (Enzyme Process) Vit. A 4000 U., ketocholeonic acids 62 mg./Tab. Bot. 100s, 250s.

Use: Vitamin A supplement and therapy.

ACHROMYCIN. (Lederle) Tetracycline HCl.

Intramuscular: (100 mg., 250 mg., buffered w/ascorbic acid) Vial. Box 1s, 100s.

Intravenous: (250 mg., 500 mg., buffered w/ascorbic acid) Vial.

Oint. (3%) Tube 0.5 oz. & 1 oz.

Oint. (Ophth.) 1% Tubes 1/8 oz.

Ophth. Susp. 1%, Plastic dropper bot. 4 ml.

ACHROMYCIN-V. (Lederle) Tetracycline HCl. Cap.: 250

mg. Bot. 100s, 1000s, Unit dose 10 $\times$ 10s, Unit of use 50 $\times$ 28s; 500 mg. Bot. 100s, 1000s; Unit of use 50 $\times$ 20s; Unit dose 10 $\times$ 10s. **Oral Susp.:**

125 mg./5 cc. w/methylparaben 0.12%,

propylparaben 0.03%. Bot. 2 oz., 1 pt.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria.

ACID ACRIFLAVINE.

See: Acriflavine Hydrochloride (Various Mfr.)

ACID CITRATE DEXTROSE SOLUTION.

See: A.C.D. Sol. (Various Mfr.)

ACID CITRATE DEXTROSE ANTICOAGULANT SOLUTION MODIFIED.

See: A-C-D Solution Modified (Squibb)

ACID HISTAMINE PHOSPHATE.

See: Histamine Phosphate. (Various Mfr.)

ACID MANTLE CREME. (Dorsey) Aluminum acetate in specially prep. water-soluble hydrophilic cream at pH 4.2. Tube 1 oz., Jar 4 oz., 1 lb.

Use: Skin prophylactic agent in restoring normal skin acidity.

ACID MANTLE LOTION. (Dorsey) Aluminum acetate lotion. pH 4.5 Bot. 4 oz.

Use: Dermatitis.

ACIDOL.

See: Betaine Hydrochloride (Various Mfr.)

ACIDOPHILUS W/PECTIN. (Barth's) *Lactobacillus acidophilus* w/natural citrus pectin 100 mg./Cap. Bot. 100s.

Use: Aid to digestion.

ACID TRYPAFLAVINE.

See: Acriflavine Hydrochloride. (Various Mfr.)

ACIDULATED PHOSPHATE FLUORIDE. (Scherer) Flavored soln. containing 0.31% fluoride ion in 0.1 molar phosphate. Bot. 64 oz. Office Product.

Use: Dental caries prevention.

ACIDULIN. (Lilly) Glutamic acid HCl 340 mg./Pulvule. Bot. 100s, 1000s.

Use: Achlorhydria.

• **ACIFRAN.** USAN.

Use: Antihyperlipoproteinemic.

ACIGLUMIN.

See: Glutamic Acid Hydrochloride. (Various Mfr.)

ACI-JEL. (Ortho) Glacial acetic acid 0.921%, ricinoleic acid 0.70%, oxyquinoline sulfate 0.025%, glycerine 5%. Tube 85 Gm. w/dose applicator.

Use: Vaginal acidifier.

ACINITRAZOLE. B.A.N. 2-Acetamido-5-nitrothiazole, Aminitrozole (I.N.N.)

Use: Treatment of trichomoniasis.

See: Trichorad

Tritheon

• **ACLARUBICIN.** USAN.

Use: Antineoplastic.

ACLOBROM. Propantheline Bromide, B.A.N.

• **ACLOMETASONE DIPROPIONATE,** U.S.P. XXI.
Cream, oint., U.S.P. XXI.

A.C.N. (Person & Covey) Vitamin A palmitate 25M I.U., ascorbic acid 250 mg., niacinamide 25 mg./Tab. Bot. 100s.

Use: Dietary supplement.

ACNAVEEN. (CooperCare)

See: Aveenobar Medicated (CooperCare)

ACNA-VITE. (Cenci) Vit. A 10,000 I.U., C 250 mg., hesperidin 50 mg., niacinamide 25 mg./Cap. Bot. 75s.

Use: Acne vulgaris with Vit. A & C deficiency.

ACNE-5. (Goldline) Benzoyl peroxide 5%. Bot. 1 oz.

Use: Acne treatment.

ACNE-10. (Goldline) Benzoyl peroxide 10%. Bot. 1 oz.

Use: Acne treatment.

ACNE-AID CLEANSING BAR. (Stiefel) Neutral soap of 6.3% surfactant blend. Non-medicated. Bar 4.0 oz., 5.8 oz.

Use: Cleanse oily skin and acne.

ACNE-AID CREAM. (Stiefel) Benzoyl peroxide 10% in special cover-up base. Tube, 1.8 oz.

Use: Absorbs oil, hides blemishes.

ACNE-AID LOTION. (Stiefel) Sulfur 10%, alcohol 10% in a washable base. Clear or Tinted Bot. plastic 2 oz.

Use: Treatment of acne.

ACNEDERM. (Lannett) **Lotion:** Dispersable sulfur 5%, zinc sulfate 1%, zinc oxide 10%, alcohol 21%. Bot. 2 oz. **Soap:** Zinc oxide 2%, zinc sulfate 1%, colloidal sulfur 5%. Cake 3.2 oz.

Use: Skin conditions.

ACNE-DOME CREAM & LOTION. (Miles Pharm) **Regu-**lar: sulfur 4%, resorcinol monoacetate 3%, in a vehicle buffered to the pH range of normal skin.

Cream: Tube 1 oz. **Lotion:** Bot. 2 oz.

Use: Treatment of acne.

ACNE-DOME MEDICATED CLEANSER. (Miles Pharm) Sulfur 2%, salicylic acid 2% in a colloidal soybean complex. Medicated w/sponge 4 oz.

Use: Medicated skin cleanser.

ACNE LOTION. (Weeks & Leo) Benzoyl peroxide 10% in odorless, greaseless, vanishing lotion. Bot. 2 oz.

Use: Treatment for acne.

ACNO CLEANSER. (Baker/Cummins) Isopropyl alcohol 60%. Bot. 8 oz.

Use: Antiseptic astringent for the treatment of acne.

ACNO LOTION. (Baker/Cummins) Micronized sulfur 3%, salicylic acid 2%. Bot. 4 oz.

Use: Treatment of acne.

ACNOMEL. (Menley & James) Resorcinol 2%, sulfur 8%, alcohol 11%.

Cream. Tube 1 oz.

Use: Acne therapy.

ACNOPHILL. (Torch) Sulfur 5%, Pot. and zinc sulfides and polysulfides 5%, zinc oxide 10% in hydrophilic vehicle. Plain or tinted. Jar 1 lb. and 2 oz.

Use: Acne treatment

ACNOTEX. (C & M Pharmacal) Sulfur 8%, salicylic acid 2.25% in lotion base. Bot. 2 oz.

Use: Anti-acne cover-up lotion.

• **ACODAZOLE HYDROCHLORIDE.** USAN.

Use: Antineoplastic.

ACOTUS. (Whorton) Phenylephrine HCl 5 mg., guaiaicol glyceryl ether 100 mg., menthol 1 mg., alcohol, by vol. 10%/5 cc. Bot. 4 oz., 12 oz., gal.

Use: Colds, coughs.

ACR. (Western Research) Ammonium chloride 7.5 gr./Tab. Handicount 28s (36 bags of 28 tab.)

Use: Diuretic.

ACRIFLAVINE. Euflavine, Gonacrine, Neutroflavine Acriflavine neutral; a mixture of 2, 8-diamino-10-methylacridinium chloride & 2,8-diaminoacridine. (Lilly) Tab. 1.5 gr. Bot. 100s.

Use: External antiseptic.

ACRIFLAVINE HYDROCHLORIDE. (Various Mfr.) Hydrochloride form of acriflavine. Acid acriflavine, acid tryptaflavine, flavine, tryptaflavine.

National Aniline—Pow., Bot. (1, 5, 10, 25, & 50 Gm.). Tab. (1.5 gr.) Bot. 50s & 100s.

Use: External bacteriostatic.

- **ACRISORCIN**, U.S.P. XXI. Cream U.S.P. XXI. 9-Aminoacridinium 4-hexylresorcinolate.
See: Akrinol Cream (Schering)
Use: Topical, tinea versicolor. Local antifungal.
- **ACRONINE**, USAN. 3, 12-Dihydro-6-methoxy-3,3,12-trimethyl-7H-pyrano[2,3,-c]acridin-7-one. Under study.
Use: Antineoplastic.
- A-C TABLET**. (Century) Aspirin 6 gr., caffeine 0.5 gr./Tab.
Bot. 100s, 1000s
Use: Analgesic.
- ACTACIN-C SYRUP**. (Vangard) Codeine phosphate 10 mg., triprolidine HCl 2 mg., pseudoephedrine HCl 20 mg., guaifenesin 100 mg./ 5 ml. Bot. pt., gal.
Use: Antihistamine, decongestant, cough depressant.
- ACTACIN SYRUP**. (Vangard) Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./ 5 ml. Bot. pt., gal.
Use: Antihistamine, decongestant.
- ACTACIN TABLETS**. (Vangard) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg./Tab. Bot. 100s, 1000s.
Use: Antihistamine/decongestant.
- ACTAGEN**. (Generix) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg. Tab. Bot. 100s, 1000s.
Use: Symptomatic treatment of allergic rhinitis.
- ACTAGEN-C COUGH SYRUP**. (Goldline) Triprolidine HCl, pseudoephedrine HCl, codeine phosphate/Bot. 4 oz., pt., gal.
Use: Decongestant, cough suppressor.
- ACTAGEN TABLETS**. (Goldline) Triprolidine HCl, pseudoephedrine HCl. Bot. 100s, 1000s.
Use: Decongestant.
- ACTAL SUSPENSION**. (Winthrop Products) Aluminum hydroxide.
Use: Antacid.
- ACTAL PLUS TABLETS**. (Winthrop Products) Aluminum hydroxide, magnesium hydroxide.
Use: Antacid, demulcent.
- ACTAL TABLETS**. (Winthrop Products) Aluminum hydroxide.
Use: Antacid.
- ACTAMIN**. (Buffington) Acetaminophen 325 mg./Tab. Sugar, lactose, and salt free. Dispens-A-Kit: 500s. Aidpaks 100s. Unit boxes 20s. Medipaks 200s.
Use: Analgesic, anti-pyretic.
- ACTAMIN EXTRA**. (Buffington) Acetaminophen 500 mg./Tab. Sugar, lactose, and salt free. Dispens-A-Kit 500s. Medipaks 200s.
Use: Extra strength nonaspirin analgesic and antipyretic.
- ACTAMIN SUPER**. (Buffington) Acetaminophen 500 mg. Tab. w/caffeine. Sugar, salt, and lactose free. Dispens-A-Kits 500s, Medipaks 200s.
Use: Maximum strength analgesic, antipyretic.
- **ACTAPLANIN**, USAN.
Use: Growth stimulant.
- ACTH-ACTEST GEL** (Fellows) Repository corticotropin 40 or 80 u./cc. Vial 5 cc.
- ACTH**. (Adrenocorticotropin hormone. Adrenocorticotrop(h)in).
See: Corticotropin, U.S.P.
Mallard—(40 u./ml., 5 ml.)
Forest—(40 u. or 80 u./ml., 5 ml.)
Parke, Davis—25 units/Vial; 40 units/Vial.
Pharmex—(40 u. or 80 u./ml., 5 ml.)
- ACTH GEL, PURIFIED**. Arcum.—40 or 80 units/ml., vial 5 ml. Conal.—40 or 80 units/ml., vial 5 ml. Hart Labs.—40 units/ml., vial 5 ml. Bowman—Adrenocorticotropin hormone 40 u., aqueous gelatin 16%, phenol 0.5%/cc. Vial 5 cc.
Use: Repository corticotropin.
Arcum—(40 u. or 80 u./ml., 5 ml.)
Bell—(40 u. or 80 u./ml., 5 ml.)
Bowman—(40 u./ml., 5 ml.)
Hyrex—(40 u. or 80 u./ml., 5 ml.)
Jenkins—(40 u. or 80 u./ml., 5 ml.)
Vortech—40 units/ml., 80 units/ml., Vial 5 ml.
Rocky Mtn. Pharmacal—40 units/ml., 80 units/ml. Vial 5 ml.
Wesley Pharm.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.
Wyeth—(40 u. or 80 u./ml. or Tubex)
- ACTHAR**. (Armour) Corticotropin for Injection. Vial 25, 40 u./Vial. (Lyophilized w/gelatin). Pkg. 10s.
Use: I.M., S.C. or I.V. stimulation of adrenal cortex and diseases related to it.
- ACTICORT LOTION 100**. (Baker/Cummins) Hydrocortisone alcohol 1% in a specially formulated vehicle. Bot. 2 oz.
Use: Anti-inflammatory & antipruritic.
- ACTIDIL** (Burroughs Wellcome) Triprolidine HCl Tab. 2.5 mg./Tab. Bot. 100s. Syr. 1.25 mg./5 ml. Bot. 1 pt.
Use: Antihistamine.
- ACTI-DIONE**. (Upjohn) Cycloheximide. (3-[2-(3,5-Dimethyl-2-oxocyclohexyl)-2-hydroxyethyl]-glutarimide) 85-100% w/w.
Not for medicinal use. For professional use only.
Bot. 4 Gm.
- ACTIDOSE**. (Paddock) Activated charcoal solution 25 Gm./120 ml. or 50 Gm./240 ml.
Use: Orally or through gastric tube to adsorb toxins in GI tract in drug overdose or poisoning emergency.
- ACTIDOSE-AQUA**. (Paddock) Activated charcoal aqueous suspension 25 Gm./120 ml. or 50 Gm./240 ml.
Use: Orally or through gastric tube to adsorb toxins in GI tract in drug overdose or poisoning.
- ACTIFED**. (Burroughs Wellcome) Tab.: Triprolidine HCl. 2.5 mg., pseudoephedrine HCl. 60 mg./Tab. Bot. 30s, 100s, 1000s. Boxes 12s, 24s. Unit dose pack 100s. Unit of use 90s. Ctn. 24s. Cap.: Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg./Cap. Box 10s, 20s. Syr.: Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./5 ml. Bot. 4 oz., 1 pt., gal.
Use: Common cold, allergic rhinitis.
- ACTIFED CAPSULES**. (Burroughs Wellcome) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg./Cap. Box 10s, 20s.
Use: Nasal decongestant, antihistamine.

ACTIFED WITH CODEINE COUGH SYRUP.

(Burroughs Wellcome) Codeine phosphate 10 mg., triprolidine HCl 2 mg., pseudoephedrine HCl 30 mg., dextromethorphan 100 mg./5 ml. Bot. pt., gal.

Use: Treatment of the common cold.

ACTINOMYCIN. B.A.N. Antimicrobial substances with antitumor activity produced by *Streptomyces* antibioticus and *Streptomyces chrysomallus*. Sanamycin is Actinomycin C.

ACTINOMYCIN C. Name previously used for Cactinomycin.

ACTINOMYCIN D. Dactinomycin, U.S.P. XXI.

See: Cosmegen (Merck Sharp & Dohme)

• **ACTINOQUINOL SODIUM.** USAN.

Use: Ultraviolet screen.

ACTINOSPECTOCIN. Name previously used for Spectinomycin.

ACTIVATED ATTAPULGITE.

W/Aluminum hydroxide, magnesium carbonate coprecipitate, compressed gel.

See: Hykasil, Cream (Philips Roxane)

W/Polysorbate 80, colloidal sulfur, salicylic acid, propylene glycol.

See: Sebasorb Lotion (Summers Labs.)

ACTIVATED CHARCOAL TABLETS. (Cowley) 5 gr./Tab. Bot. 1000s.

ACTIVATED 7-DEHYDROCHOLESTEROL

See: Vitamin D-3 (Various Mfr.)

ACTIVATED ERGOSTEROL

See: Calciferol.

ACTIVE LIFE CLOSED OSTOMY POUCH.

(ConvaTec) One-piece closed-end pouch with skin barrier, picture frame support adhesive, odor proof pouch.

ACTIVE LIFE OPEN OSTOMY POUCH. (ConvaTec)

One-piece disposable ostomy appliance with stomahesive peristomal skin covering and adhesive support collar. Box 10s.

ACTOL EXPECTORANT. (Beecham Labs) Noscapine

30 mg., guaifenesin 200 mg./5 ml. or Tab. Syr.

W/alcohol 12.5%. Bot. pt. Tab. Bot. 100s.

Use: Treatment of symptoms associated with common cold.

ACTOMOL. Mebanazine, B.A.N.

ACTOQUINOL SODIUM. 8-Ethoxy-5-quinoline-sulfonic acid sodium salt.

Use: Ultraviolet screen.

ACTOS 50. Corticotrophin, B.A.N.

ACTRAPID HUMAN INSULIN. (Squibb Novo) Human insulin 100 Units/ml. Vial 10 ml.

Use: Human insulin (rapid action).

ACTRIOL. Epicoestriol, B.A.N.

ACTYLATE. (Kinney) Ammonium salicylate 80 mg., pot. salicylate 80 mg., strontium salicylate 80 mg., pot. para-aminobenzoate 0.32 Gm., ascorbic acid 20 mg./Tab. Bot. 100s.

Use: Rheumatoid arthritis.

ACUCRON. (Seatrace) Acetaminophen 300 mg., salicylamide 200 mg., phenyltoloxamine 20 mg./Tab. Bot. 100s, 1000s, 5000s.

Use: Antihistaminic, analgesic.

• **ACYCLOVIR.** USAN. 9-(2-hydroxyethoxymethyl)-guanine.

Use: Management of initial herpes genitalis and mucocutaneous herpes simplex virus infections.

See: Zovirax Caps., Oint. (Burroughs Wellcome)

• **ACYCOLOVIR SODIUM.** USAN.

Use: Antiviral.

ACYLPYRIN.

See: Aspirin

ADALAN LANATABS. (Lannett) Amobarbital 50 mg., homatropine methylbromide 7.5 mg., methamphetamine HCl 10 mg./T.D. Tab. Bot. 100s, 1000s.

ADAMANTANAMINE HCl. Amantadine HCl. Anti-flu capsule. This drug is thought to protect cells against entry of the flu virus without actually destroying the virus. Also used in the treatment of Parkinson's disease.

See: Symmetrel, Cap., Syr. (Du Pont)

ADANON. Methadone, B.A.N.

ADAPETTES. (Alcon) EDTA 0.1%, chimerosal 0.004%. Pkg. 15 ml.

Use: Contact lens rewetting and lubricating solution.

ADAPIN. (Pennwalt) Doxepin HCl 10 mg., 25 mg. 50 mg., 75 mg., 100 mg./Cap. Bot. 100s, 1000s. 25 mg., 50 mg./Cap. also available in 5000s.

Use: Antianxiety, antidepressant.

ADAPT. (Alcon) Povidone, EDTA 0.1%, thimerosal 0.004%. Bot. 15 ml.

Use: Contact lens cushioning solution.

ADATUS. (Merchant) Hydrocodone bitartrate 5 mg., guaifenesin 100 mg., alcohol 5%/5 ml. Syr. Bot. pt., gal.

Use: Cough relief.

A.D.C. DROPS. (CMC) Vit. A 3000 I.U., D 400 I.U., C 60 mg./0.6 cc. Bot. 60 cc.

ADCCORTYL. Triamcinolone, B.A.N.

ADEECON. (CMC) Vit. A 5000 I.U., D 1000 I.U./Cap. Bot. 1000s.

Use: Supplement.

ADEFLOL CHEWABLE. (Upjohn) Fluoride 1 mg. & 0.5 mg., Vitamin A 4000 I.U., D 400 I.U., C 75 mg., thiamine monitrate (B-1) 2 mg., B-2 2 mg., niacinamide 18 mg., B-6 1 mg., calcium pantothenate 5 mg., B-12 2 mcg./0.5 or 1 mg. Chewable Tab. Bot. 100s, 500s.

Use: Prevention of dental caries.

ADEFLOL DROPS. (Upjohn) Vit. A 2000 I.U., D 400 I.U., C 50 mg., B-6 1 mg., fluoride (as sodium fluoride) 0.5 mg./0.6 ml. Bot. 50 ml. w/dropper.

Use: Prevention of dental caries.

ADEFLOL M TABLETS. (Upjohn) Vit. A 6000 I.U., D 400 I.U., B-1 1.5 mg., B-2 2.5 mg., C 100 mg., niacinamide 20 mg., Vit. B-6 10 mg., calcium pantothenate 10 mg., Vit. B-12 2 mcg., folic acid 0.1 mg., fluoride (as sodium fluoride) 1 mg., calcium (as calcium carbonate) 250 mg., iron 30 mg./Tab. Bot. 100s, 500s.

Use: Dietary supplement.

• **ADENINE.** U.S.P. XXI. 6-Aminopurine.

ADENO TWELVE GEL INJECTION. (Fellows)

Adenosine-5-monophosphate 25 mg., methionine 25 mg., niacin 10 mg./cc. Vial 10 cc.

ADENOCREST. (Nutrition) Adenosine-5-monophosphate sod. 25 mg., nicotinic acid as sod. salt 25 mg., gelatin 100 mg. w/benzyl alcohol 1.5%/cc. Vials 10 cc.

Use: Capillary and arterial vasodilation.

ADENOLIN FORTE. (Lincoln) Adenosine-5-monophosphate, 25 mg., methionine 25 mg., niacin 10 mg./cc. Vial 15 cc.

Use: Arthritis, bursitis, tendinitis & other degenerative diseases.

• **ADENOSINE.** F.D.A. 6-Amino-9-beta-D-Ribofuranosyl-9H-purine.

ADENOSINE IN GELATIN. (Fellows) Forte:

Adenosine-5-monophosphate 50 mg./cc. Super:

Adenosine-5-monophosphate 100 mg./cc.

ADENOSINE-5-MONOPHOSPHATE AS THE SODIUM SALT.

See: Adenocrest, Amp. (Nutrition)

• **ADENOSINE PHOSPHATE.** USAN. 5-Adenylic acid. B.A.N. Adenosine-5'-(dihydrogen phosphate). Adenosine monophosphate, AMP.

Use: Nutrient.

See: Adenyl

Cobalazine, Inj. (Keene)

My-B-Den (Miles Pharm)

ADENYL. Adenosine Phosphate, B.A.N.

ADEPSINE OIL.

See: Petrolatum Liquid (Various Mfr.)

ADICILLIN. B.A.N. 6-[D(+)-5-Amino-5-carboxy-valeramide]penicillanic acid. Penicillin N.

Use: Antibiotic.

• **ADINAZOLAM.** USAN.

Use: Antidepressant, sedative.

• **ADINAZOLAM MESYLATE.** USAN.

Use: Antidepressant, sedative.

ADIPEX-P. CAPSULES. (Lemmon) Phentermine HCl 37.5 mg./Cap. Bot. 100s, 400s.

Use: Anorectic agent.

ADIPEX-P TABLETS. (Lemmon) Phentermine HCl 37.5 mg./Tab. Bot. 100s, 400s, 1000s.

Use: Anorectic agent.

• **ADIPHENINE HYDROCHLORIDE.** USAN. (2-Diethylaminoethyl diphenylacetate hydrochloride.) Pow. 1 lb. for mfg. use.

Use: Antispasmodic.

ADIPIODONE (I.N.N.). Iodipamide, B.A.N.

3,3'-(ADIPOYLDIIMINO)-BIS[2,4,6-TRI-IODOBENZOIC ACID]. Iodipamide, U.S.P. XXI.

ADIPOST. (Ascher) Phendimetrazine tartrate 105 mg./Slow-rel. Cap. Bot. 100s.

Use: Appetite suppressant.

ADISOL TAB. (Major) Disulfiram 250 mg./Tab. Bot. 100s. 500 mg./Tab. Bot. 50s.

Use: Treatment of alcoholism.

ADLERIKA. (Last) Magnesium sulfate 4.0 Gm./15 ml. Bot. 12 oz.

Use: Laxative.

ADONIDINE. (City Chem.) Bot. 1 Gm.

Use: Cardiac stimulant.

ADPHEN TABLETS. (Ferndale) Phendimetrazine tartrate 35 mg./Tab. Bot. 100s.

Use: Anorexiatic.

A-D-R. (Lincoln) dl-Methionine 250 mg., corn starch lubricant 42 mg./Cap. Bot. 100s, 500s.

Use: Amino acid supplement.

ADRENALIN(E).

See: Epinephrine. (Various Mfr.)

ADRENALIN CHLORIDE SOLUTION. (Parke-Davis)

Epinephrine HCl. Principle of the medullary portion of suprarenal glands. Amp.: 1:1000-1 ml. Epinephrine

1 mg./ml. with not more than 0.1% sodium bisulfite as preservative. Amps. 10s. Steri-Vial 1:1000-1 ml.

Epinephrine in isotonic sodium chloride solution with 0.5% chlorobutanol and not more than 0.15% sodium bisulfite as preservative. Vial 30 ml. Soln.

1:1000-1 ml. Screw cap Bot. 1 oz. (Same desc. as Steri-Vial). Soln. 1:100. Each 100 ml. contains 1 Gm.

epinephrine HCl dissolved in sodium chloride-citrate buffer solution with Phemerol chloride 0.2 mg./ml.

and sodium bisulfite 0.2% as preservative. Bot. 0.25 oz.

Use: Sympathomimetic agent.

ADRENALINE HYDROCHLORIDE.

See: Epinephrine Hydrochloride. (Various Mfr.)

• **ADRENALONE.** USAN. 3',4'-Dihydroxy-2-(methylamino)-acetophenone.

Use: Adrenergic (ophthalmic).

ADRENAMINE.

See: Epinephrine (Various Mfr.)

ADRENERGIC AGENTS.

See: Sympathomimetic agents

ADRENERGIC-BLOCKING AGENTS.

See: Sympatholytic agents

ADRENINE.

See: Epinephrine. (Various Mfr.)

ADRENOCHROME.

See: Carbazochrome Salicylate.

ADRENOCHROMAZONE.

See: Carbazochrome Salicylate.

ADRENOCHROME MONOSEMICARBAZONE.

See: Carbazochrome Salicylate

ADRENOCORTICOTROPIC HORMONE. ACTH acts

by stimulating the endogenous production of cortisone and the final result is similar whichever substance is administered. The nature of action of

each is unknown. Both are necessary in the metabolism of carbohydrates, protein & fat & exert

profound effects on neuromuscular metabolism.

See: ACTH gel, Inj. (Bowman)

Corticotropin, U.S.P.

Cortigel (Savage)

ADRENOMIST INHALANT AND NEBULIZERS.

(Nephron) Epinephrine 1%, Bot. 0.5 oz., 1.25 oz.

Use: Inhalant for bronchial asthma.

ADRENUCLEO. (Enzyme Process) Vit. C 250 mg., d-cal. pantothenate 12.5 mg., bioflavonoids 62.5 mg./Tab. Bot. 100s, 250s.

Use: Vitamin-C supplement therapy.

ADRIAMYCIN. (Adria) Doxorubicin HCl 10 mg., lactose 50 mg./Vial Pkg. 10s. Doxorubicin HCl 50 mg., lactose 250 mg./Vial Ctn. 1s.

Use: Antineoplastic.

ADRIAN TABS. (Major) Nylidrin 6 mg. or 12 mg./Tab. Bot. 100s, 1000s.

Use: Vasodilator.