Clinical Pharmacology in Pregnancy

Fundamentals and Rational Pharmacotherapy

Helmut P. Kuemmerle, M.D., F.R.S.M.

Clinical Pharmacology in **Pregnancy**

Fundamentals and Rational Pharmacotherapy

Helmut P. Kuemmerle, M.D., F.R.S.M.


Specialist in Obstetrics and
Gynecology, Clinical Pharmacology
and Chemotherapy
Visiting Professor; Tokyo Medical
College, Tokyo
Visiting Professor; Clinical
Pharmacology, Hanyang University
College of Medicine, Seoul

Co-editor:

Klaus Brendel, Ph.D.

Department of Pharmacology and
Toxicology
University of Arizona Health Sciences
Center
Tucson, Arizona





Characteristica and an armore in the control of

Publisher: Thieme-Stratton Inc. 381 Park Avenue South New York, New York 10016

Cover design by M. Losaw

Clinical Pharmacology in Pregnancy Edited by Helmut P. Kuemmerle

0-86577-074-3 (Thieme-Stratton Inc.) 3-13-627501-2 (Georg Thieme Verlag)

Copyright © 1984 by Thieme-Stratton Inc. All rights reserved. No part of this publication may be reused without the express written consent of the publisher. Library of Congress Catalog Card Number 82-51140.

Preface

Clinical pharmacology is the theory of the rational formation of judgments in pharmacotherapy. In other words, it is concerned with experimental and ap-

plied clinical testing of therapeutic effects on human subjects.

Pregnancy is a particular phase of life that requires special attention in every respect, especially from the clinico-pharmacologic point of view. The maternal-placental-fetal unit, the triad, has to be understood as a biologic, pharmaco-kinetic, and pharmacodynamic unit. Consequently the concern of the mother cannot be dealt with alone, but rather both mother and embryo/fetus have to be included in all prophylactic and therapeutic considerations.

The principal target of drugs administered in pregnancy is the mother; the fetus is essentially an unwanted recipient. This feature will be sustained throughout pregnancy up into the early neonatal life, as the possibility for the drug transfer via breast milk in the nursing mother remains. The risk/benefit ratio has to be considered in every respect in pregnancy because fetal effects primarily of a toxic or negative nature have become a major problem in pregnancy-associated

therapeutics.

The concern of this book is to consider the current status and the actual role of clinical pharmacology in pregnancy for the rational use and for the evaluation of drug prophylaxis and drug therapy in pregnancy. It should be considered as a basic constituent for today's drug development, drug prophylaxis, drug ther-

apy, and drug safety in pregnancy.

All aspects and principles that are important for the rational use of drugs in pregnancy are discussed and interpreted from the fundamental up to the practical use of the various drug classes. This book will be a reliable guide and a valuable source of information for all involved in prescribing and using drugs in pregnancy. It is a clinically oriented and well documented text providing all physicians with an indispensable guide to contemporary drug therapy in pregnancy available and known today.

Clinical Pharmacology in Pregnancy brings together the knowledge, experience, and understanding of the fundamentals and principles of drugs and pharmacology.

macotherapy of 29 eminent international contributors.

We are very grateful to our publisher, especially Dr. Günther Hauff, Mr. Achim Menge, Mrs. Jill Rudansky, and Mr. James Costello for their good cooperation as well as for their patience.

Finally, we wish to thank Mrs. Kuemmerle for her excellent assistance in the

coordination of the work.

H. P. Kuemmerle K. Brendel

Contributing Authors

Ingrid Amon, M.D.

Professor of Clinical Pharmacology Institute of Clinical Pharmacology University of Greifswald Greifswald, German Democratic Republic

William Banner, Jr., M.D., Ph.D.
Professor of Pediatrics
Department of Pediatrics
University of Arizona Health Sciences
Center

Tucson, Arizona

Klaus Brendel, Ph.D.

Professor of Pharmacology and
Toxicology
Department of Pharmacology
University of Arizona Health Sciences
Center
Tucson, Arizona

Erwin Deutsch Dr.jur.

Professor of Law
Forschungsstelle für Arzt-und
Arzneimittelrecht des Juristischen
Seminars der Universität
Federal Republic of Germany

Raymond C. Duhamel, Ph.D.

Department of Pharmacology
University of Arizona Health Sciences
Center
Tucson, Arizona

D. F. Hawkins, M.D., Ph.D.
Professor of Gynecology and Obstetrics
Hammersmith Hospital
Institute of Obstetrics and Gynecology
London, United Kingdom

S. Hayashi, M.D.

Department of Obstetrics and Gynecology Kyoto University School of Medicine Tatsuno City, Japan

Keith Hillier, Ph.D.

University of Southampton Southampton, United Kingdom

G. Hitzenberger, M.D.

Professor of Clinical Pharmacology Department of Clinical Pharmacology First Medical University Clinic Vienna, Austria

Hansgeorg Hüller, M.D.

Professor of Clinical Pharmacology Institute of Clinical Pharmacology, Charite Humboldt University Berlin, German Democratic Republic

S. Katagiri, M.D.

Department of Obstetrics and Gynecology Hirosaki University School of Medicine 'Hirosaki, Japan

K. Kido, M.D.

Department of Obstetrics and Gynecology Kyoto University School of Medicine Tatsuno City, Japan

Hans Kopera, M.D.

Professor of Clinical Pharmacology Department of Experimental and Clinical Pharmacology University of Graz Graz, Austria

Helmut P. Knemmerle, M.D., F.R.S.M.

Prof. (vis.) of Clinical Pharmacology, Tokyo Medical College Prof. (vis.) of Clinical Pharmacology, Hanyang University College of Medicine, Seoul Specialist in Obstetrics and Gynecology, Clinical Pharmacology and Chemotherapy Bad Heilbrunn, Federal Republic of Germany .

G. Laudahn, M.D.

Professor of Internal Medicine Clinical Research Department Schering AG Berlin, Germany

Seiji Matsuda, M.D.

Professor of Gynecology and Obstetrics Kohto Hospital Juntendo University Tokyo, Japan

Teresita Mazzei, M.D.

Professor of Pharmacology Cattedra di Chemioterapia Univerita degli studi di Firenze Firenze, Italy

Domenico Pecorari, M.D.

Professor of Gynecology and Obstetrics Department of Obstetrics and Gynecology University of Trieste Ospedale Maggiore Trieste, Italy

Roger M. Pinder, Ph.D. CNS Research Organon International BV Oss. The Netherlands

D. Reinhardt, M.D.

Professor of Pediatrics University Children-Hospital A Duesseldorf, Federal Republic of Germany

O. Richter

Professor of Biometrics and Mathematics Institut für Landwirtschaftliche Betriebslehre Abteilung für Angewandte Statistik und Mathematik d. Universität Bonn Bonn, Federal Republic of Germany

W. A. Ritschel, Ph.D.

Professor of Pharmacokinetics University of Cincinnati Medical Center Cincinnati, Ohio

Roberto Serra, M.D.

Professor of Phthisiology Invited Professor University of Louvain Belgium

Shinryo Shinagawa, M.D.

Professor of Obstetrics and Gynecology Department of Obstetrics and Gynecology Hirosaki University School of Medicine Hirosaki, Japan

Hiroaki Soma, M.D.

Professor of Gynecology and Obstetrics Department of Obstetrics and Gynecology Tokyo Medical College Hospital Tokyo, Japan

Heini Stamm, M.D.

Professor of Gynecology and Obstetrics Clinic of Obstetrics and Gynecology Kantons-Hospital Baden, Switzerland

H. Tuchmann-Duplessis

Professor of Embryology Laboratory of Embryology Rene Descartes University Paris, France

Leo Wislicki, M.D. Professor of Gynecology and Obstetrics The Hebrew University Hadassah Medical School Jerusalem, Israel

Noritsume Yamada, M.D. Professor of Gynecology and Obstetrics Department of Obstetrics and Gynecology Kyoto University School of Medicine Tatsuno City, Japan

Contents

I:	Fundamentals	of	Drug	Therapy	in
	Pregnancy				

1.	Introduction	3
	H. P. Kuemmerle	
2.	Human Embryogenesis	5
	H. Tuchmann-Duplessis	
3.	The Maternal-Placental-Fetal Triad	21
	H. P. Kuemmerle	
4.	The Placental Morphological Structures and Their Meaning for Pharmacokinetics and Pharmacodynamics	23
	H. Soma	
5.	Structural Changes of the Placenta in Relation to Placental Transfer Alterations	29
	H. Soma	
6.	Normal and Abnormal Development of Human Embryos and Fetuses: Embryonic-Fetal Growth and Intrauterine Environment.	34
	D. Pecorari	
7.	Developmental Pharmacology and Toxicology	40
	H. Hüller and I. Amon	
8.	General Clinical Pharmacology of Diaplacental and Paraplacental Transfer	50
	H. P. Kuemmerle	
9.	Pharmacokinetic Characterization of Drugs	59
	W. A. Ritschel	

10.	Dose-Response Relationship As a Basis for Choice of Drugs in Pregnancy and for Dosage Calculation	64
	I. Amon and H. Hüller	
11.	Drug Metabolism in the Pregnant Woman	74
	I. Amon and H. Hüller	
12.	Drug Metabolism in the Fetus	87
٠.	K. Brendel and R. C. Duhamel	
13.	Transplacental Toxicology	101
14.	Teratogenesis in Humans	113
	D. F. Hawkins	
15.	Factors Affecting the Mode of Drug Action in Pregnancy.	124
	L. Wislicki	
16.	General Principles of Drug Therapy During Pregnancy	129
	H. P. Kuemmerle	
17.	General Principles of Clinical Pharmacology of the Fetus.	132
	W. Banner, Jr.	
18.	General Principles of Clinicopharmacological Evaluation of New Agents in Pregnancy	143
	H. P. Kuemmerle	
19.	Clinical Pharmacology of the Pregnant Human Uterus	145
	D. F. Hawkins and K. Hillier	
20.	Epidemiology of Drug-Induced Effects on the Embryo and Fetus	158
	T Marzei	

21.	Drugs and Maternal Death	168
	S. Shinagawa and S. Katagiri	
22.	Ethical and Medical Considerations in the Use of Drugs in Pregnancy	174
* .	D. F. Hawkins	
23.	Legal and Juridical Considerations in Administering Drugs in Pregnancy	179
	E. Deutsch	
	pplied Clinical Pharmacology in regnancy	
	Lag of the original state of the control of the con	
24.	The Real Need for Drug Prophylaxis and Therapy During Pregnancy	185
	H. Stamm	
25.	Rationale of Drug Therapy in Pregnancy	197
	N. Yamada, K. Kido, and S. Hayashi	
26a.	Pharmacokinetics of Transplacental Passage	210
	H. P. Kuemmerle	
26b.	Kinetic Model and Dosage Regimen	217
	x9nommoli x7r	
27.	i	229
	D. F. Hawkins	
28.	Embryotoxic Drugs	235
	K. Brendel and R. C. Duhamel	
29.	Translactal Passage and Pharmacotherapy of Lactation	252
	D. Pecorari	
30.	Pharmacokinetics of Translactal Passage	260
	D. Reinhardt and O. Richter	

Appe	Drug Classes. Drug Groups—Subclassification	
	The state of the s	
31.	Chemotherapeutic Agents	270
	T Mazzei	
	Legal and Legalical considerations in the first to Administrative Driver Charge space.	
32.		274
	S. Matsuda	
33.	Hormones of the Pituitary Gland	. 281
	H. Stamm	
3.4	Thyroid and Parathyroid Hormones;	
34.	Thyrostatic Drugs	. 285
	H. P. Kuemmerle	
2 =		205
35.	Adrenal Corticoid Hormones	
1	I. Amon	
36.	Catecholamines	. 290
	H. Stamm	
	The Property of the Contract o	
37.	Hormones of the Pancreas	
	H. P. Kuemmerle	
38.	Sex Hormones	294
30.	G. Laudahn	
	G. Laudani	
39.	Cardiovascular—Renal Drugs	. 312
	G. Hitzenberger	
6.	and the second s	
40.	Beta-Adrenergic Stimulants	. 325
	H. Stamm	
41	Drugs Acting on the Respiratory System	
41.	R. Serra	. 32(
42	Vitamins	331
44.	H P Knommerle	. 334

此为试读,需要完整PDF请访问: www.ertongbook.com

43.	Non-narcotic Analgesics and Antipyretics, and Nonsteroidal Antirheumatics	333
	L. Wislicki	
44.	Anticonvulsants	337
	S. Matsuda	
45.	Stimulants of the Central Nervous System	339
	R. Pinder and H. Kopera	
85		
46.	Hypnotics and Sedatives	342
	D. Pecorari	
	The second secon	
47.	Psychotomimetic and Psychotherapeutic Drugs	345
	H. Kopera and R. Pinder	
48.	Anesthetics, Including Local Anesthetics	352
	D. Pecorari	
49.	Cholinergic Drugs	355
	H. Stamm	
=0		
50.	Anticholinergic Drugs	357
	I. Amon	
F1		3.50
51.	Alkaloids	359
-	I. Amon	
52	Antiemetics	262
54.		303
	H. Stamm	
53	Laxatives	265
55.		302
	H. Stamm	
54	Immunologic Agents	367
J4.	I. Amon	307
	1. Amon	
55.	Diagnostic Agents and Agents Used in	
200	Roentgenography, Including Radioactive Agents	370
	U D Vuommarlo	

Com	ients .	
56.	Blood and Antianemic Agents	372
	D. Pecorari	
57.	Coagulants and Anticoagulants	375
	H. Stamm	
58.	Stomachics and Antacids	378
	D. Pecorari	
59.	Miscellaneous Chemical Agents Affecting Pregnancy	380
	H. P. Kuemmerle	
	Index	381

Fundamentals of Drug Therapy in Pregnancy

committed pharmer recovery as the form of the control of the control of graph means and pharmer control of the control of the

viet on a material factor, represent to the solid form of the solid factor of the soli

efection of m. mark is to be destrout translation to a succession of the modern effective and the succession of the succ

the first of the continues like the continues of the cont

en estate opposit 20 destroon yn head en heed op destroop op 100 yn generalen. De den de gegende et de heed op de heed en heed de heed

official section of the section of t

Program of the

Introduction

H. P. Kuemmerle

Introductory Remarks on the Problem Complex

For pharmacotherapy (and thus also for clinical drug trials) pregnancy is a special phase of life that, in general and in particular, differs from all others on medical, ethical, and, above all, legal grounds. However brutally, the wellknown drug scandals (Thalidomide, Menocil) have made the public acutely aware of the complex problem. Still, it is once again evident how briefly public attention focuses on such events. The term drug safety was initially taken very seriously, but in recent times it has become of symbolic importance only. The problem has not yet been completely grasped in all its sociopolitical ramifications and in the light of its inevitable and serious consequences. To best protect and preserve each individual from questionable treatment, harm, and exposure to arbitrary decisions, the concept of drug safety must be included among genuine sociopolitical necessities.

In this context, the human being must be seen not only as a biologic but also as a social unit, although the various factors cannot always be isolated. Within the physician's realm of responsibility there are few opportunities to counter harmful influences that arise exogenously, except by means of chemical substances—i.e., drugs. In the vast majority of cases this is the only way to affect such influences. This is particularly true for all pharmacotherapeutic measures and all clinical drug

trials during the childbearing years.

In the early stages the physician cannot be sure whether a woman of childbearing age is pregnant or not. Precisely because of this, the strictest standards must be observed during this life phase. If medication is required, it must be given only under conditions of the greatest precision and expertise, and always with critical distance. However, it would cer-

tainly be a mistake to assume an attitude of therapeutic nihilism because of these strictures. A good rule of thumb ought to be "as few drugs as possible," and those few drugs should be tried and tested ones, never new substances, nor combinations whose characteristic effects are little or not at all known.

The maternal-placental-fetal triad must be understood as a biologic, pharmacokmetic, and pharmacodynamic unit. Consequently, one may not deal with the concerns of the mother alone, but rather mother and child have to be included simultaneously in all therapeutic considerations.

As regards the clinicopharmacologic or clinicotherapeutic aspects of a drug, the embryo or fetus has a uniquely specific sensitivity that is not at all omparable with that of the adult. Since adult dosage must be administered to the mother, serious and often irreversible side effects and damage may become manifest in the embryo or fetus, or in the newborn. Thus, any risk of possible injury to the embryo or fetus must be judged too great.

Of course, one exception can be considered permissible: namely, new drugs that are specifically developed for therapeutic use in gravidas and have a predictable chance of success. This exception, however, is only permissible if under two conditions: (a) All legal provisions and the guidelines laid down by professional medical organizations and internationally recognized organizations (e.g., the Declaration of Helsinki, WHO); must be adhered to, and (b) preclinical data that are complete according to the current state of knowledge must be available. If there is the slightest suspicion of possible harm, the trials should be immediately terminated and not resumed until findings from new, nonobjectionable animal experiments approved by an unbiased committee of experts are available.

All those involved in clinical investigations,

whether clinical pharmacologists or specialists, whether engaged in clinical medicine, general medicine, or industry, when participating in special investigations in women of childbearing age and during pregnancy, must let their actions and thinking be guided by the keenest sense of responsibility and duty to conscientiousness. They must foreswear all avant-garde, careerist, and egoistic motiva-

In the maternal-placental-fetal unit the interests of the embryo or fetus and mother can be diametrically opposed, i.e., drug concentrations that are harmless for the mother can clearly influence or damage the embryo or fetus, and vice versa. There is still no completely reliable way known, on the basis of animal experiments, to preclude teratogenic effects, much less to exclude them. Retrospective investigations on humans are the only way to ensure relative innocuousness.

A plethora of questions arises from these considerations. The decisive question howadays is no longer whether a drug can or cannot be transferred dia- or paraplacentally, but rather: How rapidly does this passage take place? What concentrations are reached in the embryo or fetus? Is the concentration changed or is the drug even totally destroyed during its passage into the placental membrane? The most important question to each physician is: What changes can the drug cause in the embrvo or fetus?

The time of initiation and the duration of drug therapy are directly related to the injury triggered. Also of importance are the pharmacokinetic and pharmacodynamic qualities of the substance itself. Special attention must be paid to the fact that the placental membrane behaves selectively toward compounds that are closely related structurally and that have similar pharmacologic and toxic effects on the fetus. The stage of development of the fetal detoxifying and catabolic mechanisms crucially affects the drug's duration of action. Since our present knowledge of the effects of different drugs on the fetus is severely limited, we must refrain wherever possible from administering any drug during pregnancy, especially if there is some vague reason to suspect gravidity.

The therapeutic or prophylactic benefits of a medication must always be carefully weighed against possible damage to the fetus. Pharmacokinetics and pharmacodynamics are directly related to diaplacental passage, since the placenta seems permeable to most drugs. One might assume that only the drug's concentration in the maternal blood and how long that concentration will be maintained will affect the concentrations of the drug reached in the fetal circulation. But one should always remember that (a) a concentration that produces a specific effect in the mother will not always be the same in the fetal circulation, and (b) the sensitivity of the embryo or fetus can vary throughout pregnancy. Indeed, concentrations and duration of retention in the fetal blood vary over a broad range.

Drugs can be divided into three groups: 1. Drugs that do not pass through the placenta, and thus can do no direct damage.

2. Drugs with diaplacental passage in the sense of a shallow compartment with which damage is possible.

3. Drugs with both diaplacental passage and fetal accumulation, with which damage is

slightly possible.

A general knowledge of the distribution ratio, i.e., the relationship between serum and tissue concentration at any given time (time-dependent), is needed as a parameter to determine the passage of a substance from the mother to the embryo or fetus. Most often, however, a time shift in the concentration curves of both is observed; thus, knowledge of the distribution ratio but rarely permits firm conclusions about the actual and effectual intrauterine

concentrations of the drug.

The pharmacologic consequences of drugs administered during the first and second halves of pregnancy can be recognized in changes or developmental disturbances in the fetus or newborn. At the present time they at least allow us to draw certain conclusions about exogenous noxa, e.g., drugs. We now have a reasonably certain understanding of many specific effects on certain organ rudiments, as well as the pronounced phase specificity of drugs in the sequential steps of organ genesis. However, this information is valid only for morphologic research; about the functional stages, i.e., the biochemical mechanisms, we know almost nothing. The indirect effects exerted on the fetus by influences on the maternal function have also been seldom investigated and little is known about them. We can justifiably assume, however, that an insufficient detoxifying function of the maternal liver or kidneys is not without effect on the fetus.