

**Grundlehren der
mathematischen Wissenschaften 248**

A Series of Comprehensive Studies in Mathematics

Michio Suzuki

Group Theory II

Michio Suzuki

Group Theory II



Springer-Verlag
New York Berlin Heidelberg Tokyo

MICHIO SUZUKI
Department of Mathematics
University of Illinois
1409 West Green Street
Urbana, IL 61801
U.S.A.

AMS Classification: 20XX

Library of Congress Cataloging in Publication Data
(Revised for volume 2)

Suzuki, Michio.

Group theory.

(Grundlehren der mathematischen Wissenschaften;
247-248)

Translation of; Gunron.

Bibliography: v. 1, p. [419]-427; v. 2, p.

Includes indexes.

1. Groups, Theory of. I. Title. II. Series.

QA171.S79 512'.22 81-9159

With 1 Illustration.

© 1986 by Springer-Verlag New York Inc.

All rights reserved. No part of this book may be translated or reproduced in any form without written permission from Springer-Verlag (75 Fifth Avenue, New York, New York 10010, U.S.A.).

Group Theory is a translation of the Japanese edition: *Gunron* by Michio Suzuki.
© 1977, 1978 by Michio Suzuki. Originally published in Japanese by Iwanami Shoten, Publishers, Tokyo in 1977 and 1978. Translated by Michio Suzuki.

Typeset by J. W. Arrowsmith Ltd., Bristol, England.

Printed and bound by R. R. Donnelley & Sons, Harrisonburg, Virginia.

Printed in the United States of America.

9 8 7 6 5 4 3 2 1

ISBN 0-387-10916-1 Springer-Verlag New York Berlin Heidelberg Tokyo
ISBN 3-540-10916-1 Springer-Verlag Berlin Heidelberg New York Tokyo

Grundlehren der mathematischen Wissenschaften 248

A Series of Comprehensive Studies in Mathematics

Editors

M. Artin S. S. Chern J. M. Fröhlich E. Heinz
H. Hironaka F. Hirzebruch L. Hörmander S. Mac Lane
W. Magnus C. C. Moore J. K. Moser M. Nagata
W. Schmidt D. S. Scott Ya. G. Sinai J. Tits
B. L. van der Waerden M. Waldschmidt S. Watanabe

Managing Editors

M. Berger E. Eckmann S. R. S. Varadhan

Grundlehren der mathematischen Wissenschaften

A Series of Comprehensive Studies in Mathematics

A Selection

200. Dold: Lectures on Algebraic Topology
201. Beck: Continuous Flows in the Plane
202. Schmetterer: Introduction to Mathematical Statistics
203. Schoeneberg: Elliptic Modular Functions
204. Popov: Hyperstability of Control Systems
205. Nikol'skii: Approximation of Functions of Several Variables and Imbedding Theorems
206. André: Homologie des Algèbres Commutatives
207. Donoghue: Monotone Matrix Functions and Analytic Continuation
208. Lacey: The Isometric Theory of Classical Banach Spaces
209. Ringel: Map Color Theorem
210. Gihman/Skorohod: The Theory of Stochastic Processes I
211. Comfort/Negreponis: The Theory of Ultrafilters
212. Switzer: Algebraic Topology—Homotopy and Homology
213. Shafarevich: Basic Algebraic Geometry
214. van der Waerden: Group Theory and Quantum Mechanics
215. Schaefer: Banach Lattices and Positive Operators
216. Pólya/Szegő: Problems and Theorems in Analysis II
217. Stenström: Rings of Quotients
218. Gihman/Skorohod: The Theory of Stochastic Process II
219. Duvant/Lions: Inequalities in Mechanics and Physics
220. Kirillov: Elements of the Theory of Representations
221. Mumford: Algebraic Geometry I: Complex Projective Varieties
222. Lang: Introduction to Modular Forms
223. Bergh/Löfström: Interpolation Spaces. An Introduction
224. Gilbarg/Trudinger: Elliptic Partial Differential Equations of Second Order
225. Schütte: Proof Theory
226. Karoubi: K-Theory, An Introduction
227. Grauert/Remmert: Theorie der Steinschen Räume
228. Segal/Kunze: Integrals and Operators
229. Hasse: Number Theory
230. Klingenberg: Lectures on Closed Geodesics
231. Lang: Elliptic Curves: Diophantine Analysis
232. Gihman/Skorohod: The Theory of Stochastic Processes III
233. Stroock/Varadhan: Multi-dimensional Diffusion Processes
234. Aigner: Combinatorial Theory
235. Dynkin/Yushkevich: Markov Control Processes and Their Applications
236. Grauert/Remmert: Theory of Stein Spaces
237. Köthe: Topological Vector-Spaces II
238. Graham/McGehee: Essays in Commutative Harmonic Analysis
239. Elliott: Probabilistic Number Theory I
240. Elliott: Probabilistic Number Theory II
241. Rudin: Function Theory in the Unit Ball of C^n
242. Huppert/Blackburn: Finite Groups I
243. Huppert/Blackburn: Finite Groups II
244. Kubert/Lang: Modular Units

Preface to the English Edition

This is a translation from the Japanese of the second volume (chapters four through six) of my book "Gunron" (Iwanami Shoten, 1978). After discussing the concept of commutators in the fourth chapter, we turn to a discussion of the methods and theorems pertaining to finite groups. The last chapter is intended as an introduction to the recent progress in the theory of simple groups. For the translation, I have kept the main body of the text unchanged, however I have added a few comments in the last chapter in order to inform the readers of the most recent progress.

I would like to express my appreciation to Kazuko Suzuki for her devoted help in translating this book. Finally, it gives me great pleasure to acknowledge my indebtedness to my wife, Naoko, for her constant support and understanding, and for converting the long and often illegible manuscript into a beautifully typed one. To her, I express my sincere thanks and appreciation.

July, 1985

Michio Suzuki

Contents (Part II)

List of Notation	ix
Chapter 4	
Commutators	1
§1. Commutator Subgroups	1
§2. Nilpotent Groups	13
§3. Commutator Calculations	37
§4. Finite p -Groups	54
§5. Finite Solvable Groups	100
Chapter 5	
Finite Groups I	120
§1. Elements of Order Two	120
§2. Transfer and Fusion of Elements	138
§3. Generalizations of Sylow's Theorems	166
§4. ZJ -Theorem	189
§5. Supplements	226
Chapter 6	
Finite Groups II	258
§1. Representation Theory	258
§2. Applications of Representation Theory	275
§3. Groups of Odd Order	356
§4. Strongly Embedded Subgroups	391
§5. The Classification Problem of Simple Groups	418
§6. The Components of a Group	446
§7. Signalizer Functors	470
§8. Small Simple Groups	494
§9. Two Types of Simple Groups	519
§10. Graphs and Simple Groups	543
§11. Fusions in S_2 -subgroups	582
§12. Recent Developments in the Theory of Simple Groups	602
Bibliography	611
Index	617

Contents (Part I)

Chapter 1	
Basic Concepts	1
Chapter 2	
Fundamental Theorems	86
Chapter 3	
Some Special Classes of Groups	280
Bibliography	419
Index	428

List of Notation

(The notation which appeared in the List of Notation in the first volume will be omitted.)

$ G $	I. p. 6	$\exp G$	61
$ G _p$	I. p. 101	$m(G)$	64
$[x, y]$	I. p. 119	$E(p^3)$	67
$D(G)$	I. p. 119	\mathcal{A}_p	80
G'	I. p. 119	$J(P)$	80
$G^{(i)}$	I. p. 119	$ZJ(P)$	81
$G^{(\infty)}$	I. p. 119	$O_{\pi', \pi}(G)$	109
$H \triangleleft \triangleleft G$	I. p. 123	$\mathcal{H}(H; \pi)$	111
$H \times K$	I. p. 127	$\mathcal{H}^*(H; \pi)$	112
$H \text{ wr } K$	I. p. 270	$I(G)$	120
$[H, K]$	2	$i(G)$	120
$[H; x]$	4	$Sr(G)$	120
$[x_1, x_2, \dots, x_n]$	4	$C_G^*(x)$	121
$[H_1, H_2, \dots, H_n]$	4	$\text{Foc}_G(P)$	141
$O^\pi(G)$	8	$G'(p)$	142
$C_i(G)$	13	E_π	166
$Z_i(G)$	14	C_π	166
$\Phi(G)$	27	D_π	166
$F(G)$	28	E_π^n	166
$O_\pi(G)$	28	$\text{proj}_i(H)$	174
$O_P(G)$	28	$\mathcal{M}_\pi^*(G)$	177
$C_\infty(G)$	31	$\mathcal{M}_\pi(G)$	177
$[H, K; n]$	33	$\mathcal{L}_p(G)$	189
$Z_\infty(G)$	36	$\mathcal{L}_p^*(G)$	189
$x \equiv y \pmod{N}$	44	$\mathcal{L}(G)$	189
$\Omega_n(G)$	45	$Qd(p)$	190
$U_n(G)$	45	$K^\infty(P)$	227
$M(p^n)$	54	$K_\infty(P)$	227
D_g	54	\mathcal{S}	248
Q_g	54	$(\chi, \theta)_G$	267
S_g	54	θ^G	270

K^G 282
 $gc(H; K)$ 282
 $Z^*(G)$ 315
 $SCN(P)$ 357
 $SCN(p)$ 358
 $SCN_k(P)$ 358
 $Y \text{ cc } X$ 372
 $M * N$ 373
 $\mathcal{E}_k(p; H)$ 375
 \mathcal{M} 375
 $\mathcal{M}(X)$ 375
 \mathcal{U} 375
 $\text{Syl}_p(G)$ 377

$O(G)$ 392
 $E(G)$ 450
 $F^*(G)$ 452
 $L(G)$ 454
 $B(G)$ 457
 $O_*(G)$ 461
 $C_G^Z(x)$ 466
 $\mathcal{H}_\theta(H; \pi)$ 472
 $\mathcal{H}_\theta^*(H; \pi)$ 472
 $m_p(X)$ 488
 $\mathcal{C}(G)$ 519
 $\Gamma_{P,2}(G)$ 520
 $\mathcal{H}'(H; \pi)$ 591

Commutators

§1. Commutator Subgroups

The commutator $[x, y] = x^{-1}y^{-1}xy$ of two elements x and y of a group G has been used, without formal development, several times in this book since its initial appearance in Chapter 1, §3. The readers are referred to Definition 3.11 of Chapter 2, p. 110, for the definition, and to proposition (3.12) of Chapter 2, as well as the remarks preceding it, for some of the important properties of the commutator. In this section, we will define the commutator subgroup $[H, K]$ of two arbitrary subgroups H and K of a group G . We will study the properties of the commutator subgroups and also give applications.

The following proposition gives some of the useful relationships among commutators. (Recall the notation $v^y = y^{-1}vy$ for any elements y and v .)

(1.1) *Let x , y , and u be three elements of a group G . Then, the following identities hold:*

- (i) $[x, y]^{-1} = [y, x],$
- (ii) $[xy, u] = [x, u]^y [y, u],$
- (iii) $[u, xy] = [u, y][u, x]^y.$

Proof. These identities are easily verified by using the definition of the commutator. We will verify the third one as an example. We have

$$\begin{aligned}
 [u, xy] &= u^{-1}(xy)^{-1}u(xy) \\
 &= u^{-1}y^{-1}x^{-1}uxy \\
 &= u^{-1}y^{-1}(uyy^{-1}u^{-1})x^{-1}uxy \\
 &= [u, y]y^{-1}[u, x]y.
 \end{aligned}$$

This proves identity (iii). Note that the commutator involving x comes after the commutator which involves y in the right side of identity (iii). \square

The following definition is basic to the subsequent development of the theory.

Definition 1.2. Let H and K be two subgroups of a group G . The **commutator subgroup** of H and K is defined as the subgroup generated by all the commutators $[h, k]$ where $h \in H$ and $k \in K$. We denote it by $[H, K]$:

$$[H, K] = \langle [h, k] \mid h \in H, k \in K \rangle.$$

If $H = K = G$, the subgroup $[H, K]$ defined here coincides with the commutator subgroup $D(G)$ of G defined in Definition 3.11 of Chapter 2, p. 119. In this chapter, we will call $D(G) = G'$ the **derived group** in order to lessen the probability of confusion. We remark again here that the subgroup $[H, K]$ is not just the totality of the commutators, but the subgroup generated by these commutators.

The following omnibus lemma contains some basic properties of commutator subgroups.

(1.3) Let H and K be two subgroups of a group G . Then, the following propositions hold.

- (i) We have $[H, K] = [K, H]$.
- (ii) For any homomorphism σ of G , we have

$$[H, K]^\sigma = [H^\sigma, K^\sigma].$$

- (iii) $[H, K] \subset H \Leftrightarrow K \subset N_G(H)$.
- (iv) If H_1 and K_1 are subgroups such that $H \subset H_1$ and $K_1 \subset K$, then we have

$$[H_1, K_1] \subset [H, K].$$

- (v) $[H, K] = \{1\}$ if and only if H commutes elementwise with K .

Proof. Clearly, proposition (i) follows from (1.1)(i). Since

$$[h, k]^\sigma = [h^\sigma, k^\sigma]$$

for any homomorphism σ of G (cf. (3.12) of Chapter 2), the second proposition holds. If $[H, K] \subset H$, then for any $h \in H$ and $k \in K$, we have

$$[h, k] = h^{-1}k^{-1}hk \in H.$$

Thus, we get $k^{-1}hk \in H$ and $k^{-1}Hk \subset H$. If we consider k^{-1} in place of k , we obtain $kHk^{-1} \subset H$. So, we have $k^{-1}Hk = H$ and $k \in N_G(H)$. Conversely, if $k \in N_G(H)$, then for any element h of H , we get $[h, k] \in H$. It follows

from definition that $[H, K] \subset H$ holds. This proves the equivalence in (iii). The remaining part of (1.3) follows easily from the definitions. \square

Theorem 1.4. *If two subgroups H and K are normal subgroups of a group G , then so is the commutator subgroup $[H, K]$. Furthermore, we have*

$$[H, K] \subset H \cap K.$$

A similar theorem holds for characteristic, as well as for fully invariant, subgroups.

Proof. Let Ω be an operator domain of G . We prove that if both H and K are Ω -subgroups of G , then the commutator subgroup $[H, K]$ is also an Ω -subgroup. In fact, if $\sigma \in \Omega$, then we have

$$[H, K]^\sigma = [H^\sigma, K^\sigma] \subset [H, K]$$

by (1.3)(ii) and (iv). Thus, $[H, K]$ is Ω -invariant. For $\Omega = \text{Inn } G$, we get the first statement of Theorem 1.4. The last assertions will follow by taking $\Omega = \text{Aut } G$ and $\Omega = \text{End } G$. The containment

$$[H, K] \subset H \cap K$$

is proved easily from (1.3)(iii). \square

Theorem 1.5. *Let N be a normal subgroup of a group G , and set $\bar{G} = G/N$. Then, for any two subgroups H and K of G , we have*

$$[\bar{H}, \bar{K}] = [H, K]N/N.$$

Proof. Let σ be the canonical homomorphism from G onto \bar{G} . Then, we have $\bar{H} = H^\sigma = HN/N$. So, the assertion follows from (1.3)(ii). \square

The next theorem is one of the most important properties of commutator subgroups.

Theorem 1.6. *Let H and K be two subgroups of a group G , and let L be the commutator subgroup $[H, K]$.*

- (i) *The subgroup H , as well as K , normalizes L ; that is, we have*

$$I \triangleleft \langle H, K \rangle.$$

- (ii) *The subgroup LK is the subgroup generated by all the subgroups conjugate to K in $\langle H, K \rangle$. The subgroup LK is the smallest normal subgroup of $\langle H, K \rangle$ which contains K .*

Proof. (i) We will prove that $H \subset N_G(L)$. It suffices to show that for any element u of H and for any generator $[h, k]$ ($h \in H, k \in K$) of L , we have $[h, k]^u \in L$. By (1.1)(ii), we have

$$[h, k]^u = [hu, k][u, k]^{-1}.$$

Since h and u are elements of H , we have $hu \in H$, and the right side of the above equality belongs to L . Thus, we have $H \subset N_G(L)$. By (1.3)(i), we have $L = [H, K] = [K, H]$. So, similarly, the subgroup K normalizes L . This proves the first part.

(ii) For any element h of H and any element k of K , we have

$$h^{-1}kh = k[k, h] \in KL.$$

Hence, we get $(KL)^h = K^h L^h \subset KL$. This implies that H normalizes KL and that $KL \triangleleft \langle H, K \rangle$.

Suppose that $K \subset M \triangleleft \langle H, K \rangle$ for some subgroup M of G . Then, the commutator $[h, k]$ is contained in M , and we have $KL \subset M$. Thus, KL is the smallest normal subgroup of $\langle H, K \rangle$ which contains K . The first part of (ii) follows easily because the subgroup generated by the conjugate subgroups is normal. \square

For any element x of G , we define the **commutator subgroup** $[H, x]$ as the subgroup of G which is generated by $[h, x]$ with $h \in H$. Then, a proposition similar to (1.3) holds. Furthermore, the proof of Theorem 1.6(i) shows that we have

$$(1.6)' \quad H \subset N_G([H, x]).$$

Definition 1.7. Let x_1, x_2, \dots, x_n ($n \geq 3$) be elements of a group G . We define a **higher commutator** of n elements inductively by the formula

$$[x_1, x_2, \dots, x_n] = [[x_1, \dots, x_{n-1}], x_n].$$

We will often call it simply the commutator of the n elements x_1, x_2, \dots, x_n . Similarly, a **(higher) commutator subgroup** of n subgroups H_1, H_2, \dots, H_n is defined by

$$[H_1, H_2, \dots, H_n] = [[H_1, \dots, H_{n-1}], H_n].$$

A higher commutator subgroup $[H_1, H_2, \dots, H_n]$ contains all the higher commutators $[h_1, h_2, \dots, h_n]$ of elements h_1, \dots, h_n , where $h_i \in H_i$ for each i . We remark here that if $n \geq 3$, the subgroup generated by these higher commutators of elements is not necessarily equal to the higher commutator subgroup $[H_1, H_2, \dots, H_n]$ (cf. Exercise 3). We defined the higher commu-

tator by taking the commutator of two elements successively starting from the left end. Thus, in general, we have

$$[x, y, z] \neq [x, [y, z]].$$

We may rewrite identities (1.1)(ii) and (iii) using higher commutators:

$$(1.1)(ii)' \quad [xy, u] = [x, u][x, u, y][y, u],$$

$$(1.1)(iii)' \quad [u, xy] = [u, y][u, x][u, x, y].$$

Among the higher commutators of three elements, the following remarkable identity holds.

(1.8) For three elements x, y , and z of a group, we have

$$[x, y^{-1}, z]^y [y, z^{-1}, x]^z [z, x^{-1}, y]^x = 1.$$

Proof. Express the higher commutators involved as words of the elements x, y , and z . Then, all the terms will be cancelled. In fact, from

$$[x, y^{-1}] = x^{-1} y x y^{-1},$$

we get

$$\begin{aligned} [x, y^{-1}, z] &= (x^{-1} y x y^{-1})^{-1} z^{-1} (x^{-1} y x y^{-1}) z \\ &= y x^{-1} y^{-1} x z^{-1} x^{-1} y x y^{-1} z. \end{aligned}$$

Set $u = x z x^{-1} y x$, $v = y x y^{-1} z y$, and $w = z y z^{-1} x z$. Then, we have

$$[x, y^{-1}, z]^y = u^{-1} v.$$

Similarly, we have $[y, z^{-1}, x]^z = v^{-1} w$ and $[z, x^{-1}, y]^x = w^{-1} u$. It is now clear that (1.8) holds. \square

Using formula (1.8) we can prove the following **Three Subgroup Lemma** of P. Hall which, because of its many consequences, may be called one of the most basic results in group theory.

Theorem 1.9. Let H, K , and L be three subgroups of a group G . If

$$[H, K, L] = [K, L, H] = \{1\},$$

then we have $[L, H, K] = \{1\}$.

Proof. Let x, y , and z be arbitrary elements such that $x \in H$, $y \in K$, and $z \in L$. Then, by assumption, we have

$$[x, y^{-1}, z] = [y, z^{-1}, x] = 1.$$

Hence, by (1.8), we get

$$[z, x^{-1}, y]^x = 1 = [z, x^{-1}, y].$$

By definition, the commutator subgroup $[L, H]$ is generated by the commutators of the form $[z, x^{-1}]$. The above identity shows that any element y of K commutes with any element of the form $[z, x^{-1}]$ where $z \in L$ and $x \in H$. Thus, K centralizes $[L, H]$, and we have $[L, H, K] = \{1\}$ by (1.3)(v). \square

Corollary. Let N be a normal subgroup of a group G , and let H, K , and L be arbitrary subgroups of G . If N contains two of the following three higher commutator subgroups

$$[H, K, L], \quad [K, L, H], \quad \text{and} \quad [L, H, K],$$

then N contains all three commutator subgroups.

Proof. Let $\bar{G} = G/N$. By assumption, two of the following three higher commutator subgroups

$$[\bar{H}, \bar{K}, \bar{L}], \quad [\bar{K}, \bar{L}, \bar{H}], \quad \text{and} \quad [\bar{L}, \bar{H}, \bar{K}]$$

are $\{1\}$. By Theorem 1.9, all these higher commutator subgroups are $\{1\}$. This proves the Corollary. \square

The concept of commutator subgroups can be generalized to groups with an operator group.

Definition 1.10. Let A be an operator group which acts on a group G (cf. Definition 8.1 of Chapter 1, p. 66). We define

$$[G, A] = \langle g^{-1}g^a \mid g \in G, a \in A \rangle$$

and call it the **commutator subgroup** of G and A .

If a group A acts on a group G , we can form their semidirect product S by the method discussed in Chapter 1, §8. We may consider both G and A as subgroups of S . Then, the given action of A on G is realized by conjugation in S . Thus, the commutator subgroup $[G, A]$ of Definition 1.10

coincides with the commutator subgroup of the two subgroups G and A in the semidirect product S . This explains the choice of the notation and the terminology in Definition 1.10, and assures us that there will be no distinction between the two possible meanings of $[G, A]$ when A may be considered both as a subgroup of S and, at the same time, as an operator group of G . Furthermore, the theorems which have been proved so far about commutator subgroups also hold for commutator subgroups of operator groups. In particular, we have the following result.

(1.11) *Let A be an operator group which acts on a group G . Then, the commutator subgroup $[G, A]$ is an A -invariant normal subgroup of G .*

Proof. This is a special case of Theorem 1.6(i). \square

Although the concept of commutator subgroups has not been defined formally until this section, several results of Chapter 2 concern commutator subgroups. Thus, the subgroup denoted as A_0 in the proof of (5.17) of Chapter 2, p. 156, is the commutator subgroup $[A, Q]$, and (5.17) of Chapter 2 gives the direct sum decomposition

$$A = C_A(Q) + [A, Q]$$

for the abelian group A with a regular operator group Q . The Corollary of Theorem 8.13 of Chapter 2, p. 239, shows that if a π' -group Q acts on a π -group G , then we have

$$G = C_G(Q)[G, Q].$$

The proof given there for this corollary is a special case of the proof of (1.11). The commutator subgroup appeared also in (9.2) of Chapter 2, p. 246.

The following lemma gives a useful characterization of the commutator subgroup of a group with operators.

(1.12) *Let A be an operator group which acts on a group G . Let S be the semidirect product of G and A with respect to the given action of A on G , and consider G and A as subgroups of S . Then, we have*

$$[G, A] = G \cap N,$$

where N is the smallest normal subgroup such that $A \subset N \triangleleft S$.

Proof. Let N be as defined above. Then, by Theorem 1.6(ii), we have

$$N = [G, A]A.$$