

实分析与复分析

(英文版·第3版)

REAL AND COMPLEX ANALYSIS

Walter Rudin

Third Edition



McGRAW-HILL INTERNATIONAL EDITIONS
Mathematics Series

(美) Walter Rudin 著



机械工业出版社
China Machine Press



Education

经典原版书库

实分析与复分析

(英文版·第3版)

Real and Complex Analysis

(Third Edition)

(美) Walter Rudin 著



机械工业出版社
China Machine Press

Walter Rudin: Real and Complex Analysis, Third Edition (ISBN:0-07-054234-1).

Copyright © 1987, 1974, 1966 by the McGraw-Hill Companies, Inc.

Original language published by the McGraw-Hill Companies, Inc. All rights reserved. No part of this publication may be reproduced or distributed in any means, or stored in a database or retrieval system, without the prior written permission of the publisher.

Authorized English language reprint edition jointly published by McGraw-Hill Education (Asia) Co. and China Machine Press. This edition is authorized for sale in the People's Republic of China only, excluding Hong Kong, Macao SARs and Taiwan. Unauthorized export of this edition is a violation of the Copyright Act. Violation of this Law is subject to Civil and Criminal Penalties.

本书英文影印版由机械工业出版社和美国麦格劳-希尔教育出版(亚洲)公司合作出版。此版本仅限在中华人民共和国境内(不包括香港、澳门特别行政区及台湾)销售。未经许可之出口,视为违反著作权法,将受法律之制裁。

未经出版者预先书面许可,不得以任何方式复制或抄袭本书的任何部分。

本书封面贴有McGraw-Hill公司防伪标签,无标签者不得销售。

版权所有,侵权必究。

本书版权登记号:图字:01-2003-7218

图书在版编目(CIP)数据

实分析与复分析(英文版·第3版)/(美)鲁丁(Rudin, W.)著. -北京:机械工业出版社, 2004.1

(经典原版书库)

书名原文: Real and Complex Analysis, Third Edition

ISBN 7-111-13305-6

I. 实… II. 鲁… III. ①实分析-英文 ②复分析-英文 IV. O174

中国版本图书馆CIP数据核字(2003)第100322号

机械工业出版社(北京市西城区百万庄大街22号 邮政编码 100037)

责任编辑:迟振春

北京瑞德印刷有限公司印刷·新华书店北京发行所发行

2004年1月第1版第1次印刷

787mm×1092mm 1/16·26.75印张

印数:0 001-3 000册

定价:39.00元

凡购本书,如有倒页、脱页、缺页,由本社发行部调换

本社购书热线电话:(010) 68326294

ABOUT THE AUTHOR

Walter Rudin is the author of three textbooks, *Principles of Mathematical Analysis*, *Real and Complex Analysis*, and *Functional Analysis*, whose widespread use is illustrated by the fact that they have been translated into a total of 13 languages. He wrote the first of these while he was a C.L.E. Moore Instructor at M.I.T., just two years after receiving his Ph.D. at Duke University in 1949. Later he taught at the University of Rochester, and is now a Vilas Research Professor at the University of Wisconsin–Madison, where he has been since 1959.

He has spent leaves at Yale University, at the University of California in La Jolla, and at the University of Hawaii.

His research has dealt mainly with harmonic analysis and with complex variables. He has written three research monographs on these topics, *Fourier Analysis on Groups*, *Function Theory in Polydiscs*, and *Function Theory in the Unit Ball of C^n* .

PREFACE

This book contains a first-year graduate course in which the basic techniques and theorems of analysis are presented in such a way that the intimate connections between its various branches are strongly emphasized. The traditionally separate subjects of “real analysis” and “complex analysis” are thus united; some of the basic ideas from functional analysis are also included.

Here are some examples of the way in which these connections are demonstrated and exploited. The Riesz representation theorem and the Hahn-Banach theorem allow one to “guess” the Poisson integral formula. They team up in the proof of Runge’s theorem. They combine with Blaschke’s theorem on the zeros of bounded holomorphic functions to give a proof of the Müntz-Szasz theorem, which concerns approximation on an interval. The fact that L^2 is a Hilbert space is used in the proof of the Radon-Nikodym theorem, which leads to the theorem about differentiation of indefinite integrals, which in turn yields the existence of radial limits of bounded harmonic functions. The theorems of Plancherel and Cauchy combined give a theorem of Paley and Wiener which, in turn, is used in the Denjoy-Carleman theorem about infinitely differentiable functions on the real line. The maximum modulus theorem gives information about linear transformations on L^p -spaces.

Since most of the results presented here are quite classical (the novelty lies in the arrangement, and some of the proofs are new), I have not attempted to document the source of every item. References are gathered at the end, in Notes and Comments. They are not always to the original sources, but more often to more recent works where further references can be found. In no case does the absence of a reference imply any claim to originality on my part.

The prerequisite for this book is a good course in advanced calculus (settheoretic manipulations, metric spaces, uniform continuity, and uniform convergence). The first seven chapters of my earlier book “Principles of Mathematical Analysis” furnish sufficient preparation.

Experience with the first edition shows that first-year graduate students can study the first 15 chapters in two semesters, plus some topics from 1 or 2 of the remaining 5. These latter are quite independent of each other. The first 15 should be taken up in the order in which they are presented, except for Chapter 9, which can be postponed.

The most important difference between this third edition and the previous ones is the entirely new chapter on differentiation. The basic facts about differentiation are now derived from the existence of Lebesgue points, which in turn is an easy consequence of the so-called "weak type" inequality that is satisfied by the maximal functions of measures on euclidean spaces. This approach yields strong theorems with minimal effort. Even more important is that it familiarizes students with maximal functions, since these have become increasingly useful in several areas of analysis.

One of these is the study of the boundary behavior of Poisson integrals. A related one concerns H^p -spaces. Accordingly, large parts of Chapters 11 and 17 were rewritten and, I hope, simplified in the process.

I have also made several smaller changes in order to improve certain details: For example, parts of Chapter 4 have been simplified; the notions of equicontinuity and weak convergence are presented in more detail; the boundary behavior of conformal maps is studied by means of Lindelöf's theorem about asymptotic values of bounded holomorphic functions in a disc.

Over the last 20 years, numerous students and colleagues have offered comments and criticisms concerning the content of this book. I sincerely appreciated all of these, and have tried to follow some of them. As regards the present edition, my thanks go to Richard Rochberg for some useful last-minute suggestions, and I especially thank Robert Burckel for the meticulous care with which he examined the entire manuscript.

Walter Rudin

CONTENTS

Preface	ix
Prologue: The Exponential Function	1
Chapter 1 Abstract Integration	5
Set-theoretic notations and terminology	6
The concept of measurability	8
Simple functions	15
Elementary properties of measures	16
Arithmetic in $[0, \infty]$	18
Integration of positive functions	19
Integration of complex functions	24
The role played by sets of measure zero	27
Exercises	31
Chapter 2 Positive Borel Measures	33
Vector spaces	33
Topological preliminaries	35
The Riesz representation theorem	40
Regularity properties of Borel measures	47
Lebesgue measure	49
Continuity properties of measurable functions	55
Exercises	57
Chapter 3 L^p-Spaces	61
Convex functions and inequalities	61
The L^p -spaces	65
Approximation by continuous functions	69
Exercises	71

Chapter 4	Elementary Hilbert Space Theory	76
	Inner products and linear functionals	76
	Orthonormal sets	82
	Trigonometric series	88
	Exercises	92
Chapter 5	Examples of Banach Space Techniques	95
	Banach spaces	95
	Consequences of Baire's theorem	97
	Fourier series of continuous functions	100
	Fourier coefficients of L^1 -functions	103
	The Hahn-Banach theorem	104
	An abstract approach to the Poisson integral	108
	Exercises	112
Chapter 6	Complex Measures	116
	Total variation	116
	Absolute continuity	120
	Consequences of the Radon-Nikodym theorem	124
	Bounded linear functionals on L^p	126
	The Riesz representation theorem	129
	Exercises	132
Chapter 7	Differentiation	135
	Derivatives of measures	135
	The fundamental theorem of Calculus	144
	Differentiable transformations	150
	Exercises	156
Chapter 8	Integration on Product Spaces	160
	Measurability on cartesian products	160
	Product measures	163
	The Fubini theorem	164
	Completion of product measures	167
	Convolutions	170
	Distribution functions	172
	Exercises	174
Chapter 9	Fourier Transforms	178
	Formal properties	178
	The inversion theorem	180
	The Plancherel theorem	185
	The Banach algebra L^1	190
	Exercises	193

Chapter 10	Elementary Properties of Holomorphic Functions	196
	Complex differentiation	196
	Integration over paths	200
	The local Cauchy theorem	204
	The power series representation	208
	The open mapping theorem	214
	The global Cauchy theorem	217
	The calculus of residues	224
	Exercises	227
Chapter 11	Harmonic Functions	231
	The Cauchy-Riemann equations	231
	The Poisson integral	233
	The mean value property	237
	Boundary behavior of Poisson integrals	239
	Representation theorems	245
	Exercises	249
Chapter 12	The Maximum Modulus Principle	253
	Introduction	253
	The Schwarz lemma	254
	The Phragmen-Lindelöf method	256
	An interpolation theorem	260
	A converse of the maximum modulus theorem	262
	Exercises	264
Chapter 13	Approximation by Rational Functions	266
	Preparation	266
	Runge's theorem	270
	The Mittag-Leffler theorem	273
	Simply connected regions	274
	Exercises	276
Chapter 14	Conformal Mapping	278
	Preservation of angles	278
	Linear fractional transformations	279
	Normal families	281
	The Riemann mapping theorem	282
	The class $\mathcal{S}$	285
	Continuity at the boundary	289
	Conformal mapping of an annulus	291
	Exercises	293

Chapter 15	Zeros of Holomorphic Functions	298
	Infinite products	298
	The Weierstrass factorization theorem	301
	An interpolation problem	304
	Jensen's formula	307
	Blaschke products	310
	The Müntz-Szasz theorem	312
	Exercises	315
Chapter 16	Analytic Continuation	319
	Regular points and singular points	319
	Continuation along curves	323
	The monodromy theorem	326
	Construction of a modular function	328
	The Picard theorem	331
	Exercises	332
Chapter 17	H^p-Spaces	335
	Subharmonic functions	335
	The spaces H^p and N	337
	The theorem of F. and M. Riesz	341
	Factorization theorems	342
	The shift operator	346
	Conjugate functions	350
	Exercises	352
Chapter 18	Elementary Theory of Banach Algebras	356
	Introduction	356
	The invertible elements	357
	Ideals and homomorphisms	362
	Applications	365
	Exercises	369
Chapter 19	Holomorphic Fourier Transforms	371
	Introduction	371
	Two theorems of Paley and Wiener	372
	Quasi-analytic classes	377
	The Denjoy-Carleman theorem	380
	Exercises	383
Chapter 20	Uniform Approximation by Polynomials	386
	Introduction	386
	Some lemmas	387
	Mergelyan's theorem	390
	Exercises	394

Appendix: Hausdorff's Maximality Theorem	395
Notes and Comments	397
Bibliography	405
List of Special Symbols	407
Index	409

PROLOGUE

THE EXPONENTIAL FUNCTION

This is the most important function in mathematics. It is defined, for every complex number z , by the formula

$$\exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}. \quad (1)$$

The series (1) converges absolutely for every z and converges uniformly on every bounded subset of the complex plane. Thus $\exp$ is a continuous function. The absolute convergence of (1) shows that the computation

$$\sum_{k=0}^{\infty} \frac{a^k}{k!} \sum_{m=0}^{\infty} \frac{b^m}{m!} = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=0}^n \frac{n!}{k!(n-k)!} a^k b^{n-k} = \sum_{n=0}^{\infty} \frac{(a+b)^n}{n!}$$

is correct. It gives the important addition formula

$$\exp(a) \exp(b) = \exp(a+b), \quad (2)$$

valid for all complex numbers a and b .

We define the number e to be $\exp(1)$, and shall usually replace $\exp(z)$ by the customary shorter expression e^z . Note that $e^0 = \exp(0) = 1$, by (1).

Theorem

- (a) For every complex z we have $e^z \neq 0$.
- (b) $\exp$ is its own derivative: $\exp'(z) = \exp(z)$.

(c) The restriction of $\exp$ to the real axis is a monotonically increasing positive function, and

$$e^x \rightarrow \infty \text{ as } x \rightarrow \infty, \quad e^x \rightarrow 0 \text{ as } x \rightarrow -\infty.$$

(d) There exists a positive number π such that $e^{\pi i/2} = i$ and such that $e^z = 1$ if and only if $z/(2\pi i)$ is an integer.

(e) $\exp$ is a periodic function, with period $2\pi i$.

(f) The mapping $t \rightarrow e^{it}$ maps the real axis onto the unit circle.

(g) If w is a complex number and $w \neq 0$, then $w = e^z$ for some z .

PROOF By (2), $e^z \cdot e^{-z} = e^{z-z} = e^0 = 1$. This implies (a). Next,

$$\exp'(z) = \lim_{h \rightarrow 0} \frac{\exp(z+h) - \exp(z)}{h} = \exp(z) \lim_{h \rightarrow 0} \frac{\exp(h) - 1}{h} = \exp(z).$$

The first of the above equalities is a matter of definition, the second follows from (2), and the third from (1), and (b) is proved.

That $\exp$ is monotonically increasing on the positive real axis, and that $e^x \rightarrow \infty$ as $x \rightarrow \infty$, is clear from (1). The other assertions of (c) are consequences of $e^x \cdot e^{-x} = 1$.

For any real number t , (1) shows that e^{-it} is the complex conjugate of e^{it} . Thus

$$|e^{it}|^2 = e^{it} \cdot \overline{e^{it}} = e^{it} \cdot e^{-it} = e^{it-it} = e^0 = 1,$$

or

$$|e^{it}| = 1 \quad (t \text{ real}). \quad (3)$$

In other words, if t is real, e^{it} lies on the unit circle. We define $\cos t$, $\sin t$ to be the real and imaginary parts of e^{it} :

$$\cos t = \operatorname{Re} [e^{it}], \quad \sin t = \operatorname{Im} [e^{it}] \quad (t \text{ real}). \quad (4)$$

If we differentiate both sides of Euler's identity

$$e^{it} = \cos t + i \sin t, \quad (5)$$

which is equivalent to (4), and if we apply (b), we obtain

$$\cos' t + i \sin' t = ie^{it} = -\sin t + i \cos t,$$

so that

$$\cos' = -\sin, \quad \sin' = \cos. \quad (6)$$

The power series (1) yields the representation

$$\cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \frac{t^6}{6!} + \cdots. \quad (7)$$

Take $t = 2$. The terms of the series (7) then decrease in absolute value (except for the first one) and their signs alternate. Hence $\cos 2$ is less than the sum of the first three terms of (7), with $t = 2$; thus $\cos 2 < -\frac{1}{3}$. Since $\cos 0 = 1$ and $\cos$ is a continuous real function on the real axis, we conclude that there is a smallest positive number t_0 for which $\cos t_0 = 0$. We define

$$\pi = 2t_0. \quad (8)$$

It follows from (3) and (5) that $\sin t_0 = \pm 1$. Since

$$\sin'(t) = \cos t > 0$$

on the segment $(0, t_0)$ and since $\sin 0 = 0$, we have $\sin t_0 > 0$, hence $\sin t_0 = 1$, and therefore

$$e^{\pi i/2} = i. \quad (9)$$

It follows that $e^{\pi i} = i^2 = -1$, $e^{2\pi i} = (-1)^2 = 1$, and then $e^{2\pi i n} = 1$ for every integer n . Also, (e) follows immediately:

$$e^{z+2\pi i} = e^z e^{2\pi i} = e^z. \quad (10)$$

If $z = x + iy$, x and y real, then $e^z = e^x e^{iy}$; hence $|e^z| = e^x$. If $e^z = 1$, we therefore must have $e^x = 1$, so that $x = 0$; to prove that $y/2\pi$ must be an integer, it is enough to show that $e^{iy} \neq 1$ if $0 < y < 2\pi$, by (10).

Suppose $0 < y < 2\pi$, and

$$e^{iy/4} = u + iv \quad (u \text{ and } v \text{ real}). \quad (11)$$

Since $0 < y/4 < \pi/2$, we have $u > 0$ and $v > 0$. Also

$$e^{iy} = (u + iv)^4 = u^4 - 6u^2v^2 + v^4 + 4iuv(u^2 - v^2). \quad (12)$$

The right side of (12) is real only if $u^2 = v^2$; since $u^2 + v^2 = 1$, this happens only when $u^2 = v^2 = \frac{1}{2}$, and then (12) shows that

$$e^{iy} = -1 \neq 1.$$

This completes the proof of (d).

We already know that $t \rightarrow e^{it}$ maps the real axis into the unit circle. To prove (f), fix w so that $|w| = 1$; we shall show that $w = e^{it}$ for some real t . Write $w = u + iv$, u and v real, and suppose first that $u \geq 0$ and $v \geq 0$. Since $u \leq 1$, the definition of π shows that there exists a t , $0 \leq t \leq \pi/2$, such that $\cos t = u$; then $\sin^2 t = 1 - u^2 = v^2$, and since $\sin t \geq 0$ if $0 \leq t \leq \pi/2$, we have $\sin t = v$. Thus $w = e^{it}$.

If $u < 0$ and $v \geq 0$, the preceding conditions are satisfied by $-iw$. Hence $-iw = e^{it}$ for some real t , and $w = e^{i(t+\pi/2)}$. Finally, if $v < 0$, the preceding two cases show that $-w = e^{it}$ for some real t , hence $w = e^{i(t+\pi)}$. This completes the proof of (f).

If $w \neq 0$, put $\alpha = w/|w|$. Then $w = |w|\alpha$. By (c), there is a real x such that $|w| = e^x$. Since $|\alpha| = 1$, (f) shows that $\alpha = e^{iy}$ for some real y . Hence $w = e^{x+iy}$. This proves (g) and completes the theorem. ////

We shall encounter the integral of $(1 + x^2)^{-1}$ over the real line. To evaluate it, put $\varphi(t) = \sin t/\cos t$ in $(-\pi/2, \pi/2)$. By (6), $\varphi' = 1 + \varphi^2$. Hence φ is a monotonically increasing mapping of $(-\pi/2, \pi/2)$ onto $(-\infty, \infty)$, and we obtain

$$\int_{-\infty}^{\infty} \frac{dx}{1 + x^2} = \int_{-\pi/2}^{\pi/2} \frac{\varphi'(t) dt}{1 + \varphi^2(t)} = \int_{-\pi/2}^{\pi/2} dt = \pi.$$

ABSTRACT INTEGRATION

Toward the end of the nineteenth century it became clear to many mathematicians that the Riemann integral (about which one learns in calculus courses) should be replaced by some other type of integral, more general and more flexible, better suited for dealing with limit processes. Among the attempts made in this direction, the most notable ones were due to Jordan, Borel, W. H. Young, and Lebesgue. It was Lebesgue's construction which turned out to be the most successful.

In brief outline, here is the main idea: The Riemann integral of a function f over an interval $[a, b]$ can be approximated by sums of the form

$$\sum_{i=1}^n f(t_i)m(E_i)$$

where $E_1, \dots, E_n$ are disjoint intervals whose union is $[a, b]$, $m(E_i)$ denotes the length of E_i , and $t_i \in E_i$ for $i = 1, \dots, n$. Lebesgue discovered that a completely satisfactory theory of integration results if the sets E_i in the above sum are allowed to belong to a larger class of subsets of the line, the so-called "measurable sets," and if the class of functions under consideration is enlarged to what he called "measurable functions." The crucial set-theoretic properties involved are the following: The union and the intersection of any countable family of measurable sets are measurable; so is the complement of every measurable set; and, most important, the notion of "length" (now called "measure") can be extended to them in such a way that

$$m(E_1 \cup E_2 \cup E_3 \cup \cdots) = m(E_1) + m(E_2) + m(E_3) + \cdots$$

for every countable collection $\{E_i\}$ of pairwise disjoint measurable sets. This property of m is called *countable additivity*.

The passage from Riemann's theory of integration to that of Lebesgue is a process of completion (in a sense which will appear more precisely later). It is of the same fundamental importance in analysis as is the construction of the real number system from the rationals.

The above-mentioned measure m is of course intimately related to the geometry of the real line. In this chapter we shall present an abstract (axiomatic) version of the Lebesgue integral, relative to *any* countably additive measure on *any* set. (The precise definitions follow.) This abstract theory is not in any way more difficult than the special case of the real line; it shows that a large part of integration theory is independent of any geometry (or topology) of the underlying space; and, of course, it gives us a tool of much wider applicability. The existence of a large class of measures, among them that of Lebesgue, will be established in Chap. 2.

Set-Theoretic Notations and Terminology

1.1 Some sets can be described by listing their members. Thus $\{x_1, \dots, x_n\}$ is the set whose members are $x_1, \dots, x_n$; and $\{x\}$ is the set whose only member is x . More often, sets are described by properties. We write

$$\{x: P\}$$

for the set of all elements x which have the property P . The symbol $\emptyset$ denotes the empty set. The words *collection*, *family*, and *class* will be used synonymously with *set*.

We write $x \in A$ if x is a member of the set A ; otherwise $x \notin A$. If B is a subset of A , i.e., if $x \in B$ implies $x \in A$, we write $B \subset A$. If $B \subset A$ and $A \subset B$, then $A = B$. If $B \subset A$ and $A \neq B$, B is a *proper* subset of A . Note that $\emptyset \subset A$ for every set A .

$A \cup B$ and $A \cap B$ are the union and intersection of A and B , respectively. If $\{A_\alpha\}$ is a collection of sets, where α runs through some index set I , we write

$$\bigcup_{\alpha \in I} A_\alpha \quad \text{and} \quad \bigcap_{\alpha \in I} A_\alpha.$$

for the union and intersection of $\{A_\alpha\}$:

$$\begin{aligned} \bigcup_{\alpha \in I} A_\alpha &= \{x: x \in A_\alpha \text{ for at least one } \alpha \in I\} \\ \bigcap_{\alpha \in I} A_\alpha &= \{x: x \in A_\alpha \text{ for every } \alpha \in I\}. \end{aligned}$$

If I is the set of all positive integers, the customary notations are

$$\bigcup_{n=1}^{\infty} A_n \quad \text{and} \quad \bigcap_{n=1}^{\infty} A_n.$$