


Principles of Optics

光学原理

第7版



7th (expanded) edition

**Max Born and
Emil Wolf**

CAMBRIDGE

世界图书出版公司

Principles of optics

*Electromagnetic theory of propagation,
interference and diffraction of light*

MAX BORN

MA, Dr Phil, FRS

Nobel Laureate

Formerly Professor at the Universities of Göttingen and Edinburgh

and

EMIL WOLF

PhD, DSc

Wilson Professor of Optical Physics, University of Rochester, NY

with contributions by

A.B. BHATIA, P.C. CLEMMOW, D. GABOR, A.R. STOKES,
A.M. TAYLOR, P.A. WAYMAN AND W.L. WILCOCK

SEVENTH (EXPANDED) EDITION

CAMBRIDGE
UNIVERSITY PRESS

世界图书出版公司

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK www.cup.cam.ac.uk
40 West 20th Street, New York, NY 10011-4211, USA www.cup.org
10 Stamford Road, Oakleigh, Melbourne 3166, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain

Seventh (expanded) edition © Margaret Farley-Born and Emil Wolf 1999

First published 1959 by Pergamon Press Ltd, London
Sixth edition 1980
Reprinted (with corrections) 1983, 1984, 1986, 1987, 1989, 1991, 1993
Reissued by Cambridge University Press 1997
Seventh (expanded) edition 1999

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this book is available from the British Library

Library of Congress Cataloguing in Publication data

Born, Max
Principles of Optics—7th ed.
I. Optics
I. Title II. Wolf Emil
535 QC351 80-41470

ISBN 0 521 642221 hardback

This edition of *Principles of Optics* 7th Edition by Max Born and Emil Wolf is published by arrangement with the Syndicate of the Press of University of Cambridge, Cambridge, England.

Licensed edition for sale in the People's Republic of China only. Not for export elsewhere.

Principles of optics

Principles of Optics is one of the classic science books of the twentieth century, and probably the most influential book in optics published in the past 40 years. The new edition is the first ever thoroughly revised and expanded edition of this standard text.

Among the new material, much of which is not available in any other optics text, is a section on the CAT scan (computerized axial tomography), which has revolutionized medical diagnostics. The book also includes a new chapter on scattering from inhomogeneous media which provides a comprehensive treatment of the theory of scattering of scalar as well as of electromagnetic waves, including the Born series and the Rytov series. The chapter also presents an account of the principles of diffraction tomography – a refinement of the CAT scan – to which Emil Wolf, one of the authors, has made a basic contribution by formulating in 1969 what is generally regarded to be the basic theorem in this field. The chapter also includes an account of scattering from periodic potentials and its connection to the classic subject of determining the structure of crystals from X-ray diffraction experiments, including accounts of von Laue equations, Bragg's law, the Ewald sphere of reflection and the Ewald limiting sphere, both generalized to continuous media. These topics, although originally introduced in connection with the theory of X-ray diffraction by crystals, have since become of considerable relevance to optics, for example in connection with deep holograms.

Other new topics covered in this new edition include interference with broad-band light, which introduces the reader to an important phenomenon discovered relatively recently by Emil Wolf, namely the generation of shifts of spectral lines and other modifications of spectra of radiated fields due to the state of coherence of a source. There is also a section on the so-called Rayleigh–Sommerfeld diffraction theory which, in recent times, has been finding increasing popularity among optical scientists. There are also several new appendices, including one on energy conservation in scalar wavefields, which is seldom discussed in books on optics.

The new edition of this standard reference will continue to be invaluable to advanced undergraduates, graduate students and researchers working in most areas of optics.

**To the Memory of
Sir Ernest Oppenheimer**

Preface to the seventh edition

Forty years ago this month Max Born and I dispatched to the publishers the Preface to the first edition of *Principles of Optics*. Since that time the book has been published in six editions and has been reprinted seventeen times (not counting unauthorized editions and several translations), usually with only a small number of corrections. A recent change to a new publisher, who expressed willingness to reset the whole text, has given me the opportunity to make more substantial changes.

The first edition was published a year before the invention of the laser, an event which triggered an explosion of activities in optics and soon led to the creation of entirely new fields, such as non-linear optics, fiber optics and opto-electronics. Numerous applications followed, in medicine, in optical data storage, in information transfer and in many other areas. On a more fundamental level, quantum optics emerged as a vibrant and rapidly expanding field, which has provided new ways of testing some basic assumptions of quantum physics relating, for example, to localization and indistinguishability. The progress made in these fields has been rapid and broad and some of the newer areas have themselves become the subjects of books.

It is clear that a fully updated new edition of *Principles of Optics* would require that it be expanded into several volumes. Consequently, in order to preserve a single-volume, only a few new topics have been added; they were selected to some extent so as not to necessitate major revisions of the original text. Specifically, the following new material has been added:

- (1) Section 4.11, which presents the principles of computerized axial tomography, generally referred to as CAT. This subject originated in the early 1960s and has revolutionized diagnostic medicine. The section also includes an account of the Radon transform, introduced as early as 1917, which underlies the theory of computerized axial tomography, although this was not known to its inventors. The fact that three Nobel Prizes have been awarded for this invention and its applications attests to its importance. More recently the theory underlying the CAT scan is finding much broader usage, for example in connection with the reconstruction of quantum states.
- (2) Section 8.11 gives an account of the so-called Rayleigh–Kirchhoff diffraction theory. This theory has become rather popular after it was introduced in the book *Optics* by A. Sommerfeld, published in 1954. It is preferred by some optical scientists to the much older classic Kirchhoff diffraction theory. However, which of these theories better describes various diffraction effects is still an open question.
- (3) Section 10.5 discusses some effects discovered relatively recently arising on

superposition of broad-band light beams of any state of coherence. The analysis of such effects has demonstrated that even though under such circumstances interference fringes may not be seen in the region of superposition, the light distribution in that region may nevertheless contain important physical information which is revealed when the light is spectrally analyzed. One may then find that the spectrum of the light is different at different points of observation and from such spectral changes one may determine coherence properties of the light. This effect is an example of a coherence phenomenon in the space-frequency domain, which must be distinguished from the more familiar coherence effects in the space-time domain. The quantitative measure of space-frequency coherence phenomena is the so-called spectral degree of coherence, which is introduced in this new edition in the somewhat revised Section 10.5. The experiment which is analyzed in that section also provides an elementary introduction to the phenomenon of correlation-induced spectral changes, discovered just over 10 years ago and studied extensively since that time.

(4) A new Chapter 13 presents the theory of scattering of light by inhomogeneous media. In the context of optics this subject was largely developed in relatively recent times although essentially the same theory was well established many years ago in connection with quantum mechanical potential scattering. The chapter presents the basic integral equation of light scattering on linear isotropic bodies, discusses the solution of the equation in series form and includes a detailed account of the first Born and the first Rytov approximations. The chapter includes a brief description of the classic theory of scattering by a medium with periodic structure, which is the basis of the theory of X-ray diffraction by crystals. The chapter also covers the von Laue equations, Bragg's law, the Ewald sphere of reflection and the Ewald limiting sphere. In recent years these subjects have become important in the broad area of inverse light scattering and they have found new applications, for example in connection with holographic gratings. The chapter also contains a detailed account of the optical cross-section theorem (usually known as the optical theorem), which, in spite of its name, is seldom discussed in the optical literature.

Another topic treated in Chapter 13 is diffraction tomography. In computerized axial tomography discussed in the new Section 4.11, the finite wavelength of the radiation is ignored. This is usually justified when the technique is used with X-rays, but the approximation is often inadequate in applications where light waves or sound waves are used. Diffraction tomography takes the finite wavelength of the radiation into account and, therefore, provides better resolution.

In addition to the new material already mentioned, this new edition also contains several new appendices (VIII, XI and XII), many new references, some of which replace older ones, and a few relatively small changes have been made in the text, usually with the aim of improving clarity and updating information.

It is a pleasure to thank several colleagues, friends and students for their help. I am obliged to Professor Harrison H. Barrett for helpful suggestions which led to improvements of Sections 4.11 and 13.2 on computerized axial tomography and on diffraction tomography. I am indebted to my long-time friend and colleague Professor Leonard Mandel for useful comments on some parts of the manuscript. Dr Doo Cho and Dr V. N. Mahajan were kind enough to draw my attention to some inaccuracies which they found in earlier editions and which have now been taken care of. My collaborators Dr

Yajun Li, Dr Taco D. Visser and my former students Dr Avshalom Gamliel, Dr David G. Fischer and Dr Kisik Kim have read some of the new sections and have made helpful comments leading to improvements. My former student Dr Weijian Wang kindly prepared most of the new figures. I am particularly grateful to two of my present graduate students, P. Scott Carney and Greg J. Gbur, who provided invaluable help by checking calculations and references, weeding out inaccuracies, suggesting improvements in the presentation and helping with proof-reading.

I wish to express my appreciation to Mrs Patricia Sulouff, Head Librarian of the Physics–Optics–Astronomy library at the University of Rochester for helping to trace some of the more obscure references and for locating papers and books that were not easily accessible. I am also obliged to Mrs Ellen Calkins for typing and re-typing several drafts of the manuscript of the new sections and for preparing the author index.

Most of the revisions were prepared whilst I was on a visiting appointment at the Center for Research and Education in Optics and Research (CREOL) at the University of Central Florida during the 1998 Spring Semester. I wish to thank Professors M. J. Soileau, M. G. Moharam and G. I. Stegeman of the CREOL faculty for providing a congenial environment and facilities for this work.

The family life of most authors suffers during the time when one of the marriage partners is engaged in book writing. Mine was no exception, but I am glad to say that my wife, Marlies, survived this period cheerfully and without complaints. She has helped with checking the manuscript and the proofs, for which I am grateful.

I acknowledge with thanks the fine cooperation that I received from the staff of Cambridge University Press. I am particularly appreciative of the considerable help provided by Dr Simon Capelin, the publishing director for Physical Sciences. I am also obliged to Mrs Maureen Storey for thorough copy-editing of the whole book and to Ms Miranda Fyfe and particularly to Mrs Jayne Aldhouse for their cooperation in meeting the rather stringent production deadlines.

I am sometimes asked about my collaboration with Max Born, which resulted in the publication of *Principles of Optics*. Interested readers can find it discussed in my article ‘Recollections of Max Born,’ published in *Optics News* 9, 10–16 (November/December, 1983).*

Rochester, New York
January 1999

Emil Wolf

*The article has been reprinted in *Technology of Our Times*, ed. F. Sue (SPIE Optical Engineering Press, Bellington, WA, 1990), 15–26 and also in *Astrophysics and Space Science* 227, ed. A. L. Peratt (Kluwer Academic Publishers, Dordrecht, The Netherlands, 1995), 277–97.

Preface to the first edition

THE idea of writing this book was a result of frequent enquiries about the possibility of publishing in the English language a book on optics written by one of us* more than twenty-five years ago. A preliminary survey of the literature showed that numerous researches on almost every aspect of optics have been carried out in the intervening years, so that the book no longer gives a comprehensive and balanced picture of the field. In consequence it was felt that a translation was hardly appropriate; instead a substantially new book was prepared, which we are now placing before the reader. In planning this book it soon became apparent that even if only the most important developments which took place since the publication of *Optik* were incorporated, the book would become impracticably large. It was, therefore, deemed necessary to restrict its scope to a narrower field. *Optik* itself did not treat the whole of optics. The optics of moving media, optics of X-rays and γ -rays, the theory of spectra and the full connection between optics and atomic physics were not discussed; nor did the old book consider the effects of light on our visual sense organ – the eye. These subjects can be treated more appropriately in connection with other fields such as relativity, quantum mechanics, atomic and nuclear physics, and physiology. In this book not only are these subjects excluded, but also the classical molecular optics which was the subject matter of almost half of the German book. Thus our discussion is restricted to those optical phenomena which may be treated in terms of Maxwell's phenomenological theory. This includes all situations in which the atomistic structure of matter plays no decisive part. The connection with atomic physics, quantum mechanics, and physiology is indicated only by short references wherever necessary. The fact that, even after this limitation, the book is much larger than *Optik*, gives some indication about the extent of the researches that have been carried out in classical optics in recent times.

We have aimed at giving, within the framework just outlined, a reasonably complete picture of our present knowledge. We have attempted to present the theory in such a way that practically all the results can be traced back to the basic equations of Maxwell's electromagnetic theory, from which our whole consideration starts.

In Chapter I the main properties of the electromagnetic field are discussed and the effect of matter on the propagation of the electromagnetic disturbance is described formally, in terms of the usual material constants. A more physical approach to the

* Max Born, *Optik* (Berlin, Springer, 1933).

question of influence of matter is developed in Chapter II: it is shown that in the presence of an external incident field, each volume element of a material medium may be assumed to give rise to a secondary (scattered) wavelet and that the combination of these wavelets leads to the observable, macroscopic field. This approach is of considerable physical significance and its power is illustrated in a later chapter (Chapter XII) in connection with the diffraction of light by ultrasonic waves, first treated in this way by A. B. Bhatia and W. J. Noble; Chapter XII was contributed by Prof. Bhatia himself.

A considerable part of Chapter III is devoted to showing how geometrical optics follows from Maxwell's wave theory as a limiting case of short wavelengths. In addition to discussing the main properties of rays and wave-fronts, the vectorial aspects of the problem (propagation of the directions of the field vectors) are also considered. A detailed discussion of the foundations of geometrical optics seemed to us desirable in view of the important developments made in recent years in the related field of microwave optics (optics of short radio waves). These developments were often stimulated by the close analogy between the two fields and have provided new experimental techniques for testing the predictions of the theory. We found it convenient to separate the mathematical apparatus of geometrical optics – the calculus of variations – from the main text; an appendix on this subject (Appendix I) is based in the main part on unpublished lectures given by D. Hilbert at Göttingen University in the early years of this century. The following appendix (Appendix II), contributed by Prof. D. Gabor, shows the close formal analogy that exists between geometrical optics, classical mechanics, and electron optics, when these subjects are presented in the language of the calculus of variations.

We make no apology for basing our treatment of geometrical theory of imaging (Chapter IV) on Hamilton's classical methods of characteristic functions. Though these methods have found little favour in connection with the design of optical instruments, they represent nevertheless an essential tool for presenting in a unified manner the many diverse aspects of the subject. It is, of course, possible to derive some of the results more simply from *ad hoc* assumptions; but, however valuable such an approach may be for the solution of individual problems, it cannot have more than illustrative value in a book concerned with a systematic development of a theory from a few simple postulates.

The defect of optical images (the influence of aberrations) may be studied either by geometrical optics (appropriate when the aberrations are large), or by diffraction theory (when they are sufficiently small). Since one usually proceeds from quite different starting points in the two methods of treatment, a comparison of results has in the past not always been easy. We have attempted to develop a more unified treatment, based on the concept of the deformation of wave-fronts. In the geometrical analysis of aberrations (Chapter V) we have found it possible and advantageous to follow, after a slight modification of his eikonal, the old method of K. Schwarzschild. The chapter on diffraction theory of aberrations (Chapter IX) gives an account of the Nijboer–Zernike theory and also includes an introductory section on the imaging of extended objects, in coherent and in incoherent illumination, based on the techniques of Fourier transforms.

Chapter VI, contributed by Dr P. A. Wayman, gives a brief description of the main image-forming optical systems. Its purpose is to provide a framework for those parts of the book which deal with the theory of image formation.

Chapter VII is concerned with the elements of the theory of interference and with interferometers. Some of the theoretical sections have their nucleus in the corresponding sections of *Optik*, but the chapter has been completely re-written by Dr W. L. Wilcock, who has also considerably broadened its scope.

Chapter VIII is mainly concerned with the Fresnel–Kirchhoff diffraction theory and with some of its applications. In addition to the usual topics, the chapter includes a detailed discussion of the central problem of optical image formation – the analysis of the three-dimensional light distribution near the geometrical focus. An account is also given of a less familiar alternative approach to diffraction, based on the notion of the boundary diffraction wave of T. Young.

The chapters so far referred to are mainly concerned with perfectly monochromatic (and therefore completely coherent) light, produced by point sources. Chapter X deals with the more realistic case of light produced by sources of finite extension and covering a finite frequency range. This is the subject of partial coherence, where considerable progress has been made in recent years. In fact, a systematic theory of interference and diffraction with partially coherent light has now been developed. This chapter also includes an account of the closely related subject of partial polarization, from the standpoint of coherence theory.

Chapter XI deals with rigorous diffraction theory, a field that has witnessed a tremendous development over the period of the last twenty years,* stimulated largely by advances in the ultra-shortwave radio techniques. This chapter was contributed by Dr P. C. Clemmow who also prepared Appendix III, which deals with the mathematical methods of steepest descent and stationary phase.

The last two chapters, ‘Optics of metals’ (Chapter XIII) and ‘Optics of crystals’ (Chapter XIV) are based largely on the corresponding chapters of *Optik*, but were revised and extended with the help of Prof. A. M. Taylor and Dr A. R. Stokes respectively. These two subjects are perhaps discussed in less detail than might seem appropriate. However, the optics of metals can only be treated adequately with the help of quantum mechanics of electrons, which is outside the scope of this book. In crystal optics the centre of interest has gradually shifted from visible radiation to X-rays, and the progress made in recent years has been of a technical rather than theoretical nature.

Though we have aimed at producing a book which in its methods of presentation and general approach would be similar to *Optik*, it will be evident that the present book is neither a translation of *Optik*, nor entirely a compilation of known data. As regards our own share in its production, the elder coauthor (M. B.) has contributed that material from *Optik* which has been used as a basis for some of the chapters in the present treatise, and has taken an active part in the general planning of the book and in numerous discussions concerning disputable points, presentation, etc. Most of the compiling, writing and checking of the text was done by the younger coauthor (E. W.).

Naturally we have tried to use systematic notation throughout the book. But in a book that covers such a wide field, the number of letters in available alphabets is far too limited. We have, therefore, not always been able to use the most elegant notation but we hope that we have succeeded, at least, in avoiding the use in any one section of the same symbol for different quantities.

* The important review article by C. J. Bouwkamp, *Rep. Progr. Phys.* (London, Physical Society), 17 (1954), 35, records more than 500 papers published in the period 1940–1954.

In general we use vector notation as customary in Great Britain. After much reflection we rejected the use of the nabla operator alone and employed also the customary 'div', 'grad' and 'curl'. Also, we did not adopt the modern electrotechnical units, as their main advantage lies in connection with purely electromagnetic measurements, and these play a negligible part in our discussions; moreover, we hope, that if ever a second volume (*Molecular and Atomic Optics*) and perhaps a third volume (*Quantum Optics*) is written, the C.G.S. system, as used in theoretical physics, will have returned to favour. Although, in this system of units, the magnetic permeability μ of most substances differs inappreciably from unity at optical frequencies, we have retained it in some of the equations. This has the advantage of greater symmetry and makes it possible to derive 'dual' results by making use of the symmetry properties of Maxwell's equations. For time periodic fields we have used, in complex representation, the factor $\exp(-i\omega t)$ throughout.

We have not attempted the task of referring to all the relevant publications. The references that are given, and which, we hope, include the most important papers, are to help the reader to gain some orientation in the literature; an omission of any particular reference should not be interpreted as due to our lack of regard for its merit.

In conclusion it is a pleasure to thank many friends and colleagues for advice and help. In the first place we wish to record our gratitude to Professor D. Gabor for useful advice and assistance in the early stages of this project, as well as for providing a draft concerning his ingenious method of reconstructed wave-fronts (§8.10). We are also greatly indebted to Dr F. Abelès, who prepared a draft, which is the backbone of §1.6, on the propagation of electromagnetic waves through stratified media, a field to which he himself has made a substantial contribution. We have also benefited by advice on this subject from Dr B. H. Billings.

We are much indebted to Dr H. H. Hopkins, Dr R. A. Silverman, Dr W. T. Welford and Dr G. Wyllie for critical comments and valuable advice, and to them and also to Dr G. Black, Dr H. J. J. Braddick, Dr N. Chako, Dr F. D. Kahn, Mr A. Nisbet, Dr M. Ross and Mr R. M. Sillitto for scrutinizing various sections of the manuscript. We are obliged to Polaroid Corporation for information concerning dichroic materials. Dr F. D. Kahn helped with proof-reading and Dr P. Roman and Mrs M. Podolanski with the preparation of the author index.

The main part of the writing was done at the Universities of Edinburgh and Manchester. The last stages were completed whilst one of the authors (E. W.) was a guest at the Institute of Mathematical Sciences, New York University. We are grateful to Professor M. Kline, Head of its Division of Electromagnetic Research, for his helpful interest and for placing at our disposal some of the technical facilities of the Institute.

We gratefully acknowledge the loan of original photographs by Professor M. Françon and Dr M. Cagnet (Figs 7.4, 7.26, 7.28, 7.60, 15.24, 15.26), Professor H. Lipson and his coworkers at the Manchester College of Science and Technology (Figs. 8.10, 8.12, 8.15), Dr O. W. Richards (Figs. 8.34, 8.35), and Professor F. Zernike and Dr K. Nienhus (Figs. 9.4, 9.5, 9.8, 9.10, 9.11). Fig. 7.66 is reproduced by courtesy of the Director of the Mount Wilson and Palomar Observatories. The blocks of Fig. 7.42 were kindly loaned by Messrs Hilger and Watts, Ltd, and those of Figs. 7.64 and 7.65 by Dr K. W. Meissner.

Financial assistance was provided by Messrs Industrial Distributors Ltd, London,

and we wish to acknowledge the generosity of the late Sir Ernest Oppenheimer, its former head.

Finally, it is a pleasure to thank our publishers and in particular Mr E. J. Buckley, Mr D. M. Lowe and also Dr P. Rosbaud, who as a former Director of Pergamon Press was closely associated with this project in its early stages, for the great care they have taken in the production of the book. It is a pleasure to pay tribute also to the printers, Pitman Press of Bath, for the excellence of their printing.

Bad Pyrmont and Manchester
January 1959

Max Born
Emil Wolf

Preface to the second edition

ADVANTAGE has been taken in the preparation of a new edition of this work to make a number of corrections of errors and misprints, to make a few minor additions and to include some new references.

Since the appearance of the first edition almost exactly three years ago, the first optical masers (lasers) have been developed. By means of these devices very intense and highly coherent light beams may be produced. Whilst it is evident that optical masers will prove of considerable value not only for optics but also for other sciences and for technology, no account of them is given in this new edition. For the basic principles of maser action have roots outside the domain of classical electromagnetic theory on which considerations of this book are based. We have, however, included a few references to recent researches in which light generated by optical masers was utilized or which have been stimulated by the potentialities of these new optical devices.

We wish to acknowledge our gratitude to a number of readers who drew our attention to errors and misprints. We are also obliged to Dr B. Karczewski and Mr C. L. Mehta for assistance with the revisions.

Bad Pyrmont and Rochester
November 1962

M.B.
E.W.

Preface to the sixth edition

THIS edition differs from its immediate predecessor chiefly in that it contains corrections of a small number of errors and misprints.

Rochester
September 1985

E.W.

Historical introduction

THE physical principles underlying the optical phenomena with which we are concerned in this treatise were substantially formulated before 1900. Since that year, optics, like the rest of physics, has undergone a thorough revolution by the discovery of the quantum of energy. While this discovery has profoundly affected our views about the nature of light, it has not made the earlier theories and techniques superfluous; rather, it has brought out their limitations and defined their range of validity. The extension of the older principles and methods and their applications to very many diverse situations has continued, and is continuing with undiminished intensity.

In attempting to present in an orderly way the knowledge acquired over a period of several centuries in such a vast field it is impossible to follow the historical development, with its numerous false starts and detours. It is therefore deemed necessary to record separately, in this preliminary section, the main landmarks in the evolution of ideas concerning the nature of light.*

The philosophers of antiquity speculated about the nature of light, being familiar with burning glasses, with the rectilinear propagation of light, and with refraction and reflection. The first systematic writings on optics of which we have any definite knowledge are due to the Greek philosophers and mathematicians [Empedocles (c. 490–430 BC), Euclid (c. 300 BC)].

Amongst the founders of the new philosophy, René Descartes (1596–1650) may be singled out for mention as having formulated views on the nature of light on the basis of his metaphysical ideas.† Descartes considered light to be essentially a pressure transmitted through a perfectly elastic medium (the aether) which fills all space, and he attributed the diversity of colours to rotary motions with different velocities of the particles in this medium. But it was only after Galileo Galilei (1564–1642) had, by his

* For a more extensive account of the history of optics, reference may be made to: J. Priestley, *History and Present State of Discoveries relating to Vision, Light and Colours* (2 Vols., London, 1772); Thomas Young, *A Course of Lectures on Natural Philosophy and the Mechanical Arts* Vol. I (London, 1845), pp. 374–385; E. Wilde, *Geschichte der Optik vom Ursprung dieser Wissenschaft bis auf die gegenwärtige Zeit*, 2 Vols. (Berlin, 1838, 1843); Ernst Mach, *The Principles of Physical Optics*, A historical and philosophical treatment (First German edition 1913. English translation 1926, reprinted by Dover Publications, New York, 1953); E. Hoppe, *Geschichte der Optik* (Leipzig, Weber, 1926); V. Ronchi, *Storia della Luce* (Bologna: Zanichelli, 2nd. Ed., 1952). A comprehensive historical account up to recent times is E. T. Whittaker's *A History of the Theories of Aether and Electricity*, Vol. I (*The Classical Theories*), revised and enlarged edition 1952; Vol. II (*The Modern Theories 1900–1926*), 1953, published by T. Nelson and Sons, London and Edinburgh. The first volume was used as the chief source for this introductory section.

† R. Descartes, *Dioptrique, Météores* (published (anonymously) in Leyden in 1637 with prefatory essay 'Discours de la méthode'). *Principia Philosophiae* (Amsterdam, 1644).

development of mechanics, demonstrated the power of the experimental method that optics was put on a firm foundation. The law of reflection was known to the Greeks; the law of refraction was discovered experimentally in 1621 by Willebrord Snell* (Snellius, c. 1580–1626). In 1657 Pierre de Fermat (1601–1665) enunciated the celebrated *Principle of Least Time*† in the form ‘Nature always acts by the shortest course’. According to this principle, light always follows that path which brings it to its destination in the shortest time, and from this, in turn, and from the assumption of varying ‘resistance’ in different media, the law of refraction follows. This principle is of great philosophical significance, and because it seems to imply a teleological manner of explanation, foreign to natural science, it has raised a great deal of controversy.

The first phenomenon of interference, the colours exhibited by thin films now known as ‘Newton’s rings’, was discovered independently by Robert Boyle‡ (1627–1691) and Robert Hooke§ (1635–1703). Hooke also observed the presence of light in the geometrical shadow, the ‘diffraction’ of light but this phenomenon had been noted previously by Francesco Maria Grimaldi|| (1618–1663). Hooke was the first to advocate the view that light consists of rapid vibrations propagated instantaneously, or with a very great speed, over any distance, and believed that in an homogeneous medium every vibration will generate a sphere which will grow steadily.¶ By means of these ideas Hooke attempted an explanation of the phenomenon of refraction, and an interpretation of colours. But the basic quality of colour was revealed only when Isaac Newton (1642–1727) discovered** in 1666 that white light could be split up into component colours by means of a prism, and found that each pure colour is characterized by a specific refrangibility. The difficulties which the wave theory encountered in connection with the rectilinear propagation of light and of polarization (discovered by Huygens††) seemed to Newton so decisive that he devoted himself to the development of an emission (or corpuscular) theory, according to which light is propagated from a luminous body in the form of minute particles.

At the time of the publication of Newton’s theory of colour it was not known whether light was propagated instantaneously or not. The discovery of the finite speed of light was made in 1675 by Olaf Römer (1644–1710) from the observations of the eclipses of Jupiter’s satellites.‡‡

The wave theory of light which, as we saw, had Hooke amongst its first champions was greatly improved and extended by Christian Huygens†† (1629–1695). He enunciated the principle, subsequently named after him, according to which every point of the ‘aether’ upon which the luminous disturbance falls may be regarded as the centre of a new disturbance propagated in the form of spherical waves; these secondary waves

* Snell died in 1626 without making his discoveries public. The law was first published by Descartes in his *Dioptrique* without an acknowledgement to Snell, though it is generally believed that Descartes had seen Snell’s manuscript on this subject.

† In a letter to Cureau de la Chambre. It is published in *Oeuvres de Fermat*, Vol. 2 (Paris, 1891) p. 354.

‡ *The Philosophical Works of Robert Boyle* (abridged by P. Shaw), Vol. II (Second ed. London, 1738), p. 70.

§ R. Hooke, *Micrographia* (1665), 47.

|| F. M. Grimaldi, *Physico-Mathesis de lumine, coloribus, et iride* (Bologna, 1665).

¶ The early wave theories of Hooke and Huygens operate with single ‘pulses’ rather than with wave trains of definite wavelengths.

** I. Newton, *Phil. Trans.* No. 80 (Feb. 1672), 3075.

†† Chr. Huygens, *Traité de la lumière* (completed in 1678, published in Leyden in 1690).

‡‡ Olaf Römer, *Mém. de l’Acad. Sci. Paris*, 10 (1666–1699), 575; *J. de Sav.* (1676), 223.