

John W. Morgan

The Seiberg-Witten Equations
and Applications to the
Topology of Smooth
Four-Manifolds

塞伯格-威顿方程及其在
光滑四流形拓扑中的应用



PRINCETON UNIVERSITY PRESS

世界图书出版公司
www.wpcbj.com.cn



John W. Morgan



The Seiberg-Witten Equations
and Applications to the
Topology of Surfaces
Four-Manifolds

塞伯格-威顿方程及其在
四维流形拓扑中的应用



世界图书出版公司
www.wppub.com

The Seiberg-Witten Equations and Applications to the Topology of Smooth Four-Manifolds

by

John W. Morgan

Mathematical Notes 77

PRINCETON UNIVERSITY PRESS

PRINCETON, NEW JERSEY

1996

图书在版编目 (CIP) 数据

塞伯格-威顿方程及其在光滑四流形拓扑中的应用 = The Seiberg - Witten equations and applications to the topology of smooth four - manifolds; 英文/(美)摩根 (Morgan, J. W.) 著. —影印本. —北京: 世界图书出版公司北京公司, 2013. 10
ISBN 978 - 7 - 5100 - 7028 - 0

I. ①S… II. ①摩… III. ①映射—流形拓扑—英文 IV. ①O189

中国版本图书馆 CIP 数据核字 (2013) 第 249303 号

-
- 书 名: The Seiberg - Witten Equations and Applications to the Topology of Smooth Four - Manifolds
- 作 者: John W. Morgan
- 中 译 名: 塞伯格-威顿方程及其在光滑四流形拓扑中的应用
- 责任编辑: 高蓉 刘慧
-
- 出 版 者: 世界图书出版公司北京公司
- 印 刷 者: 三河市国英印务有限公司
- 发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)
- 联系电话: 010 - 64021602, 010 - 64015659
- 电子信箱: kjb@wpcbj.com.cn
-
- 开 本: 24 开
- 印 张: 6
- 版 次: 2014 年 3 月
- 版权登记: 图字: 01 - 2013 - 1437
-
- 书 号: 978 - 7 - 5100 - 7028 - 0 定 价: 39.00 元
-

The Seiberg-Witten Equations and Applications to the Topology of Smooth Four-Manifolds

Copyright © 1996 by Princeton University Press
All Rights Reserved

Princeton University Press books are printed on acid-free paper
and meet the guidelines for permanence and durability of the
Committee on Production Guidelines for Book Longevity
of the Council on Library Resources

The Princeton Mathematical Notes are edited by
Luis A. Caffarelli, John N. Mather, and Elias M. Stein

Library of Congress Cataloging-in-Publication Data

Morgan, John W., 1946–

The Seiberg-Witten equations and applications to the topology of smooth
four-manifolds / by John W. Morgan.

p. cm. — (Mathematical notes ; 44)

Includes bibliographical references.

ISBN 0-691-02597-5 (pb : alk. paper)

1. Four-manifolds (Topology) 2. Seiberg-Witten invariants.

3. Mathematical physics. I. Title.

II. Series : Mathematical notes (Princeton University Press) ; 44

QA613.2.M67 1995

514'.2—dc20 95-43748

All rights reserved. No part of this book may be reproduced or transmitted in any form or
by any means, electronic or mechanical, including photocopying, recording or by any
information storage and retrieval system, without permission in writing from the Publisher.

The Seiberg-Witten Equations and
Applications to the Topology of
Smooth Four-Manifolds

Contents

1	Introduction	1
2	Clifford Algebras and Spin Groups	5
2.1	The Clifford Algebras	5
2.2	The groups $Pin(V)$ and $Spin(V)$	9
2.3	Splitting of the Clifford Algebra	13
2.4	The complexification of the $Cl(V)$	16
2.5	The Complex Spin Representation	19
2.6	The Group $Spin^c(V)$	20
3	Spin Bundles and the Dirac Operator	23
3.1	Spin Bundles and Clifford Bundles	23
3.2	Connections and Curvature	28
3.3	The Dirac Operator	40
3.4	The Case of Complex Manifolds	48
4	The Seiberg-Witten Moduli Space	55
4.1	The Equations	55
4.2	Space of Configurations	57
4.3	Group of Changes of Gauge	59
4.4	The Action	59
4.5	The Quotient Space	60
4.6	The Elliptic Complex	65
5	Curvature Identities and Bounds	69
5.1	Curvature Identities	70
5.2	<i>A Priori</i> bounds	77
5.3	The Compactness of the Moduli Space	80

6	The Seiberg-Witten Invariant	87
6.1	The Statement	87
6.2	The Parametrized Moduli Space	88
6.3	Reducible Solutions	91
6.4	Compactness of the Perturbed Moduli Space	92
6.5	Variation of the Metric and Self-dual Two-form	94
6.6	Orientability of the Moduli Space	95
6.7	The Case when $b_2^+(X) > 1$	99
6.8	An Involution in the Theory	100
6.9	The Case when $b_2^+(X) = 1$	104
7	Invariants of Kähler Surfaces	109
7.1	The Equations over a Kähler Manifold	109
7.2	Holomorphic Description of the Moduli Space	112
7.3	Evaluation for Kähler Surfaces	118
7.4	Computation for Kähler Surfaces	122
7.5	Final Remarks	126
	Bibliography	127

Chapter 1

Introduction

Beginning with the groundbreaking work of Donaldson in about 1980 it became clear that gauge-theoretic invariants of principal bundles and connections were an important tool in the study of smooth four-dimensional manifolds. Donaldson showed the importance of the moduli space of anti-self-dual connections. The next fifteen years saw an explosion of work in this area leading to computations of Donaldson polynomial invariants for a wide class of four-dimensional manifolds, especially algebraic surfaces. These computations yielded many powerful topological consequences including, for example, the diffeomorphism classification of elliptic surfaces. For some of the results obtained by using these techniques see [1], [6], and [2].

Last fall, motivated by new work in quantum field theory, Seiberg and Witten [9], introduced a different gauge-theoretic invariant which they claimed should be closely related to Donaldson's invariants. Indeed they gave an explicit formula for the relationship of their invariant to Donaldson's. While the link claimed by Seiberg-Witten between their invariants and Donaldson's has not yet been established mathematically, it is clearly true and can be shown to hold in all computed examples. Nevertheless, one can forget this supposed link and work directly with the new invariants as a substitute for the anti-self-dual invariants. This has been the approach during the last year or so.

It was clear from the beginning that the new invariants would be easier to work with since they involved principal bundles with structure group the circle instead of the non-abelian groups such as $SU(2)$ which arise in Donaldson theory. The surprise was that such simple invariants could capture the subtlety that Donaldson's invariants revealed. But in short

order, Witten [17] and then Taubes, Kronheimer, and Mrowka showed that indeed these new invariants did capture these subtleties and that they were easier to compute, at least in many cases. They did this by explicitly solving the Seiberg-Witten equations over Kähler surfaces. (See [3] for one account of the results for Kähler surfaces.) There followed in quick succession a series of remarkable theorems, each extending in a different way partial results from Donaldson's anti-self-dual theory. In fact, conjectures which seemed reasonable from the perspective of Donaldson theory but technically difficult, if not unreachable within that theory, suddenly became the standard test cases for the power of the new invariants. One by one these conjectures were established – leaving only one classical conjecture outstanding. The remaining one, called the *11/8ths-Conjecture*, deals with the quadratic forms that arise as intersection forms of simply connected spin four-manifolds.

It is the purpose of these notes to lay the groundwork for the Seiberg-Witten theory and then to show how one computes these invariants for most Kähler surfaces. We begin with the basics of Clifford algebras, spin structures and their cousins $Spin^c$ -structures, spin representations, and spinor bundles. We then consider the Dirac operator on the spin bundles over a riemannian four-manifold. The connection with Kähler geometry is facilitated because of the close connection on a Kähler manifold between the Dirac operator and $\bar{\partial}$. All of this is examined in detail. The book of Lawson-Michelson [7] gives a complete introduction to this material as well as an elaborate treatment of many more topics in the spinor geometry.

We then exhibit the Seiberg-Witten equations and show how to use these equations to produce a finite dimensional manifold, the moduli space of solutions to these equations modulo changes of gauge, inside an infinite dimensional configuration space. The homology class of this moduli space is then the value of the Seiberg-Witten invariant. We establish analogues of familiar theorems from anti-self-dual theory. The equations defining the moduli space are elliptic modulo the group of changes of gauge. A generic perturbation of the equations leads to a smooth orientable moduli space whose dimension can be computed by the Atiyah-Singer index theorem. To orient this moduli space it suffices to choose an orientation of H_+^2 and H^1 of the underlying four-manifold. As long as $b_2^+ > 1$ the moduli space varies smoothly as we vary the metric and perturbation. When $b_2^+ = 1$ the generic perturbation yields a smooth moduli space, but there will be singularities when we vary the metric and perturbation. These singularities are reducible solutions to the equations. This leads to a chamber structure for the Seiberg-Witten invariants in case $b_2^+ = 1$, similar to what happens

in the anti-self-dual theory.

We go on to establish one special property of this theory, namely the compactness of the moduli space of solutions. This is a consequence of *a priori* bounds for the pointwise norms of solutions to the equations. This result has no analogue for the anti-self-dual equations. Much of the geometric richness and much of the technical complexity of the anti-self-dual theory is directly related to the non-compactness of the moduli space. For both better and worse, that is all missing here. The Seiberg-Witten moduli spaces are compact and vary by a compact bordism as we vary the metric.

Having presented all these technical results, it is now clear that the homology class of the moduli space in the ambient space of configurations is an invariant of the underlying smooth four-manifold and the isomorphism class of the $Spin^c$ structure on that manifold. By definition, the Seiberg-Witten invariant of the $Spin^c$ structure is this homology class. We finish by explicitly computing the moduli spaces of solutions on ‘most’ algebraic surfaces, leading immediately to a computation of the Seiberg-Witten invariant for any $Spin^c$ structure over the surface. There are some special cases that we do not treat. These can, however, be treated by an elaboration of the techniques that we introduce. A complete discussion is contained in [3].

There is much more to be said about the Seiberg-Witten equations and the invariants that are defined from these equations. There are analogues of the theorems for Kähler manifolds which hold for symplectic manifolds, see [12, 13, 14]. There are also gluing theorems which lead to Meyer-Vietoris principles for solutions to the equations, see [8]. These then lead to product formulas for the invariants, which is one approach to gaining a more topological understanding of the invariants. In spite of the striking progress over the last year, much remains to be done. It is my hope that these notes will serve as an introduction making it possible for more mathematicians to contribute to this progress.

These notes are a written version of lecture series given at Columbia University and Princeton University during the past year. I wish to thank the graduate students at both Universities who attended these lectures and who, by their comments and questions, helped shape these notes.

Chapter 2

Clifford Algebras and Spin Groups

For any $n > 2$ the orthogonal group $SO(n)$ has fundamental group $\mathbf{Z}/2\mathbf{Z}$ and hence has a universal covering group called $Spin(n)$ which is a non-trivial double covering. There is a beautiful algebraic construction which yields the Spin groups (and much more as well). This is the subject of Clifford algebras.

2.1 The Clifford Algebras

An example. Before delving into the complexities of Clifford algebras in general, let us consider a simple example. Consider the unit sphere S^3 inside the quaternion algebra $\mathbf{H}$. Multiplication of quaternions induces a group structure on S^3 . Let us consider the action of this group on $\mathbf{H}$ by conjugation

$$\begin{aligned} S^3 \times \mathbf{H} &\rightarrow \mathbf{H} \\ (\alpha, \lambda) &\mapsto \alpha \lambda \alpha^{-1}. \end{aligned}$$

This action preserves the norm, i.e., is an orthogonal action. It also leaves invariant the center of $\mathbf{H}$ which is $\mathbf{R} \subset \mathbf{H}$, and hence it leaves invariant the perpendicular complement to $\mathbf{R}$ which is the three-dimensional space $\mathbf{Im} \mathbf{H}$ of purely imaginary quaternions. Of course, $\mathbf{Im} \mathbf{H}$ is naturally identified with the Lie algebra of S^3 and the action we are considering is the adjoint action of S^3 on its Lie algebra.

It is an easy geometric exercise to see that a unit quaternion α , different from ± 1 , acts by conjugation on $\mathbf{H}$ preserving the complex plane $\mathbf{C}\alpha$ spanned by α and its perpendicular $\mathbf{C}\alpha j$. On the first complex plane conjugation by α acts trivially and on the second it acts by rotation through twice the angle θ between α and 1. It follows that the conjugation action of α on $\text{Im } \mathbf{H}$ leaves invariant the line tangent to the circle generated by α and acts by rotation through 2θ on the perpendicular complement and thus that every rotation of $\text{Im } \mathbf{H}$ is in the image of the representation

$$S^3 \rightarrow SO(\text{Im } \mathbf{H}) = SO(3).$$

It is also clear that the kernel of this representation is the intersection of S^3 with the center of $\mathbf{H}$, which is the center of S^3 and is $\{\pm 1\}$.

In this way we construct the double covering group of $SO(3)$ and identify it with the group of quaternions of length one under multiplication.

The definition of the Clifford algebra associated to an positive-definite inner product space. Now let us turn to the general case. Let V be a finite dimensional vector space over $\mathbf{R}$ with a positive definite inner product $\langle \cdot, \cdot \rangle$ leading to a norm denoted $\| \cdot \|$. We consider the tensor algebra

$$T(V) = \bigoplus_{n \geq 0} \underbrace{V \otimes \cdots \otimes V}_{n \text{ times}}$$

generated by V . It is an associative algebra with unit 1. The Clifford algebra $Cl(V)$ generated by V is the quotient of $T(V)$ by the two-sided ideal generated by all elements of the form

$$v \otimes v + \|v\|^2 1$$

for $v \in V$. Notice that the grading on $T(V)$ descends to a $\mathbf{Z}/2\mathbf{Z}$ grading on $Cl(V)$, giving a decomposition

$$Cl(V) = Cl_0(V) \oplus Cl_1(V)$$

where $Cl_0(V)$ is a subalgebra and $Cl_1(V)$ is a module over this subalgebra. Corresponding to this splitting, we can write any $v \in Cl(V)$ as $v_0 + v_1$. We denote by

$$\epsilon: Cl(V) \rightarrow Cl(V)$$

the algebra homomorphism that is multiplication by $+1$ on $Cl_0(V)$ and is multiplication by -1 on $Cl_1(V)$; i.e., $\epsilon(v_0 + v_1) = v_0 - v_1$.

Choosing an orthonormal basis $\{e_1, \dots, e_n\}$ for V we can write $Cl(V)$ in terms of generators and relations. Namely, $Cl(V)$ is the algebra over $\mathbf{R}$ generated by $\{e_1, \dots, e_n\}$ subject to the relations $e_i^2 = -1$ for all $i \leq n$ and $e_i \cdot e_j = -e_j \cdot e_i$ for all $i \neq j$. In particular, it follows that every element of $Cl(V)$ can be written uniquely as a sum of products of the form

$$e_{i_1} \cdots e_{i_t}$$

where $i_1 < \cdots < i_t$. Thus, the dimension of $Cl(V)$ as a real vector space is 2^d where d is the dimension of V over $\mathbf{R}$.

Examples. (i) Let $\mathbf{R}^n$ denote the usual Euclidean space of dimension n . Then $Cl(\mathbf{R}^1) = \mathbf{R}[x]/(x^2 + 1) \cong \mathbf{C}$. The subalgebra $Cl_0(\mathbf{R}^1)$ is identified with the reals and $Cl_1(\mathbf{R}^1)$ with the purely imaginary complex numbers.

(ii) Similarly, $Cl(\mathbf{R}^2)$ is the algebra generated by x, y subject to the relations

$$x^2 = -1; y^2 = -1; xy = -yx.$$

Hence, $Cl(\mathbf{R}^2)$ is isomorphic to the quaternion algebra $\mathbf{H}$. The subalgebra $Cl_0(\mathbf{R}^2)$ is generated by xy and can be identified with $\mathbf{C} \subset \mathbf{H}$.

(iii) $Cl(\mathbf{R}^3)$ is of dimension 8 over $\mathbf{R}$. It is generated by x, y, z with $x^2 = y^2 = z^2 = -1$ and $xy = -yx, xz = -zx, yz = -zy$. This algebra is isomorphic to $\mathbf{H} \oplus \mathbf{H}$. The isomorphism between $\mathbf{H}$ and the first factor, resp., the second factor, is given by sending $1, i, j, k$ to

$$\frac{1 + xyz}{2}, \frac{xy - z}{2}, \frac{yz - x}{2}, \frac{zx - y}{2},$$

resp., to

$$\frac{1 - xyz}{2}, \frac{xy + z}{2}, \frac{yz + x}{2}, \frac{zx + y}{2}$$

The subalgebra $Cl_0(\mathbf{R}^3)$ is identified with the diagonal copy of $\mathbf{H}$ in this decomposition.

(iv) For any inner product space V , we have an isomorphism of algebras $Cl(V) \cong Cl_0(V \oplus \mathbf{R})$. Letting e be a unit vector in $\mathbf{R}$, the isomorphism is given by

$$v_0 + v_1 \mapsto v_0 + v_1 e.$$

It is an easy exercise to show that this map is an isomorphism of algebras. In particular, $Cl_0(\mathbf{R}^4)$ is isomorphic to $\mathbf{H} \oplus \mathbf{H}$.

Comparison with the Exterior Algebra. The grading on $T(V)$ induces an increasing filtration

$$0 \subset \mathcal{F}_1 \subset \mathcal{F}_2 \subset \cdots$$

of $Cl(V)$ by linear subspaces; namely, we set $\mathcal{F}_t$ equal to the image in $Cl(V)$ of

$$\bigoplus_{n \leq t} \underbrace{V \otimes \cdots \otimes V}_{n \text{ times}}.$$

Clearly, the multiplication in $Cl(V)$ preserves this filtration in the sense that it induces maps

$$\mathcal{F}_i \otimes \mathcal{F}_j \rightarrow \mathcal{F}_{i+j}.$$

Thus, there is an associated graded algebra

$$\text{Gr}_{\mathcal{F}_*}(Cl(V)) = \bigoplus_{n=0}^{\infty} \mathcal{F}_n / \mathcal{F}_{n-1}$$

with the induced multiplication.

Claim 2.1.1 $\text{Gr}_{\mathcal{F}_*}(Cl(V))$ is naturally isomorphic to the exterior algebra $\Lambda^*(V)$.

Proof. Let $\tilde{\mathcal{F}}_*$ be the increasing filtration of $T(V)$ coming from the grading. Clearly, there is a natural algebra isomorphism between $T(V)$ and $\text{Gr}_{\tilde{\mathcal{F}}_*}(T(V))$. Consequently, we have a commutative diagram

$$\begin{array}{ccccccc} \mathcal{I}(v \otimes v + \|v\|^2 1) & \longrightarrow & T(V) & \longrightarrow & Cl(V) & \longrightarrow & 0 \\ \downarrow & & \downarrow \cong & & \downarrow & & \\ \mathcal{I}(v \otimes v) & \longrightarrow & \text{Gr}_{\tilde{\mathcal{F}}_*}(T(V)) & \longrightarrow & \text{Gr}_{\mathcal{F}_*}(Cl(V)) & \longrightarrow & 0. \end{array}$$

Clearly, the quotient of $T(V)$ by the two-sided ideal generated by $v \otimes v$ for $v \in V$ is exactly the exterior algebra $\Lambda^*(V)$. Thus, this diagram induces the claimed isomorphism. $\square$

There is a natural splitting σ of the map $Cl(V) \rightarrow \Lambda^*(V)$. The map σ is linear but not multiplicative. It is defined as follows. Consider an elementary element in $\Lambda^k V$ (; i.e., one contained in $\Lambda^k(W) \subset \Lambda^k(V)$ for some k -dimensional subspace $W \subset V$. Such an element can be written as $re_1 \wedge \cdots \wedge e_k$ where the e_i form an orthonormal basis for W and where $r > 0$. We define

$$\sigma(re_1 \wedge \cdots \wedge e_k) = re_1 \cdots e_k.$$