


概率论基础教程

(英文版·第9版)



(美) 谢尔登 M. 罗斯 (Sheldon M. Ross) 著
南加州大学



华章数学原版精品系列

概率论基础教程

(英文版·第9版)

A First Course in Probability
(Ninth Edition)

(美) 谢尔登 M. 罗斯 (Sheldon M. Ross) 著
南加州大学



机械工业出版社
China Machine Press

图书在版编目 (CIP) 数据

概率论基础教程 (英文版·第9版) / (美) 谢尔登 M. 罗斯 (Sheldon M. Ross) 著. —北京: 机械工业出版社, 2017.2

(华章数学原版精品系列)

书名原文: A First Course in Probability, Ninth Edition

ISBN 978-7-111-56148-4

I. 概… II. 谢… III. 概率论-教材-英文 IV. 0211

中国版本图书馆 CIP 数据核字 (2017) 第 032263 号

本书版权登记号: 图字: 01-2016-9715

Authorized Adaptation from the English Language edition, entitled *A First Course in Probability, Ninth Edition* (ISBN 9780321794772) by Sheldon M. Ross, Copyright © 2014, 2010, 2006 by Pearson Education, Inc.

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage retrieval system, without permission from Pearson Education, Inc.

English language adaptation edition published by Pearson Education Asia Ltd., and China Machine Press Copyright © 2017.

English language adaptation edition is manufactured in the People's Republic of China and is authorized for sale only in People's Republic of China excluding Taiwan, Hong Kong SAR and Macau SAR.

本书英文影印版由 Pearson Education Asia Ltd. 授权机械工业出版社独家出版。未经出版者书面许可, 不得以任何方式复制或抄袭本书内容。

仅限于中华人民共和国境内 (不包括香港、澳门特别行政区及台湾地区) 销售发行。

本书封面贴有 Pearson Education (培生教育出版集团) 激光防伪标签, 无标签者不得销售。

出版发行: 机械工业出版社 (北京市西城区百万庄大街 22 号 邮政编码: 100037)

责任编辑: 和 静

责任校对: 殷 虹

印 刷: 北京诚信伟业印刷有限公司

版 次: 2017 年 3 月第 1 版第 1 次印刷

开 本: 186mm×240mm 1/16

印 张: 29.75

书 号: ISBN 978-7-111-56148-4

定 价: 79.00 元

凡购本书, 如有缺页、倒页、脱页, 由本社发行部调换

客服热线: (010) 88378991 88361066

投稿热线: (010) 88379604

购书热线: (010) 68326294 88379649 68995259

读者信箱: hzjsj@hzbook.com

版权所有·侵权必究

封底无防伪标均为盗版

本书法律顾问: 北京大成律师事务所 韩光 / 邹晓东

前 言

“我们看到，概率论实际上只是将常识归结为计算，它使我们能够用理性的头脑精确地评价凭某种直觉感受到的、往往又不能解释清楚的见解……引人注意的是，概率论这门起源于对机会游戏进行思考的科学，早就应该成为人类知识中最重要的组成部分……生活中那些最重要的问题绝大部分其实只是概率论的问题。”著名的法国数学家和天文学家拉普拉斯侯爵（人称“法国的牛顿”）如是说。尽管许多人认为，这位对概率论的发展作出过重大贡献的著名侯爵说话夸张了一些，但是概率论已经成为几乎所有的科学工作者、工程师、医务人员、法律工作者和企业家们手中的基本工具，这是一个不争的事实。实际上，有见识的人们不再问：“是这样吗？”而是问：“有多大的概率是这样？”

一般方法和数学水平

本书是概率论的入门教材，适用于具备初等微积分知识的数学、统计、工程和其他学科（包括计算机科学、生物学、社会科学和管理科学）的学生。本书不仅介绍概率论的数学理论，而且通过大量例子来展示这门学科的广泛应用。

内容和课程计划

第1章阐述了组合分析的基本原理，它是计算概率的最有用的工具。

第2章介绍了概率论的公理体系，并且阐明如何应用这些公理进行概率计算。

第3章讨论概率论中极为重要的两个概念，即事件的条件概率和事件的独立性。通过一系列例子说明：当部分信息可利用时，条件概率就会起作用；即使在没有部分信息时，条件概率也可以使概率的计算变得容易。利用“条件”计算概率这一极为重要的技巧还将出现在第7章，在那里我们用它来计算期望。

第4~6章引入随机变量的概念。第4章讨论离散型随机变量，第5章讨论连续型随机变量，第6章讨论随机变量的联合分布。在第4章和第5章中讨论了两个重要概念，即随机变量的期望值和方差，并且对许多常见的随机变量求出了相应的期望值和方差。

第7章进一步讨论了期望值的一些重要性质。书中引入了许多例子，解释如何利用随机变量和的期望等于随机变量期望的和这一重要规律来计算随机变量的期望值。本章中还有几节介绍条件期望（包括它在预测方面的应用）和矩母函数。本章最后一节介绍了多元正态分布，同时给出了来自正态总体的样本均值和样本方差的联合分布的简单证明。

在第8章我们介绍了概率论的主要理论结果。特别地，我们证明了强大数定律和中心极限定理。在强大数定律的证明中，我们假定了随机变量具有有限的四阶矩，因为在这种假定之下，证明非常简单。在中心极限定理的证明中，我们假定了莱维连续性定理成立。在本章中，我们

还介绍了若干概率不等式，如马尔可夫不等式、切比雪夫不等式和切尔诺夫界。在本章最后一节，我们给出用有相同期望值的泊松随机变量的相应概率去近似独立伯努利随机变量和的相关概率的误差界。

第 9 章阐述了一些额外的论题，如马尔可夫链、泊松过程以及信息编码理论初步。第 10 章介绍了统计模拟。

与以前的版本一样，在每章末给出了三组练习题——习题、理论习题和自检习题。自检习题的全部解答在附录 B 给出，这部分练习题可以帮助学生检测他们对知识的掌握程度并为考试作准备。

第 9 版的特色

第 9 版继续对教材进行微调和优化，除了大量的小修改使得教材更加清晰外，本版还包括了很多新的或更新的练习题和正文内容，内容的选择不仅因为它们本身的趣味性，更是为了用它们来建立学生对概率的直觉。第 3 章的例 3h 和例 4k 就是这个目标的最好例证，例 3h 介绍双胞胎同卵的比例的估计，例 4k 分析发球和接球游戏。

致谢

我要感谢下面这些为了改进本教材而慷慨地与我联系并提出意见的人们：Amir Ardestani（德黑兰理工大学），Joe Blitzstein（哈佛大学），Peter Nuesch（洛桑大学），Joseph Mitchell（纽约州立大学石溪分校），Alan Chambless（精算师），Robert Kriner、Israel David（本-古里安大学），T.Lim（乔治梅森大学），Wei Chen（罗格斯大学），D.Monrad（伊利诺伊大学），W.Rosenberger（乔治梅森大学），E.Ionides（密歇根大学），J.Corvino（拉法叶学院），T.Seppalainen（威斯康星大学），Jack Goldberg（密歇根大学），Sunil Dhar（新泽西理工学院），Vladislav Kargin（斯坦福大学），Marlene Miller、Ahmad Parsian 和 Fritz Scholz（华盛顿大学）。

我也要特别感谢第 9 版的审查者：Richard Laugesen（伊利诺伊大学），Stacey Hancock（克拉克大学），Stefan Heinz（怀俄明大学），Brian Thelen（密歇根大学）。准确性的审查者 Keith Friedman（得克萨斯大学奥斯汀分校）和 Stacey Hancock（克拉克大学）非常仔细地审查了书稿内容，在此也要特别感谢他们。

最后，我要感谢下面这些审查者提出很有用的评论意见，其中第 9 版的审查者用星号标记。

K.B.Athreya（爱荷华州立大学）

Richard Bass（康涅狄格大学）

Robert Bauer（伊利诺伊大学厄巴纳-尚佩恩分校）

Phillip Beckwith（密歇根科技大学）

Arthur Benjamin（哈佛姆德学院）

Geoffrey Berresford（长岛大学）

Baidurya Bhattacharya（特拉华大学）

Howard Bird（圣克劳德州立大学）

Shahar Boneh（丹佛大都会州立学院）

Jean Cadet（纽约州立大学石溪分校）

Steven Chiappari（圣塔克拉拉大学）

Nicolas Christou（加州大学洛杉矶分校）

James Clay（亚利桑那大学图森分校）

Francis Conlan (圣克拉拉大学)
 Justin Corvino (拉法叶学院)
 Jay DeVore (加州州立理工大学圣路易斯奥比
 斯波分校)
 Scott Emerson (华盛顿大学)
 Thomas R.Fischer (德州农工大学)
 Anant Godbole (密歇根科技大学)
 Zakkula Govindarajulu (肯塔基大学)
 Richard Groeneveld (爱荷华州立大学)
 Stacey Hancock (克拉克大学)
 Mike Hardy (麻省理工学院)
 Bernard Harris (威斯康星大学)
 Larry Harris (肯塔基大学)
 David Heath (康奈尔大学)
 Stefan Heinz (怀俄明大学)
 Stephen Herschkorn (罗格斯大学)
 Julia L.Higle (亚利桑那大学)
 Mark Huber (杜克大学)
 Edward Ionides (密歇根大学)
 Anastasia Ivanova (北卡罗来纳大学)
 Hamid Jafarkhani (加州大学欧文分校)
 Chuanshu Ji (北卡罗来纳大学教堂山分校)
 Robert Keener (密歇根大学)
 Richard Laugesen (伊利诺伊大学)
 Fred Leysieffer (佛罗里达州立大学)
 Thomas Liggett (加州大学洛杉矶分校)
 Helmut Mayer (佐治亚大学)
 Bill McCormick (佐治亚大学)
 Ian McKeague (佛罗里达州立大学)
 R.Miller (斯坦福大学)
 Ditlev Monrad (伊利诺伊大学)
 Robb J.Muirhead (密歇根大学)
 Joe Naus (罗格斯大学)
 Nhu Nguyen (新墨西哥州立大学)
 Ellen O'Brien (乔治梅森大学)
 N.U.Prabhu (康奈尔大学)
 Kathryn Prewitt (亚利桑那州立大学)
 Jim Propp (威斯康星大学)
 William F.Rosenberger (乔治梅森大学)
 Myra Samuels (普度大学)
 I.R.Savage (耶鲁大学)
 Art Schwartz (密歇根大学安阿伯分校)
 Therese Shelton (西南大学)
 Malcolm Sherman(纽约州立大学奥尔巴尼分校)
 Murad Taqqu (波士顿大学)
 *Brian Thelen (密歇根大学)
 Eli Upfal (布朗大学)
 Ed Wheeler (田纳西大学)
 Allen Webster (布拉德利大学)

S. R. smross@usc.edu

Contents

Preface ix

1 COMBINATORIAL ANALYSIS 1

- 1.1 Introduction 1
- 1.2 The Basic Principle of Counting 2
- 1.3 Permutations 3
- 1.4 Combinations 5
- 1.5 Multinomial Coefficients 9
- 1.6 The Number of Integer Solutions of Equations 12

2 AXIOMS OF PROBABILITY 21

- 2.1 Introduction 21
- 2.2 Sample Space and Events 21
- 2.3 Axioms of Probability 25
- 2.4 Some Simple Propositions 28
- 2.5 Sample Spaces Having Equally Likely Outcomes 32
- 2.6 Probability as a Continuous Set Function 42
- 2.7 Probability as a Measure of Belief 46

3 CONDITIONAL PROBABILITY AND INDEPENDENCE 56

- 3.1 Introduction 56
- 3.2 Conditional Probabilities 56
- 3.3 Bayes's Formula 62
- 3.4 Independent Events 75
- 3.5 $P(\cdot|F)$ Is a Probability 89

4 RANDOM VARIABLES 112

- 4.1 Random Variables 112
- 4.2 Discrete Random Variables 116

- 4.3 Expected Value 119
- 4.4 Expectation of a Function of a Random Variable 121
- 4.5 Variance 125
- 4.6 The Bernoulli and Binomial Random Variables 127
- 4.7 The Poisson Random Variable 135
- 4.8 Other Discrete Probability Distributions 147
- 4.9 Expected Value of Sums of Random Variables 155
- 4.10 Properties of the Cumulative Distribution Function 159

5 CONTINUOUS RANDOM VARIABLES 176

- 5.1 Introduction 176
- 5.2 Expectation and Variance of Continuous Random Variables 179
- 5.3 The Uniform Random Variable 184
- 5.4 Normal Random Variables 187
- 5.5 Exponential Random Variables 197
- 5.6 Other Continuous Distributions 203
- 5.7 The Distribution of a Function of a Random Variable 208

6 JOINTLY DISTRIBUTED RANDOM VARIABLES 220

- 6.1 Joint Distribution Functions 220
- 6.2 Independent Random Variables 228
- 6.3 Sums of Independent Random Variables 239
- 6.4 Conditional Distributions: Discrete Case 248
- 6.5 Conditional Distributions: Continuous Case 250

- 6.6 Order Statistics 256
- 6.7 Joint Probability Distribution of Functions of Random Variables 260
- 6.8 Exchangeable Random Variables 267

7 PROPERTIES OF EXPECTATION 280

- 7.1 Introduction 280
- 7.2 Expectation of Sums of Random Variables 281
- 7.3 Moments of the Number of Events that Occur 298
- 7.4 Covariance, Variance of Sums, and Correlations 304
- 7.5 Conditional Expectation 313
- 7.6 Conditional Expectation and Prediction 330
- 7.7 Moment Generating Functions 334
- 7.8 Additional Properties of Normal Random Variables 345
- 7.9 General Definition of Expectation 349

8 LIMIT THEOREMS 367

- 8.1 Introduction 367
- 8.2 Chebyshev's Inequality and the Weak Law of Large Numbers 367

- 8.3 The Central Limit Theorem 370
- 8.4 The Strong Law of Large Numbers 378
- 8.5 Other Inequalities 382
- 8.6 Bounding the Error Probability When Approximating a Sum of Independent Bernoulli Random Variables by a Poisson Random Variable 388

9 ADDITIONAL TOPICS IN PROBABILITY 395

- 9.1 The Poisson Process 395
- 9.2 Markov Chains 397
- 9.3 Surprise, Uncertainty, and Entropy 402
- 9.4 Coding Theory and Entropy 405

10 SIMULATION 415

- 10.1 Introduction 415
- 10.2 General Techniques for Simulating Continuous Random Variables 417
- 10.3 Simulating from Discrete Distributions 424
- 10.4 Variance Reduction Techniques 426

Answers to Selected Problems 433

Solutions to Self-Test Problems and Exercises 435

目 录

前 言

第 1 章 组合分析	1
1.1 引言	1
1.2 计数基本法则	2
1.3 排列	3
1.4 组合	5
1.5 多项式系数	9
1.6 方程的整数解个数	12
第 2 章 概率论公理	21
2.1 引言	21
2.2 样本空间和事件	21
2.3 概率论公理	25
2.4 几个简单命题	28
2.5 等可能结果的样本空间	32
*2.6 概率: 连续集函数	42
2.7 概率: 确信程度的度量	46
第 3 章 条件概率和独立性	56
3.1 引言	56
3.2 条件概率	56
3.3 贝叶斯公式	62
3.4 独立事件	75
3.5 $P(\cdot F)$ 是概率	89
第 4 章 随机变量	112
4.1 随机变量	112
4.2 离散型随机变量	116
4.3 期望	119
4.4 随机变量函数的期望	121
4.5 方差	125
4.6 伯努利随机变量和二项随机变量	127

4.7 泊松随机变量	135
4.8 其他离散型概率分布	147
4.9 随机变量和的期望	155
4.10 分布函数的性质	159
第 5 章 连续型随机变量	176
5.1 引言	176
5.2 连续型随机变量的期望和方差	179
5.3 均匀随机变量	184
5.4 正态随机变量	187
5.5 指数随机变量	197
5.6 其他连续型概率分布	203
5.7 随机变量函数的分布	208
第 6 章 随机变量的联合分布	220
6.1 联合分布函数	220
6.2 独立随机变量	228
6.3 独立随机变量的和	239
6.4 离散情形下的条件分布	248
6.5 连续情形下的条件分布	250
*6.6 次序统计量	256
6.7 随机变量函数的联合分布	260
*6.8 可交换随机变量	267
第 7 章 期望的性质	280
7.1 引言	280
7.2 随机变量和的期望	281
7.3 试验序列中事件发生次数的矩	298
7.4 随机变量和的协方差、方差及 及相关系数	304
7.5 条件期望	313
7.6 条件期望及预测	330
7.7 矩母函数	334
7.8 正态随机变量的更多性质	345

7.9	期望的一般定义	349	9.2	马尔可夫链	397
第 8 章	极限定理	367	9.3	惊奇、不确定性及熵	402
8.1	引言	367	9.4	编码定理及熵	405
8.2	切比雪夫不等式及弱大数定律	367	第 10 章	模拟	415
8.3	中心极限定理	370	10.1	引言	415
8.4	强大数定律	378	10.2	模拟连续型随机变量的 一般方法	417
8.5	其他不等式	382	10.3	模拟离散分布	424
8.6	用泊松随机变量逼近独立的伯努 利随机变量和的概率误差界	388	10.4	方差缩减技术	426
第 9 章	概率论的其他课题	395	附录 A	部分习题答案	433
9.1	泊松过程	395	附录 B	自检习题解答	435

COMBINATORIAL ANALYSIS

Chapter

1

Contents

- | | |
|-------------------------------------|--|
| 1.1 Introduction | 1.5 Multinomial Coefficients |
| 1.2 The Basic Principle of Counting | 1.6 The Number of Integer Solutions of Equations |
| 1.3 Permutations | |
| 1.4 Combinations | |

1.1 Introduction

Here is a typical problem of interest involving probability: A communication system is to consist of n seemingly identical antennas that are to be lined up in a linear order. The resulting system will then be able to receive all incoming signals—and will be called *functional*—as long as no two consecutive antennas are defective. If it turns out that exactly m of the n antennas are defective, what is the probability that the resulting system will be functional? For instance, in the special case where $n = 4$ and $m = 2$, there are 6 possible system configurations, namely,

0 1 1 0
0 1 0 1
1 0 1 0
0 0 1 1
1 0 0 1
1 1 0 0

where 1 means that the antenna is working and 0 that it is defective. Because the resulting system will be functional in the first 3 arrangements and not functional in the remaining 3, it seems reasonable to take $\frac{3}{6} = \frac{1}{2}$ as the desired probability. In the case of general n and m , we could compute the probability that the system is functional in a similar fashion. That is, we could count the number of configurations that result in the system's being functional and then divide by the total number of all possible configurations.

From the preceding discussion, we see that it would be useful to have an effective method for counting the number of ways that things can occur. In fact, many problems in probability theory can be solved simply by counting the number of different ways that a certain event can occur. The mathematical theory of counting is formally known as *combinatorial analysis*.

1.2 The Basic Principle of Counting

The basic principle of counting will be fundamental to all our work. Loosely put, it states that if one experiment can result in any of m possible outcomes and if another experiment can result in any of n possible outcomes, then there are mn possible outcomes of the two experiments.

The basic principle of counting

Suppose that two experiments are to be performed. Then if experiment 1 can result in any one of m possible outcomes and if, for each outcome of experiment 1, there are n possible outcomes of experiment 2, then together there are mn possible outcomes of the two experiments.

Proof of the Basic Principle: The basic principle may be proven by enumerating all the possible outcomes of the two experiments; that is,

$$\begin{array}{c} (1,1), (1,2), \dots, (1,n) \\ (2,1), (2,2), \dots, (2,n) \\ \vdots \\ (m,1), (m,2), \dots, (m,n) \end{array}$$

where we say that the outcome is (i, j) if experiment 1 results in its i th possible outcome and experiment 2 then results in its j th possible outcome. Hence, the set of possible outcomes consists of m rows, each containing n elements. This proves the result.

Example 2a

A small community consists of 10 women, each of whom has 3 children. If one woman and one of her children are to be chosen as mother and child of the year, how many different choices are possible?

Solution By regarding the choice of the woman as the outcome of the first experiment and the subsequent choice of one of her children as the outcome of the second experiment, we see from the basic principle that there are $10 \times 3 = 30$ possible choices. ■

When there are more than two experiments to be performed, the basic principle can be generalized.

The generalized basic principle of counting

If r experiments that are to be performed are such that the first one may result in any of n_1 possible outcomes; and if, for each of these n_1 possible outcomes, there are n_2 possible outcomes of the second experiment; and if, for each of the possible outcomes of the first two experiments, there are n_3 possible outcomes of the third experiment; and if ..., then there is a total of $n_1 \cdot n_2 \cdots n_r$ possible outcomes of the r experiments.

Example 2b

A college planning committee consists of 3 freshmen, 4 sophomores, 5 juniors, and 2 seniors. A subcommittee of 4, consisting of 1 person from each class, is to be chosen. How many different subcommittees are possible?

Solution We may regard the choice of a subcommittee as the combined outcome of the four separate experiments of choosing a single representative from each of the classes. It then follows from the generalized version of the basic principle that there are $3 \times 4 \times 5 \times 2 = 120$ possible subcommittees. ■

**Example
2c**

How many different 7-place license plates are possible if the first 3 places are to be occupied by letters and the final 4 by numbers?

Solution By the generalized version of the basic principle, the answer is $26 \cdot 26 \cdot 26 \cdot 10 \cdot 10 \cdot 10 \cdot 10 = 175,760,000$. ■

**Example
2d**

How many functions defined on n points are possible if each functional value is either 0 or 1?

Solution Let the points be $1, 2, \dots, n$. Since $f(i)$ must be either 0 or 1 for each $i = 1, 2, \dots, n$, it follows that there are 2^n possible functions. ■

**Example
2e**

In Example 2c, how many license plates would be possible if repetition among letters or numbers were prohibited?

Solution In this case, there would be $26 \cdot 25 \cdot 24 \cdot 10 \cdot 9 \cdot 8 \cdot 7 = 78,624,000$ possible license plates. ■

1.3 Permutations

How many different ordered arrangements of the letters a, b , and c are possible? By direct enumeration we see that there are 6, namely, abc, acb, bac, bca, cab , and cba . Each arrangement is known as a *permutation*. Thus, there are 6 possible permutations of a set of 3 objects. This result could also have been obtained from the basic principle, since the first object in the permutation can be any of the 3, the second object in the permutation can then be chosen from any of the remaining 2, and the third object in the permutation is then the remaining 1. Thus, there are $3 \cdot 2 \cdot 1 = 6$ possible permutations.

Suppose now that we have n objects. Reasoning similar to that we have just used for the 3 letters then shows that there are

$$n(n - 1)(n - 2) \cdots 3 \cdot 2 \cdot 1 = n!$$

different permutations of the n objects.

Whereas $n!$ (read as “ n factorial”) is defined to equal $1 \cdot 2 \cdots n$ when n is a positive integer, it is convenient to define $0!$ to equal 1.

**Example
3a**

How many different batting orders are possible for a baseball team consisting of 9 players?

Solution There are $9! = 362,880$ possible batting orders. ■

**Example
3b**

A class in probability theory consists of 6 men and 4 women. An examination is given, and the students are ranked according to their performance. Assume that no two students obtain the same score.

(a) How many different rankings are possible?

- (b) If the men are ranked just among themselves and the women just among themselves, how many different rankings are possible?

Solution (a) Because each ranking corresponds to a particular ordered arrangement of the 10 people, the answer to this part is $10! = 3,628,800$.

(b) Since there are 6! possible rankings of the men among themselves and 4! possible rankings of the women among themselves, it follows from the basic principle that there are $(6!)(4!) = (720)(24) = 17,280$ possible rankings in this case. ■

Example
3c

Ms. Jones has 10 books that she is going to put on her bookshelf. Of these, 4 are mathematics books, 3 are chemistry books, 2 are history books, and 1 is a language book. Ms. Jones wants to arrange her books so that all the books dealing with the same subject are together on the shelf. How many different arrangements are possible?

Solution There are $4! 3! 2! 1!$ arrangements such that the mathematics books are first in line, then the chemistry books, then the history books, and then the language book. Similarly, for each possible ordering of the subjects, there are $4! 3! 2! 1!$ possible arrangements. Hence, as there are 4! possible orderings of the subjects, the desired answer is $4! 4! 3! 2! 1! = 6912$. ■

We shall now determine the number of permutations of a set of n objects when certain of the objects are indistinguishable from one another. To set this situation straight in our minds, consider the following example.

Example
3d

How many different letter arrangements can be formed from the letters *PEPPER*?

Solution We first note that there are $6!$ permutations of the letters $P_1E_1P_2P_3E_2R$ when the $3P$'s and the $2E$'s are distinguished from one another. However, consider any one of these permutations—for instance, $P_1P_2E_1P_3E_2R$. If we now permute the P 's among themselves and the E 's among themselves, then the resultant arrangement would still be of the form *PPEPER*. That is, all $3! 2!$ permutations

$$\begin{array}{ll} P_1P_2E_1P_3E_2R & P_1P_2E_2P_3E_1R \\ P_1P_3E_1P_2E_2R & P_1P_3E_2P_2E_1R \\ P_2P_1E_1P_3E_2R & P_2P_1E_2P_3E_1R \\ P_2P_3E_1P_1E_2R & P_2P_3E_2P_1E_1R \\ P_3P_1E_1P_2E_2R & P_3P_1E_2P_2E_1R \\ P_3P_2E_1P_1E_2R & P_3P_2E_2P_1E_1R \end{array}$$

are of the form *PPEPER*. Hence, there are $6!/(3! 2!) = 60$ possible letter arrangements of the letters *PEPPER*. ■

In general, the same reasoning as that used in Example 3d shows that there are

$$\frac{n!}{n_1! n_2! \cdots n_r!}$$

different permutations of n objects, of which n_1 are alike, n_2 are alike, \dots , n_r are alike.

Example
3e

A chess tournament has 10 competitors, of which 4 are Russian, 3 are from the United States, 2 are from Great Britain, and 1 is from Brazil. If the tournament result lists just the nationalities of the players in the order in which they placed, how many outcomes are possible?

Solution There are

$$\frac{10!}{4! 3! 2! 1!} = 12,600$$

possible outcomes. ■

Example 3f

How many different signals, each consisting of 9 flags hung in a line, can be made from a set of 4 white flags, 3 red flags, and 2 blue flags if all flags of the same color are identical?

Solution There are

$$\frac{9!}{4! 3! 2!} = 1260$$

different signals. ■

1.4 Combinations

We are often interested in determining the number of different groups of r objects that could be formed from a total of n objects. For instance, how many different groups of 3 could be selected from the 5 items $A, B, C, D,$ and E ? To answer this question, reason as follows: Since there are 5 ways to select the initial item, 4 ways to then select the next item, and 3 ways to select the final item, there are thus $5 \cdot 4 \cdot 3$ ways of selecting the group of 3 when the order in which the items are selected is relevant. However, since every group of 3—say, the group consisting of items $A, B,$ and C —will be counted 6 times (that is, all of the permutations $ABC, ACB, BAC, BCA, CAB,$ and CBA will be counted when the order of selection is relevant), it follows that the total number of groups that can be formed is

$$\frac{5 \cdot 4 \cdot 3}{3 \cdot 2 \cdot 1} = 10$$

In general, as $n(n-1) \cdots (n-r+1)$ represents the number of different ways that a group of r items could be selected from n items when the order of selection is relevant, and as each group of r items will be counted $r!$ times in this count, it follows that the number of different groups of r items that could be formed from a set of n items is

$$\frac{n(n-1) \cdots (n-r+1)}{r!} = \frac{n!}{(n-r)! r!}$$

Notation and terminology

We define $\binom{n}{r}$, for $r \leq n$, by

$$\binom{n}{r} = \frac{n!}{(n-r)! r!}$$

and say that $\binom{n}{r}$ (read as “ n choose r ”) represents the number of possible combinations of n objects taken r at a time.†

† By convention, $0!$ is defined to be 1. Thus, $\binom{n}{0} = \binom{n}{n} = 1$. We also take $\binom{n}{i}$ to be equal to 0 when either $i < 0$ or $i > n$.

Thus, $\binom{n}{r}$ represents the number of different groups of size r that could be selected from a set of n objects when the order of selection is not considered relevant.

Equivalently, $\binom{n}{r}$ is the number of subsets of size r that can be chosen from a set of size n . Using that $0! = 1$, note that $\binom{n}{n} = \binom{n}{0} = \frac{n!}{0!n!} = 1$, which is consistent with the preceding interpretation because in a set of size n there is exactly 1 subset of size n (namely, the entire set), and exactly one subset of size 0 (namely the empty set). A useful convention is to define $\binom{n}{r}$ equal to 0 when either $r > n$ or $r < 0$.

Example
4a

A committee of 3 is to be formed from a group of 20 people. How many different committees are possible?

Solution There are $\binom{20}{3} = \frac{20 \cdot 19 \cdot 18}{3 \cdot 2 \cdot 1} = 1140$ possible committees. ■

Example
4b

From a group of 5 women and 7 men, how many different committees consisting of 2 women and 3 men can be formed? What if 2 of the men are feuding and refuse to serve on the committee together?

Solution As there are $\binom{5}{2}$ possible groups of 2 women, and $\binom{7}{3}$ possible groups of 3 men, it follows from the basic principle that there are $\binom{5}{2} \binom{7}{3} = \frac{(5 \cdot 4)}{(2 \cdot 1)} \frac{7 \cdot 6 \cdot 5}{3 \cdot 2 \cdot 1} = 350$ possible committees consisting of 2 women and 3 men.

Now suppose that 2 of the men refuse to serve together. Because a total of $\binom{2}{2} \binom{5}{1} = 5$ out of the $\binom{7}{3} = 35$ possible groups of 3 men contain both of the feuding men, it follows that there are $35 - 5 = 30$ groups that do not contain both of the feuding men. Because there are still $\binom{5}{2} = 10$ ways to choose the 2 women, there are $30 \cdot 10 = 300$ possible committees in this case. ■

Example
4c

Consider a set of n antennas of which m are defective and $n - m$ are functional and assume that all of the defectives and all of the functionals are considered indistinguishable. How many linear orderings are there in which no two defectives are consecutive?

Solution Imagine that the $n - m$ functional antennas are lined up among themselves. Now, if no two defectives are to be consecutive, then the spaces between the functional antennas must each contain at most one defective antenna. That is, in the $n - m + 1$ possible positions—represented in Figure 1.1 by carets—between the $n - m$ functional antennas, we must select m of these in which to put the defective antennas. Hence, there are $\binom{n - m + 1}{m}$ possible orderings in which there is at least one functional antenna between any two defective ones. ■




Figure 1.1 No consecutive defectives.

A useful combinatorial identity is

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r} \quad 1 \leq r \leq n \quad (4.1)$$

Equation (4.1) may be proved analytically or by the following combinatorial argument: Consider a group of n objects, and fix attention on some particular one of these objects—call it object 1. Now, there are $\binom{n-1}{r-1}$ groups of size r that contain object 1 (since each such group is formed by selecting $r-1$ from the remaining $n-1$ objects). Also, there are $\binom{n-1}{r}$ groups of size r that do not contain object 1. As there is a total of $\binom{n}{r}$ groups of size r , Equation (4.1) follows.

The values $\binom{n}{r}$ are often referred to as *binomial coefficients* because of their prominence in the binomial theorem.

The binomial theorem

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} \quad (4.2)$$

We shall present two proofs of the binomial theorem. The first is a proof by mathematical induction, and the second is a proof based on combinatorial considerations.

Proof of the Binomial Theorem by Induction: When $n = 1$, Equation (4.2) reduces to

$$x + y = \binom{1}{0} x^0 y^1 + \binom{1}{1} x^1 y^0 = y + x$$

Assume Equation (4.2) for $n - 1$. Now,

$$\begin{aligned} (x + y)^n &= (x + y)(x + y)^{n-1} \\ &= (x + y) \sum_{k=0}^{n-1} \binom{n-1}{k} x^k y^{n-1-k} \\ &= \sum_{k=0}^{n-1} \binom{n-1}{k} x^{k+1} y^{n-1-k} + \sum_{k=0}^{n-1} \binom{n-1}{k} x^k y^{n-k} \end{aligned}$$