

Discriminants, Resultants, and Multidimensional Determinants

I. M. Gelfand
M. M. Kapranov
A. V. Zelevinsky

判别式、结式和多维行列式



Birkhäuser

世界图书出版公司
www.wpcbj.com.cn

Job #: 111365

Author Name: Gelfand

Title of Book: Discriminants, Resultants and Multidimen

ISBN #: 9780817647704



图书在版编目 (CIP) 数据

判别式结式和多维行列式 = Discriminants, Resultants, and Multidimensional Determinants: 英文/(美) 盖尔芬德 (Gelfand, I. M.) 著. — 影印本.

—北京: 世界图书出版公司北京公司, 2013. 5

ISBN 978 - 7 - 5100 - 6145 - 5

I. ①判… II. ①盖… III. ①结式—英文②行列式—英文 IV. ①O151. 1
②O151. 22

中国版本图书馆 CIP 数据核字 (2013) 第 088359 号

书 名: Discriminants, Resultants, and Multidimensional Determinants
作 者: I. M. Gelfand, M. M. Kapranov, A. V. Zelevinsky
中 译 名: 判别式、结式和多维行列式
责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司
印 刷 者: 三河市国英印务有限公司
发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)
联系电话: 010 - 64021602, 010 - 64015659
电子信箱: kjb@wpcbj. com. cn

开 本: 24 开
印 张: 22. 5
版 次: 2013 年 10 月
版权登记: 图字: 01 - 2013 - 8538

书 号: 978 - 7 - 5100 - 6145 - 5 定 价: 79.00 元

Preface

This book has expanded from our attempt to construct a general theory of hypergeometric functions and can be regarded as a first step towards its systematic exposition. However, this step turned out to be so interesting and important, and the whole program so overwhelming, that we decided to present it as a separate work. Moreover, in the process of writing we discovered a beautiful area which had been nearly forgotten so that our work can be regarded as a natural continuation of the classical developments in algebra during the 19th century.

We found that Cayley and other mathematicians of the period understood many of the concepts which today are commonly thought of as modern and quite recent. Thus, in an 1848 note on the resultant, Cayley in fact laid out the foundations of modern homological algebra. We were happy to enter into spiritual contact with this great mathematician.

The place of discriminants in the general theory of hypergeometric functions is similar to the place of quasi-classical approximation in quantum mechanics. More precisely, in [GGZ] [GKZ2] [GZK1] a general class of special functions was introduced and studied, the so-called A -hypergeometric functions. These functions satisfy a certain holonomic system of linear partial differential equations (the A -hypergeometric equations). The A -discriminant, which is one of our main objects of study, describes singularities of A -hypergeometric functions. According to the general principles of the theory of linear differential equations, these singularities are governed by the vanishing of the highest symbols of A -hypergeometric equations. The relation between differential operators and their highest symbols is the mathematical counterpart of the relation between quantum and classical mechanics; so we can say that hypergeometric functions provide a “quantization” of discriminants.

In our work on hypergeometric functions we found connections with many questions in algebra and combinatorics. We hope that this book brings to light some of these connections. One of the algebraic concepts which seems to us particularly important is that of hyperdeterminants (analogs of determinants for multi-dimensional “matrices.”) After rediscovering hyperdeterminants in connection with hypergeometric functions, we found that they too, had been introduced by Cayley in the 1840s. Unfortunately, later on, the study of hyperdeterminants was largely abandoned in favor of another, more straightforward definition (cf. [P]). The only other work on hyperdeterminants of which we are aware is an important

paper by Schläfli [Schl]. In this volume we give a detailed treatment of hyperdeterminants with the hope of attracting the attention of other mathematicians to this subject.

We would like to thank S.I. Gelfand, M.I. Graev and V.A. Vassiliev, who, through discussions and collaboration, have much influenced our understanding of the vast and beautiful field of hypergeometric functions.

Discriminants, Resultants and Multidimensional Determinants

I.M. Gelfand
M.M. Kapranov
A.V. Zelevinsky

Reprint of the 1994 Edition

Birkhäuser
Boston • Basel • Berlin

Israel M. Gelfand
Department of Mathematics
Rutgers University
New Brunswick, NJ 08903
U.S.A.

Mikhail M. Kapranov
Department of Mathematics
Yale University
New Haven, CT 06520-8283
U.S.A.

Andrei V. Zelevinsky
Department of Mathematics
Northeastern University
Boston, MA 02115
U.S.A.

Originally published in the series *Mathematics: Theory and Applications*

ISBN-13: 978-0-8176-4770-4

e-ISBN-13: 978-0-8176-4771-1

DOL: 10.1007/978-0-8176-4771-1

Library of Congress Control Number: 2007940668

Mathematics Subject Classification (2000): 14M25, 14M12, 13D25, 13XX, 14XX, 13C40, 15A15, 13F20, 52B20

©2008 Birkhäuser Boston

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Birkhäuser Boston, c/o Springer Science+Business Media LLC, 233 Spring Street, New York, NY 10013, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use in this publication of trade names, trademarks, service marks and similar terms, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.

Reprint from English language edition:

Discriminants, Resultants, and Multidimensional Determinants

by I. M. Gelfand, M.M. Kapranov, A. V. Zelevinsky

Copyright © 2008, Birkhäuser Boston

Birkhäuser Boston is a part of Springer Science+Business Media

All Rights Reserved

This reprint has been authorized by Springer Science & Business Media for distribution in China Mainland only and not for export therefrom.

Modern Birkhäuser Classics

Many of the original research and survey monographs in pure and applied mathematics published by Birkhäuser in recent decades have been groundbreaking and have come to be regarded as foundational to the subject. Through the MBC Series, a select number of these modern classics, entirely uncorrected, are being re-released in paperback (and as eBooks) to ensure that these treasures remain accessible to new generations of students, scholars, and researchers.

Contents

Preface	ix
-------------------	----

Introduction	1
------------------------	---

I. GENERAL DISCRIMINANTS AND RESULTANTS

CHAPTER 1. Projective Dual Varieties and General Discriminants

1. Definitions and basic examples	13
2. Duality for plane curves	16
3. The incidence variety and the proof of the biduality theorem	27
4. Further examples and properties of projective duality	30
5. The Katz dimension formula and its applications	39

CHAPTER 2. The Cayley Method for Studying Discriminants

1. Jet bundles and Koszul complexes	48
2. Discriminantal complexes	54
3. The degree and the dimension of the dual	61
4. Discriminantal complexes in terms of differential forms	71
5. The discriminant as the determinant of a spectral sequence	80

CHAPTER 3. Associated Varieties and General Resultants

1. Grassmannians. Preliminary material	91
2. Associated hypersurfaces	97
3. Mixed resultants	105
4. The Cayley method for the study of resultants	112

CHAPTER 4. Chow Varieties

1. Definitions and main properties	122
2. 0-cycles, factorizable forms and symmetric products	131
3. Cayley-Green-Morrison equations of Chow varieties	146

II. A-DISCRIMINANTS AND A-RESULTANTS

CHAPTER 5. Toric Varieties

1. Projectively embedded toric varieties	165
2. Affine toric varieties and semigroups	172
3. Local structure of toric varieties	177
4. Abstract toric varieties and fans	187

CHAPTER 6. Newton Polytopes and Chow Polytopes

1. Polynomials and their Newton polytopes	193
2. Theorems of Kouchnirenko and Bernstein on the number of solutions of a system of equations	200
3. Chow polytopes	206

CHAPTER 7. Triangulations and Secondary Polytopes

1. Triangulations and secondary polytopes	214
2. Faces of the secondary polytope	227
3. Examples of secondary polytopes	233

CHAPTER 8. A-Resultants and Chow Polytopes of Toric Varieties

1. Mixed $(A_1, \dots, A_k)$ -resultants	252
2. The A-resultant	255
3. The Chow polytope of a toric variety and the secondary polytope	259

CHAPTER 9. A-Discriminants

1. Basic definitions and examples	271
2. The discriminantal complex	275
3. A differential-geometric characterization of A-discriminantal hypersurfaces	285

CHAPTER 10. Principal A-Determinants

1. Statements of main results	297
2. Proof of the prime factorization theorem	313
3. Proof of the properties of generalized A-determinants	329
4. The proof of the product formula	333

CHAPTER 11. Regular A-Determinants and A-Discriminants

1. Differential forms on a singular toric variety and the regular A -determinant	344
2. Newton numbers and Newton functions	351
3. The Newton polytope of the regular A -determinant and D -equivalence of triangulations	361
4. More on D -equivalence	370
5. Relations to real algebraic geometry	378

III. CLASSICAL DISCRIMINANTS AND RESULTANTS**CHAPTER 12. Discriminants and Resultants for Polynomials in One Variable**

1. An overview of classical formulas and properties	397
2. Newton polytopes of the classical discriminant and resultant	411

CHAPTER 13. Discriminants and Resultants for Forms in Several Variables

1. Homogeneous forms in several variables	426
2. Forms in several groups of variables	437

CHAPTER 14. Hyperdeterminants

1. Basic properties of the hyperdeterminant	444
2. The Cayley method and the degree	450
3. Hyperdeterminant of the boundary format	458
4. Schläfli's method	475

APPENDIX A. Determinants of Complexes	480
--	------------

APPENDIX B. A. Cayley: On the Theory of Elimination	498
--	------------

Bibliography	503
-------------------------------	------------

Notes and References	513
---------------------------------------	------------

List of Notations	517
------------------------------------	------------

Index	521
------------------------	------------

Introduction

I

In this book we study discriminants and resultants of polynomials in several variables. The most familiar example is the discriminant of a quadratic polynomial $f(x) = ax^2 + bx + c$. This is

$$\Delta(f) = b^2 - 4ac, \quad (1)$$

which vanishes when $f(x)$ has a double root.

More generally, we can consider a polynomial $f(x_1, \dots, x_k)$ of degree $\leq d$ in k variables. An analog of a multiple root for f is a point where f vanishes together with all its first partial derivatives $\partial f / \partial x_i$. The *discriminant* $\Delta(f)$ is a polynomial function in the coefficients of f which vanishes whenever f has such a "multiple root." The existence of Δ is not quite trivial; however, it can be shown that $\Delta(f)$ exists and is unique up to sign if we require it to be irreducible and to have relatively prime integer coefficients. For instance, the discriminant of a cubic polynomial in one variable ($k = 1$, $d = 3$) is given by

$$\Delta(a_0 + a_1x + a_2x^2 + a_3x^3) = a_1^2a_2^2 - 4a_1^3a_3 - 4a_0a_2^3 - 27a_0^2a_3^2 + 18a_0a_1a_2a_3. \quad (2)$$

There is a subtle point in the definition of $\Delta(f)$: that is, $\Delta(f)$ depends not only on f but also on the choice of a degree bound d . For instance, the formula (2) applied to a quadratic polynomial gives a different expression from (1). With this in mind, we introduce the following more general version of a discriminant. Let A be a finite set of monomials in k variables, and let $\mathbf{C}^A$ denote the space of all polynomials with complex coefficients all of whose monomials belong to A . The A -discriminant $\Delta_A(f)$ is an irreducible polynomial in the coefficients of $f \in \mathbf{C}^A$ which vanishes whenever f has a multiple root $(x_1, \dots, x_k)$ with all $x_i \neq 0$ (the last condition is added to be able to ignore trivial multiple roots which can appear if all monomials from A have high degree). The A -discriminant will be one of our main objects of study.

The notion of the A -discriminant includes as special cases several fundamental algebraic concepts. If we take $A = \{1, x, \dots, x^m, y, yx, \dots, yx^n\}$, for example, then a typical polynomial from $\mathbf{C}^A$ has the form $f(x) + yg(x)$. Its A -discriminant is the *resultant* of f and g : it vanishes whenever f and g have a common root.

More generally, the resultant of $k + 1$ polynomials $f_0, \dots, f_k$ in k variables is defined as an irreducible polynomial in the coefficients of $f_0, \dots, f_k$, which

vanishes whenever these polynomials have a common root. The resultant can be treated as a special case of the A -discriminant of an auxiliary polynomial $f_0(x) + \sum_{i=1}^k y_i f_i(x)$, $x = (x_1, \dots, x_k)$.

Another important example occurs when A consists of n^2 monomials $x_i y_j$, $i, j = 1, \dots, n$. A typical polynomial from $\mathbb{C}^A$ is now a bilinear form $f(x, y) = \sum a_{ij} x_i y_j$ whose A -discriminant is the determinant of the matrix $\|a_{ij}\|$.

The last example has a natural generalization: we can take A as the set of all multilinear monomials in three or more groups of variables. An element $f \in \mathbb{C}^A$ (i.e., a multilinear form) is represented by a higher-dimensional "matrix" $\|a_{i_1 \dots i_r}\|$. Thus the A -discriminant Δ_A in this case is a polynomial function of a "matrix" which extends the notion of a determinant. Following Cayley [Ca1], we call this Δ_A the *hyperdeterminant* of $\|a_{i_1 \dots i_r}\|$. For example, the hyperdeterminant of a $2 \times 2 \times 2$ matrix $\|a_{ijk}\|$, $i, j, k = 0, 1$, is given by

$$\begin{aligned} & (a_{000}^2 a_{111}^2 + a_{001}^2 a_{110}^2 + a_{010}^2 a_{101}^2 + a_{011}^2 a_{100}^2) \\ & - 2(a_{000} a_{001} a_{110} a_{111} + a_{000} a_{010} a_{101} a_{111} + a_{000} a_{011} a_{100} a_{111} + a_{001} a_{010} a_{101} a_{110} \\ & + a_{001} a_{011} a_{110} a_{100} + a_{010} a_{011} a_{101} a_{100}) + 4(a_{000} a_{011} a_{101} a_{110} + a_{001} a_{010} a_{100} a_{111}). \end{aligned}$$

The study of hyperdeterminants was initiated by Cayley [Ca1] and Schläfli [Schl] but then was largely abandoned for 150 years. We present a treatment of hyperdeterminants in Chapter 14.

II

Let $\nabla_A = \{f \in \mathbb{C}^A : \Delta_A(f) = 0\}$ be the hypersurface in the space of polynomials consisting of polynomials with vanishing A -discriminant. We shall be mainly concerned with the following two closely related problems:

- (a) the study of the geometric properties of the hypersurface ∇_A ;
- (b) finding an explicit algebraic expression of the discriminant Δ_A .

To illustrate the importance of problem (a), consider the special case when A consists of all monomials in $x_1, \dots, x_k$ of a given degree d . Every $f \in \mathbb{C}^A$ (i.e., a homogeneous form of degree d) defines a hypersurface $\{f = 0\}$ in the projective space P^{k-1} . It is easy to see that ∇_A consists exactly of those f for which the hypersurface $\{f = 0\}$ is *singular*. Therefore the complement $\mathbb{C}^A - \nabla_A$ parametrizes all smooth hypersurfaces of a given degree in the projective space. To understand the geometric structure of $\mathbb{C}^A - \nabla_A$ is an important instance of the general moduli problem in algebraic geometry.

Equally important is the situation over the real numbers. Hilbert's 16th problem (classifying isotopy types of smooth real hypersurfaces of given degree d)

amounts to the study of connected components of $\mathbf{R}^A - \nabla_A$, the space of real polynomials with a non-vanishing discriminant.

Problem (b) has a long and glorious history. Explicit formulas for discriminants and resultants were the focus of several remarkable mathematicians in the last century. Many ingenious formulas were found by Cayley, Sylvester and their followers. However, we are still very far from a complete understanding of discriminants. For instance, an explicit polynomial expression for Δ_A is known only in a very limited number of special cases. Such formulas would be of great importance for the problem of finding explicit solutions of systems of polynomial equations. Problems of this kind are of interest not only for theoretical reasons, but are encountered more and more on a practical level because of the progress in computer technology.

III

We will use three main approaches in our study of discriminants and resultants:

- a geometric approach via projective duality and associated hypersurfaces;
- an algebraic approach via homological algebra and determinants of complexes (Whitehead torsion);
- a combinatorial approach via Newton polytopes and triangulations.

The geometric approach to discriminants is based on the observation that the discriminantal variety ∇_A is projectively dual to a certain variety X_A defined by a simple parametric representation. For example, if A consists of all monomials of degree d in k variables then X_A is the projective space P^{k-1} in its Veronese embedding. In the general case, X_A is the projective *toric* variety associated with A . The notion of the projectively dual variety $X^\vee$ makes sense for an arbitrary projective variety $X \subset P^{n-1}$: it is the closure of the set of all hyperplanes in P^{n-1} which are tangent to X at some smooth point. Thus the problem of finding the discriminant is a particular case of a more general geometric problem: find the equation(s) of $X^\vee$. We call this equation (in the case where $X^\vee$ is a hypersurface) the *X-discriminant*.

Although the resultants can be formally treated as discriminants of a special kind (see above), they have their own interesting geometric meaning. As for discriminants, we can associate the resultant to any projective variety $X \subset P^{n-1}$. Instead of $X^\vee$, we now consider the *associated hypersurface* $\mathcal{Z}(X)$ of X . If $\dim X = k - 1$ then $\mathcal{Z}(X)$ is the locus of all codimension k projective subspaces in P^{n-1} which meet X . The equation of $\mathcal{Z}(X)$ in the appropriate Grassmannian is the classical *Chow form* of X . This can be represented as a polynomial in the

coefficients of k linear forms defining a subspace from $\mathcal{Z}(X)$. We call this polynomial the X -*resultant* (the classical resultant of polynomials in several variables is a special case of this construction).

In Part I of this book we examine X -discriminants and X -resultants (or, in other words, projective duality and associated hypersurfaces) in the general context of projective geometry.

IV

The algebraic approach to discriminants and resultants which we use here goes back to Cayley. In his breathtaking 1848 note [Ca4] * he outlined a general method of writing down the resultant of several polynomials in several variables. We were very surprised to find that Cayley introduced in this note several fundamental concepts of homological algebra: complexes, exactness, Koszul complexes, and even the invariant now sometimes called the Whitehead torsion or Reidemeister-Franz torsion of an exact complex. The latter invariant is a natural generalization of the determinant of a square matrix (which itself was a rather recent discovery back in 1848!), so we prefer to call it the determinant of a complex. Using this terminology, Cayley's main result is that the resultant is the determinant of the Koszul complex.

Cayley's method is very general: without much effort it can be adapted to the study of X -discriminants and X -resultants associated as above to an arbitrary projective variety X . To get more detailed information, we complement Cayley's method with more recent tools such as coherent sheaves, perverse sheaves, microlocal geometry and D -modules.

V

Under a combinatorial approach we treat polynomials in the most naive way: as sums of monomials. To the best of our knowledge, there were no attempts in the classical literature to understand discriminants and resultants from this point of view, i.e., to describe which monomials can appear in them and with which coefficients. This is probably because the number of occurring monomials is usually very large. For example, the discriminant of a cubic form in three variables contains 2040 monomials (we are obliged to S. Duzhin who first showed it to us some years ago). At first glance, there seems to be no structure at all in these monomials and their coefficients. However, such a structure exists! The "magic crystal" that brings it to light is the concept of a *Newton polytope*.

Every monomial $x_1^{\omega_1} \cdots x_n^{\omega_n}$ in n variables can be visualized as a lattice point $(\omega_1, \dots, \omega_n)$ in $\mathbf{R}^n$. The Newton polytope $N(F)$ of a polynomial $F(x_1, \dots, x_n)$ is

* This note is reproduced as an appendix in this book

the convex hull in $\mathbf{R}^n$ of all lattice points representing monomials occurring in F . The structure of this polytope is deeply related to the geometry of the hypersurface $\{F = 0\}$. In fact, the asymptotic behavior of this hypersurface “at infinity” is controlled by the *extreme monomials* of F which correspond to the vertices of $N(F)$.

The notion of a Newton polytope goes back to Newton, and made some isolated appearances in the 19th century, cf. [Br 2]. More recently, some spectacular applications of Newton polytopes to classical algebraic problems (the number of solutions of systems of polynomial equations) have been found by A. Kouchnirenko, D. Bernstein, A. Khovansky [Ber], [Kou], [Kh]. We make use of these results in Part II.

It was a very surprising discovery for us when we realized that the Newton polytopes of A -discriminants admit a very nice combinatorial description. We recall that A is a finite set of monomials in k variables. As before, we represent the monomials from A as lattice points in $\mathbf{R}^k$. Hence we can consider the convex hull $Q \in \mathbf{R}^k$ of the set A . Our main result (which is the central point of Part II) is a description of the Newton polytope $N(\Delta_A)$ in terms of Q and A . Roughly speaking, it turns out that vertices of $N(\Delta_A)$ (i.e., extreme monomials in the A -discriminant) correspond to some *triangulations* of Q into simplices all of whose vertices lie in A .

The extreme monomial in Δ_A corresponding to a triangulation T of Q is determined explicitly once we know all the simplices in T and their volumes. The coefficient of this monomial is the product of numbers of the form $V_i^{V_i}$ where the V_i are the volumes of the simplices of T under suitable normalization. This provides an explanation of such coefficients as $4 = 2^2$ or $27 = 3^3$ in the formulas (1) and (2) above. The expression $\prod V_i^{V_i}$ (or, rather, its logarithm $\sum V_i \log(V_i)$) brings to mind the entropy of a probability distribution. It would be interesting to find a “probabilistic” reason for its appearance in discriminants. Even more intriguing is the fact that this appearance is not isolated—entropy-like expressions enter the formula for the rational uniformization of the variety ∇_A (see Chapter 9).

To illustrate the above description, consider the simplest A -discriminant of a quadratic polynomial $ax^2 + bx + c$ given by (1). Here A consists of $0, 1, 2 \in \mathbf{Z}$, the polytope Q is the segment $[0, 2]$, with its two “triangulations”. The first one consists of just one 1-dimensional “simplex” $[0, 2]$ of length 2, corresponding to the term $-4ac$ in (1). The second “triangulation” consists of two “simplices”: $[0, 1]$ and $[1, 2]$, corresponding to the term b^2 . Similarly, for the case of a cubic polynomial in one variable, we have $A = \{0, 1, 2, 3\}$ and $Q = [0, 3]$. There are now 4 triangulations of Q which correspond to the first four terms in the discriminant (2). Our final example is the determinant of a 2×2 matrix given by