

中国科学院中国孢子植物志编辑委员会 编辑

中 国 真 菌 志

第三十四卷


拟茎点霉属

戚佩坤 姜子德 向梅梅 主编

中国科学院知识创新工程重大项目

国家自然科学基金重大项目

(国家自然科学基金委员会 中国科学院 国家科学技术部 资助)



科 学 出 版 社

北 京

CONSILIO FLORARUM CRYPTOGRAMMARUM SINICARUM
ACADEMIAE SINICAE EDITA

FLORA FUNGORUM SINICORUM

VOL. 34

PHOMOPSIS

REDACTORES PRINCIPALES

Chi Peikun Jiang Zide Xiang Meimei

**A Major Project of the Knowledge Innovation Program
of the Chinese Academy of Sciences**

A Major Project of the National Natural Science Foundation of China

(Supported by the National Natural Science Foundation of China,
the Chinese Academy of Sciences, and the Ministry of Science and Technology of China)

Science Press
Beijing

拟茎点霉属

本 卷 著 者

戚佩坤 姜子德 习平根

(华南农业大学)

向梅梅

(仲恺农业技术学院)

PHOMOPSIS

AUCTORES

Chi Peikun Jiang Zide Xi Pinggen
(*Universitas Agriculturae Austro-Sinensis*)

Xiang Meimei
(*Collegium Agritechnicum Nomine Zhongkai*)

中国孢子植物志第四届编委名单

(1998年4月)

(右上角有*者为常委)

主 编 曾呈奎*

常务副主编 魏江春*

副 主 编 余永年* 吴鹏程* 毕列爵*

编 委 (以姓氏笔画为序)

王全喜 白金铠 田金秀* 刘 波 庄文颖*

庄剑云* 齐雨藻 齐祖同* 朱浩然 应建浙*

吴继农 邵力平 陈灼华 陈健斌* 陆保仁

林永水 郑柏林 郑儒永* 姜广正 赵震宇

施之新 胡人亮 胡征宇 胡鸿钧 高 谦

夏邦美 谢树莲 臧 穆 黎兴江

序

中国孢子植物志是非维管束孢子植物志，分《中国海藻志》、《中国淡水藻志》、《中国真菌志》、《中国地衣志》及《中国苔藓志》五部分。中国孢子植物志是在系统生物学原理与方法的指导下对中国孢子植物进行考察、收集和分类的研究成果；是生物多样性研究的主要内容；是物种保护的重要依据，对人类活动与环境甚至全球变化都有不可分割的联系。

中国孢子植物志是我国孢子植物物种数量、形态特征、生理生化性状、地理分布及其与人类关系等方面的综合信息库；是我国生物资源开发利用、科学研究与教学的重要参考文献。

我国气候条件复杂，山河纵横，湖泊星布，海域辽阔，陆生和水生孢子植物资源极其丰富。中国孢子植物分类工作的发展和《中国孢子植物志》的陆续出版，必将为我国开发利用孢子植物资源和促进学科发展发挥积极作用。

随着科学技术的进步，我国孢子植物分类工作在广度和深度方面将有更大的发展，对于这部著作也将不断补充、修订和提高。

中国科学院中国孢子植物志编辑委员会

1984年10月·北京

《中国真菌志》序

《中国真菌志》是在系统生物学原理和方法指导下，对中国真菌，即真菌界的子囊菌、担子菌、壶菌及接合菌四个门以及不属于真菌界的卵菌等三个门和黏菌及其类似的菌类生物进行搜集、考察和研究的成果。本志所谓“真菌”系广义概念，涵盖上述三大菌类生物（地衣型真菌除外），即当今所称“菌物”。

中国先民认识并利用真菌作为生活、生产资料，历史悠久，经验丰富，诸如酒、醋、酱、红曲、豆豉、豆腐乳、豆瓣酱等的酿制，蘑菇、木耳、茭白作食用，茯苓、虫草、灵芝等作药用，在制革、纺织、造纸工业中应用真菌进行发酵，以及利用具有抗癌作用和促进碳素循环的真菌，充分显示其经济价值和生态效益。此外，真菌又是多种植物和人畜病害的病原菌，危害甚大。因此，对真菌物种的形态特征、多样性、生理生化、亲缘关系、区系组成、地理分布、生态环境以及经济价值等进行研究和描述，非常必要。这是一项重要的基础科学研究，也是利用益菌、控制害菌、化害为利、变废为宝的应用科学的源泉和先导。

中国是具有悠久历史的文明古国，从远古到明代的 4500 年间，科学技术一直处于世界前沿，真菌学也不例外。酒是真菌的代谢产物，中国酒文化博大精深、源远流长，有六七千年历史。约在公元 300 年的晋代，江统在其《酒诰》诗中说：“酒之所兴，肇自上皇。或云仪狄，又曰杜康。有饭不尽，委之空桑。郁结成味，久蓄气芳。本出于此，不由奇方。”作者精辟地总结了我国酿酒历史和自然发酵方法，比之意大利学者雷蒂 (Radi, 1860) 提出微生物自然发酵法的学说约早 1500 年。在仰韶文化时期(5000~3000 B. C.)，我国先民已懂得采食蘑菇。中国历代古籍中均有食用菇蕈的记载，如宋代陈仁玉在其《菌谱》(1245 年)中记述浙江台州产鹅膏菌、松蕈等 11 种，并对其形态、生态、品级和食用方法等作了论述和分类，是中国第一部地方性食用蕈菌志。先民用真菌作药材也是一大创造，中国最早的药典《神农本草经》(成书于 102~200 A. D.) 所载 365 种药物中，有茯苓、雷丸、桑耳等 10 余种药用真菌的形态、色泽、性味和疗效的叙述。明代李时珍在《本草纲目》(1578) 中，记载“三菌”、“五蕈”、“六芝”、“七耳”以及羊肚菜、桑黄、鸡枞、雪蚕等 30 多种药用真菌。李氏将菌、蕈、芝、耳集为一类论述，在当时尚无显微镜帮助的情况下，其认识颇为精深。该籍的真菌学知识，足可代表中国古代真菌学水平，堪与同时代欧洲人（如 C. Clusius, 1529~1609）的水平比拟而无逊色。

15 世纪以后，居世界领先地位的中国科学技术，逐渐落后。从 18 世纪中叶到 20 世纪 40 年代，外国传教士、旅行家、科学工作者、外交官、军官、教师以及负有特殊任务者，纷纷来华考察，搜集资料，采集标本，研究鉴定，发表论文或专辑。如法国传教士西博特 (P. M. Cibot) 1759 年首先来到中国，一住就是 25 年，对中国的植物（含真菌）写过不少文章，1775 年他发表的五棱散尾菌 (*Lysurus mokusin*)，是用现代科学方法研究发表的第一个中国真菌。继而，俄国的波塔宁 (G. N. Potanin, 1876)、意大利的吉拉迪 (P. Giraldui, 1890)、奥地利的汉德尔-马泽蒂 (H. Handel-Mazzetti,

1913)、美国的梅里尔 (E. D. Merrill, 1916)、瑞典的史密斯 (H. Smith, 1921) 等共 27 人次来我国采集标本。研究发表中国真菌论著 114 篇册, 作者多达 60 余人次, 报道中国真菌 2040 种, 其中含 10 新属、361 新种。东邻日本自 1894 年以来, 特别是 1937 年以后, 大批人员涌到中国, 调查真菌资源及植物病害, 采集标本, 鉴定发表。据初步统计, 发表论著 172 篇册, 作者 67 人次以上, 共报道中国真菌约 6000 种 (有重复), 其中含 17 新属、1130 新种。其代表人物在华北有三宅市郎 (1908), 东北有三浦道哉 (1918), 台湾有泽田兼吉 (1912); 此外, 还有斋藤贤道、伊藤诚哉、平冢直秀、山本和太郎、逸见武雄等数十人。

国人用现代科学方法研究中国真菌始于 20 世纪初, 最初工作多侧重于植物病害和工业发酵, 纯真菌学研究较少。在一二十年代便有不少研究报告和学术论文发表在中外各种刊物上, 如胡先骕 1915 年的“菌类鉴别法”, 章祖纯 1916 年的“北京附近发生最盛之植物病害调查表”以及钱穉孙 (1918)、邹钟琳 (1919)、戴芳澜 (1920)、李寅恭 (1921)、朱凤美 (1924)、孙豫寿 (1925)、俞大绂 (1926)、魏岳寿 (1928) 等的论文。三四十年代有陈鸿康、邓叔群、魏景超、凌立、周宗璜、欧世璜、方心芳、王云章、裘维蕃等发表的论文, 为数甚多。他们中有的人终生或大半生都从事中国真菌学的科教工作, 如戴芳澜 (1893~1973) 著“江苏真菌名录” (1927)、“中国真菌杂记” (1932~1946)、《中国已知真菌名录》 (1936, 1937)、《中国真菌总汇》 (1979) 和《真菌的形态和分类》 (1987) 等, 他发表的“三角枫上白粉菌一新种” (1930), 是国人用现代科学方法研究、发表的第一个中国真菌新种。邓叔群 (1902~1970) 著“南京真菌记载” (1932~1933)、“中国真菌续志” (1936~1938)、《中国高等真菌志》 (1939) 和《中国的真菌》 (1963, 1996) 等, 堪称《中国真菌志》的先导。上述学者以及其他许多真菌学工作者, 为《中国真菌志》研编的起步奠定了基础。

在 20 世纪后半叶, 特别是改革开放以来的 20 多年, 中国真菌学有了迅猛的发展, 如各类真菌学课程的开设, 各级学位研究生的招收和培养, 专业机构和学会的建立, 专业刊物的创办和出版, 地区真菌志的问世等, 使真菌学人才辈出, 为《中国真菌志》的研编输送了新鲜血液。1973 年中国科学院广州“三志”会议决定, 《中国真菌志》的研编正式启动, 1987 年由郑儒永、余永年等编辑出版了《中国真菌志》第 1 卷《白粉菌目》, 至 2000 年已出版 14 卷。自第 2 卷开始实行主编负责制, 2. 《银耳目和花耳目》 (刘波主编, 1992); 3. 《多孔菌科》 (赵继鼎, 1998); 4. 《小煤炱目 I》 (胡炎兴, 1996); 5. 《曲霉属及其相关有性型》 (齐祖同, 1997); 6. 《霜霉目》 (余永年, 1998); 7. 《层腹菌目》 (刘波, 1998); 8. 《核盘菌科和地舌菌科》 (庄文颖, 1998); 9. 《假尾孢属》 (刘锡琰、郭英兰, 1998); 10. 《锈菌目 I》 (王云章、庄剑云, 1998); 11. 《小煤炱目 II》 (胡炎兴, 1999); 12. 《黑粉菌科》 (郭林, 2000); 13. 《虫霉目》 (李增智, 2000); 14. 《灵芝科》 (赵继鼎、张小青, 2000)。盛世出巨著, 在国家“科教兴国”英明政策的指引下, 《中国真菌志》的研编和出版, 定将为中华灿烂文化做出新贡献。

余永年 谨识
庄文颖

中国科学院微生物研究所
中国·北京·中关村
公元 2002 年 09 月 15 日

Foreword of Flora Fungorum Sinicorum

Flora Fungorum Sinicorum summarizes the achievements of Chinese mycologists based on principles and methods of systematic biology in intensive studies on the organisms studied by mycologists, which include non-lichenized fungi of the Kingdom Fungi, some organisms of the Chromista, such as oomycetes etc., and some of the Protozoa, such as slime molds. In this series of volumes, results from extensive collections, field investigations, and taxonomic treatments reveal the fungal diversity of China.

Our Chinese ancestors were very experienced in the application of fungi in their daily life and production. Fungi have long been used in China as food, such as edible mushrooms, including jelly fungi, and the hypertrophic stems of water bamboo infected with *Ustilago esculenta*; as medicines, like *Cordyceps sinensis* (caterpillar fungus), *Poria cocos* (China root), and *Ganoderma* spp. (lingzhi); and in the fermentation industry, for example, manufacturing liquors, vinegar, soy-sauce, *Monascus*, fermented soya beans, fermented bean curd, and thick broad-bean sauce. Fungal fermentation is also applied in the tannery, papermaking, and textile industries. The anti-cancer compounds produced by fungi and functions of saprophytic fungi in accelerating the carbon-cycle in nature are of economic value and ecological benefits to human beings. On the other hand, fungal pathogens of plants, animals and human cause a huge amount of damage each year. In order to utilize the beneficial fungi and to control the harmful ones, to turn the harmfulness into advantage, and to convert wastes into valuables, it is necessary to understand the morphology, diversity, physiology, biochemistry, relationship, geographical distribution, ecological environment, and economic value of different groups of fungi. *Flora Fungorum Sinicorum* plays an important role from precursor to fountainhead for the applied sciences.

China is a country with an ancient civilization of long standing. In the 4500 years from remote antiquity to the Ming Dynasty, her science and technology as well as knowledge of fungi stood in the leading position of the world. Wine is a metabolite of fungi. The Wine Culture history in China goes back 6000 to 7000 years ago, which has a distant source and a long stream of extensive knowledge and profound scholarship. In the Jin Dynasty (ca. 300 A. D.), JIANG Tong, the famous writer, gave a vivid account of the Chinese fermentation history and methods of wine processing in one of his poems entitled *Drinking Games* (Jiu Gao), 1500 years earlier than the theory of microbial fermentation in natural conditions raised by the Italian scholar, Radi (1860). During the period of the Yangshao Culture (5000—3000 B. C.), our Chinese ancestors knew how to eat mushrooms. There were a great number of records of edible mushrooms in Chinese ancient books. For example, back to

the Song Dynasty, CHEN Ren-Yu (1245) published the *Mushroom Menu* (Jun Pu) in which he listed 11 species of edible fungi including *Amanita* sp. and *Tricholoma matsutake* from Taizhou, Zhejiang Province, and described in detail their morphology, habitats, taxonomy, taste, and way of cooking. This was the first local flora of the Chinese edible mushrooms. Fungi used as medicines originated in ancient China. The earliest Chinese pharmacopoeia, *Shen-Nong Materia Medica* (Shen Nong Ben Cao Jing), was published in 102—200 A. D. Among the 365 medicines recorded, more than 10 fungi, such as *Poria cocos* and *Polyporus mylittae*, were included. Their fruitbody shape, color, taste, and medical functions were provided. The great pharmacist of Ming Dynasty, LI Shi-Zhen (1578) published his eminent work *Compendium Materia Medica* (Ben Cao Gang Mu) in which more than thirty fungal species were accepted as medicines, including *Aecidium mori*, *Cordyceps sinensis*, *Morchella* spp., *Termitomyces* sp., etc. Before the invention of microscope, he managed to bring fungi of different classes together, which demonstrated his intelligence and profound knowledge of biology.

After the 15th century, development of science and technology in China slowed down. From middle of the 18th century to the 1940's, foreign missionaries, tourists, scientists, diplomats, officers, and other professional workers visited China. They collected specimens of plants and fungi, carried out taxonomic studies, and published papers, exsiccatae, and monographs based on Chinese materials. The French missionary, P. M. Cibot, came to China in 1759 and stayed for 25 years to investigate plants including fungi in different regions of China. Many papers were written by him. *Lysurus mokusin*, identified with modern techniques and published in 1775, was probably the first Chinese fungal record by these visitors. Subsequently, around 27 man-times of foreigners attended field excursions in China, such as G. N. Potanin from Russia in 1876, P. Giraldui from Italy in 1890, H. Handel-Mazzetti from Austria in 1913, E. D. Merrill from the United States in 1916, and H. Smith from Sweden in 1921. Based on examinations of the Chinese collections obtained, 2040 species including 10 new genera and 361 new species were reported or described in 114 papers and books. Since 1894, especially after 1937, many Japanese entered China. They investigated the fungal resources and plant diseases, collected specimens, and published their identification results. According to incomplete information, some 6000 fungal names (with synonyms) including 17 new genera and 1130 new species appeared in 172 publications. The main workers were I. Miyake in the Northern China, M. Miura in the Northeast, K. Sawada in Taiwan, as well as K. Saito, S. Ito, N. Hiratsuka, W. Yamamoto, T. Hemmi, etc.

Research by Chinese mycologists started at the turn of the 20th century when plant diseases and fungal fermentation were emphasized with very little systematic work. Scientific papers or experimental reports were published in domestic and international journals during the 1910's to 1920's. The best-known are "Identification of the fungi" by H. H. Hu in 1915, "Plant disease report from Peking and the adjacent regions" by C. S. Chang in 1916,

and papers by S. S. Chian (1918), C. L. Chou (1919), F. L. Tai (1920), Y. G. Li (1921), V. M. Chu (1924), Y. S. Sun (1925), T. F. Yu (1926), and N. S. Wei (1928). Mycologists who were active at the 1930's to 1940's are H. K. Chen, S. C. Teng, C. T. Wei, L. Ling, C. H. Chow, S. H. Ou, S. F. Fang, Y. C. Wang, W. F. Chiu, and others. Some of them dedicated their lifetime to research and teaching in mycology. Prof. F. L. Tai (1893—1973) is one of them, whose representative works were “List of fungi from Jiangsu” (1927), “Notes on Chinese fungi” (1932—1946), *A List of Fungi Hitherto Known from China* (1936, 1937), *Sylloge Fungorum Sinicorum* (1979), *Morphology and Taxonomy of the Fungi* (1987), etc. His paper entitled “A new species of *Uncinula* on *Acer trifidum* Hook. & Arn.” was the first new species described by a Chinese mycologist. Prof. S. C. Teng (1902—1970) is also an eminent teacher. He published “Notes on fungi from Nanking” in 1932—1933, “Notes on Chinese fungi” in 1936—1938, *A Contribution to Our Knowledge of the Higher Fungi of China* in 1939, and *Fungi of China* in 1963 and 1996. Work done by the above-mentioned scholars lays a foundation for our current project on *Flora Fungorum Sinicorum*.

In 1973, an important meeting organized by the Chinese Academy of Sciences was held in Guangzhou (Canton) and a decision was made, uniting the related scientists from all over China to initiate the long term project “Fauna, Flora, and Cryptogamic Flora of China”. Work on *Flora Fungorum Sinicorum* thus started. Significant progress has been made in development of Chinese mycology since 1978. Many mycological institutions were founded in different areas of the country. The Mycological Society of China was established, the journals *Acta Mycological Sinica* and *Mycosystema* were published as well as local floras of the economically important fungi. A young generation in field of mycology grew up through post-graduate training programs in the graduate schools. The first volume of Chinese Mycoflora on the Erysiphales (edited by R. Y. Zheng & Y. N. Yu, 1987) appeared. Up to now, 14 volumes have been published: Tremellales and Dacrymycetales edited by B. Liu (1992), Polyporaceae by J. D. Zhao (1998), Meliolales Part I (Y. X. Hu, 1996), *Aspergillus* and its related teleomorphs (Z. T. Qi, 1997), Peronosporales (Y. N. Yu, 1998), Sclerotiniaceae and Geoglossaceae (W. Y. Zhuang, 1998), *Pseudocercospora* (X. J. Liu & Y. L. Guo, 1998), Uredinales Part I (Y. C. Wang & J. Y. Zhuang, 1998), Meliolales Part II (Y. X. Hu, 1999), Ustilaginaceae (L. Guo, 2000), Entomophthorales (Z. Z. Li, 2000), and Ganodermataceae (J. D. Zhao & X. Q. Zhang, 2000). We eagerly await the coming volumes and expect the completion of *Flora Fungorum Sinicorum* which will reflect the flourishing of Chinese culture.

Y. N. Yu and W. Y. Zhuang
Institute of Microbiology, CAS, Beijing
September 15, 2002

致 谢

本卷册的研究及撰写工作是在中国科学院中国孢子植物志编辑委员会和华南农业大学的领导和支持下完成的。

感谢 1983 年以来,攻读真菌分类学的研究生林石明、曹福祥、张传飞、黄亚军、陈永青、郑晓慧、曾莉、罗利娟、常长青、程颖慧、马莉、李敏慧等在标本采集、分布调查、拟茎点霉鉴定和研究等方面所做的大量工作和仲恺农业技术学院黄江华老师所付出的辛勤劳动。

感谢中国科学院郑儒永院士多年来的支持与鼓励,并提供了重要的有关分类资料;感谢中国科学院姚一建研究员在分类资料方面所给予的帮助;感谢华南农业大学李秉滔教授对拉丁文稿的审阅,以及吴万春教授、肖绵韵高级实验师在植物分类方面给予的指导。

中国科学院微生物研究所菌物标本馆借调了馆藏的拟茎点霉属标本。在标本采集过程中,还得到了新疆农业大学赵震宇教授,云南农业大学张中义、朱有勇、刘云龙教授和胡先奇博士,云南大学张克勤教授,沈阳农业大学吕国忠教授,中南林学院曹福祥教授,西昌学院郑晓慧副教授,云南省农科院曾莉副研究员,贵州大学向红琼教授,贵州药用植物园王兰主任、于子文副主任,贵阳植物园魏德生老师的帮助,谨致以衷心感谢!

说 明

1. 本书是我国拟茎点霉属的研究总结。全书包括绪论、专论、附录、参考文献和索引五部分。
2. 绪论部分简要叙述了拟茎点霉属的经济意义、形态、研究史、属级特征、与邻近属的区别、分类标准及种的亲缘关系等。
3. 专论部分描述了我国寄生在 74 科植物上的拟茎点霉属真菌 133 个种，包括 2 个专化型。描述按寄主科进行，科名及科内真菌学名都按字母顺序排列。寄主科内有 3 个种以上者均设有分种检索表。每个种包括正名、异名、详细的形态描述、按学名字母顺序排列的寄主名称及在国内的分布和世界分布，并附有显微绘图。讨论部分包括种的历史渊源和与邻近种的区别。
4. 附录包括各科、属、种寄主上的拟茎点霉名录。
5. 参考文献按作者姓名字母顺序排列。我国的作者按汉语拼音字母顺序排列。文献按发表时的语种引用。
6. 索引包括：寄主汉名索引、真菌汉名索引、寄主学名索引、真菌学名索引。寄主汉名索引和真菌汉名索引均按汉语拼音字母顺序排列。
7. 真菌汉名主要根据 1990 年出版的《真菌名词及名称》，寄主汉名主要根据 1996 年航空工业出版社出版的《新编拉汉英植物名称》，1979 年的《英拉汉植物名称》及地方植物志。菌名命名人缩写主要根据《菌物名称的作者》(Kirk et Asell 1992)。
8. 专论部分每个种的形态描述及数据均系根据我国的标本研究和测量所得，种下引证的标本全由作者直接研究。产地后注明的标本室名称 SCHM 为华南农业大学真菌标本室。
9. 本书共有插图 135 幅，全部为作者根据标本所作的显微绘图。
10. 国内分布以我国直辖市及各行政省、自治区的市、县或区为单位。直辖市、省或自治区之间以“;”号区分，同一省或自治区下面的不同市、县、区之间则以“,”号区分。
11. 世界分布根据文献资料整理而成，各国的名称按字母顺序排列。

目 录

序

中国孢子植物志总序

《中国真菌志》序

致谢

说明

绪论	(1)
属的研究史	(1)
寄生性、致病性及专化性	(2)
所致症状	(3)
属级特征	(4)
培养性状	(5)
与拟茎点霉易混淆的属	(7)
分类标准	(7)
种的亲缘关系	(15)
有性态	(15)
分子生物学	(16)
专论	(19)
裸子植物上的拟茎点霉	(19)
柏科 (Cupressaceae)	(19)
隐藏拟茎点霉 <i>Phomopsis occulta</i> (Sacc.) Traverso	(19)
银杏科 (Ginkgoaceae)	(20)
银杏拟茎点霉 <i>Phomopsis ginkginis</i> Z. D. Jiang, C. Q. Chang et M. M. Xiang	(20)
罗汉松科 (Podocarpaceae)	(21)
竹柏拟茎点霉 <i>Phomopsis podocarpi</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(21)
被子植物上的拟茎点霉	(22)
槭树科 (Aceraceae)	(22)
欧亚槭拟茎点霉 <i>Phomopsis platanoidis</i> (Cooke) Died.	(22)
龙舌兰科 (Agavaceae)	(23)
龙血树拟茎点霉 <i>Phomopsis dracaenae</i> Sahní	(23)
龙血树生拟茎点霉 <i>Phomopsis dracaenicola</i> Z. D. Jiang, P. G. Xi et P. K. Chi	(24)
丝兰拟茎点霉 <i>Phomopsis yuccae</i> (Cooke) Grove	(25)
苋科 (Amaranthaceae)	(26)

千日红拟茎点霉 <i>Phomopsis gomphrenae</i> M. H. Li, Z. D. Jiang et P. K. Chi	(26)
漆树科 (Anacardiaceae)	(27)
芒果拟茎点霉 <i>Phomopsis mangiferae</i> Ahmad	(27)
漆树拟茎点霉 <i>Phomopsis rhois</i> (Sacc.) Traverso	(29)
番荔枝科 (Annonaceae)	(30)
番荔枝叶生拟茎点霉 <i>Phomopsis anonacearum</i> Bond.-Mont.	(30)
夹竹桃科 (Apocynaceae)	(31)
黄夹竹桃拟茎点霉 <i>Phomopsis thevetiae</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	(31)
冬青科 (Aquifoliaceae)	(32)
外皮拟茎点霉小专化型 <i>Phomopsis crustosa</i> (Sacc., Bommer et Roussel) Traverso	
f. <i>minuscule</i> Dos Santos et Camara	(32)
天南星科 (Araceae)	(34)
花烛拟茎点霉 <i>Phomopsis anthurii</i> X. H. Zheng, M. M. Xiang et P. K. Chi	(34)
喜林芋拟茎点霉 <i>Phomopsis philodendri</i> Z. D. Jiang, X. H. Zheng et P. K. Chi	(34)
五加科 (Araliaceae)	(35)
黑白拟茎点霉 <i>Phomopsis melaleuca</i> (Berk. et M. A. Curtis) Grove	(35)
鹅掌柴拟茎点霉 <i>Phomopsis schefflerae</i> P. G. Xi, P. K. Chi et Z. D. Jiang	(36)
凤仙花科 (Balsaminaceae)	(37)
凤仙拟茎点霉 <i>Phomopsis impatientis</i> Dearn. et House	(37)
小檗科 (Berberidaceae)	(38)
十大功劳拟茎点霉 <i>Phomopsis mahoniae</i> Grove	(38)
木棉科 (Bombacaceae)	(39)
猴面包树拟茎点霉 <i>Phomopsis adansoniae</i> L. Ma, P. K. Chi et Z. D. Jiang	(40)
木棉拟茎点霉 <i>Phomopsis bombacis</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	(41)
榴莲拟茎点霉 <i>Phomopsis durionis</i> H. Syd.	(42)
大果拟茎点霉 <i>Phomopsis macrocarpae</i> P. G. Xi, Z. D. Jiang et P. K. Chi	(43)
凤梨科 (Bromeliaceae)	(44)
类色二孢拟茎点霉 <i>Phomopsis diplodinoides</i> (Sacc.) Punith.	(44)
筒凤花拟茎点霉 <i>Phomopsis spectabilis</i> V. G. Rao et B. R. D. Yadav	(45)
黄杨科 (Buxaceae)	(47)
黄杨生拟茎点霉 <i>Phomopsis buxicola</i> P. K. Chi, C. Q. Chang et Z. D. Jiang	(47)
苏木科 (Caesalpiniaceae)	(48)
羊蹄甲拟茎点霉 <i>Phomopsis bauhiniae</i> Alcalde	(48)
羊蹄甲生拟茎点霉 <i>Phomopsis bauhiniicola</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	
Chi	(50)
决明拟茎点霉 <i>Phomopsis cassiae</i> Camara	(51)
喜决明拟茎点霉 <i>Phomopsis cassiophila</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	(52)
肥皂荚拟茎点霉 <i>Phomopsis gymnocladi</i> Byzova	(53)

腊梅科 (Calycanthaceae)	(54)
腊梅拟茎点霉 <i>Phomopsis chimonanthi</i> C. Q. Chang, M. M. Xiang et P. K. Chi	(54)
忍冬科 (Caprifoliaceae)	(54)
荚迷拟茎点霉 <i>Phomopsis tineae</i> (Sacc.) Died.	(54)
番木瓜科 (Caricaceae)	(55)
番木瓜拟茎点霉 <i>Phomopsis caricae-papayae</i> Petrak et Cif.	(55)
菊科 (Compositae)	(56)
旋覆花拟茎点霉 <i>Phomopsis inulina</i> (Sacc.) Petrak	(56)
埋生拟茎点霉 <i>Phomopsis immersa</i> (Sacc.) Höhn.	(58)
山茱萸科 (Cornaceae)	(59)
桃叶珊瑚拟茎点霉 <i>Phomopsis aucubae</i> (Westend.) Traverso	(59)
五桠果科 (Dilleniaceae)	(60)
五桠果拟茎点霉 <i>Phomopsis dilleniae</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	(60)
薯蓣科 (Dioscoreaceae)	(61)
山药拟茎点霉 <i>Phomopsis dioscoreae</i> Sacc.	(61)
柿科 (Ebenaceae)	(62)
柿拟茎点霉 <i>Phomopsis diospyri</i> (Sacc.) Traverso et Spessa	(62)
胡颓子科 (Elaeagnaceae)	(63)
胡颓子生拟茎点霉 <i>Phomopsis elaeagnicola</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(63)
杜鹃花科 (Ericaceae)	(64)
杜鹃拟茎点霉 <i>Phomopsis rhododendri</i> Z. D. Jiang, M. M. Xiang et P. K. Chi	(64)
杜鹃生拟茎点霉 <i>Phomopsis rhododendricola</i> L. Ma, M. M. Xiang et P. K. Chi	(65)
杜仲科 (Eucommiaceae)	(66)
杜仲拟茎点霉 <i>Phomopsis eucommiae</i> F. X. Cao et P. K. Chi	(66)
杜仲生拟茎点霉 <i>Phomopsis eucommiicola</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(67)
大戟科 (Euphorbiaceae)	(68)
土密树拟茎点霉 <i>Phomopsis brideliae</i> Chowdhury	(68)
大戟拟茎点霉 <i>Phomopsis euphorbiae</i> (Sacc.) Sacc.	(69)
橡胶生拟茎点霉 <i>Phomopsis heveicola</i> L. Ma, M. M. Xiang et P. K. Chi	(70)
叶下珠拟茎点霉 <i>Phomopsis phyllanthi</i> Punith.	(71)
叶下珠生拟茎点霉 <i>Phomopsis phyllanthicola</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(71)
壳斗科 (Fagaceae)	(72)
栗拟茎点霉 <i>Phomopsis castanea</i> (Sacc.) Petrak	(72)
夏栎生拟茎点霉 <i>Phomopsis quercicola</i> Moesz	(73)
大枫子科 (Flacourtiaceae)	(74)
大枫子拟茎点霉 <i>Phomopsis hydnocarpi</i> S. M. Lin et P. K. Chi	(74)
金缕梅科 (Hamamelidaceae)	(75)

枫香拟茎点霉 <i>Phomopsis liquidambaris</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(75)
红继木拟茎点霉 <i>Phomopsis loropetali</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(76)
胡桃科 (Juglandaceae)	(77)
胡桃楸拟茎点霉 <i>Phomopsis juglandina</i> (Sacc.) Traverso	(77)
铅色拟茎点霉 <i>Phomopsis lizivia</i> Höhn.	(78)
枫杨生拟茎点霉 <i>Phomopsis pterocaryicola</i> L. Ma, Z. D. Jiang et P. K. Chi	(79)
樟科 (Lauraceae)	(80)
樟拟茎点霉 <i>Phomopsis cinnamomi</i> S. M. Lin et P. K. Chi	(80)
樟生拟茎点霉 <i>Phomopsis cinnamomicola</i> Z. D. Jiang, M. M. Xiang et P. K. Chi ...	(81)
豆科 (Leguminosae)	(82)
菜豆拟茎点霉 <i>Phomopsis phaseoli</i> (Desm.) Sacc.	(82)
百合科 (Liliaceae)	(83)
芦笋拟茎点霉 <i>Phomopsis asparagi</i> (Sacc.) Bubák	(83)
千屈菜科 (Lythraceae)	(84)
紫薇拟茎点霉 <i>Phomopsis lagerstroemiae</i> C. Q. Chang, M. M. Xiang et P. K. Chi	(84)
木兰科 (Magnoliaceae)	(85)
鹅掌楸拟茎点霉 <i>Phomopsis liriodendri</i> Grove	(85)
木兰拟茎点霉 <i>Phomopsis magnoliae</i> M. M. Xiang, Z. D. Jiang et P. K. Chi	(87)
木兰生拟茎点霉 <i>Phomopsis magnoliicola</i> (Syd. et P. Syd.) Died.	(88)
似木兰拟茎点霉 <i>Phomopsis magnolina</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(88)
似含笑拟茎点霉 <i>Phomopsis michelina</i> C. Q. Chang, Z. D. Jiang et P. K. Chi	(89)
含笑生拟茎点霉 <i>Phomopsis micheliicola</i> Z. D. Jiang, P. K. Chi et M. M. Xiang ...	(90)
锦葵科 (Malvaceae)	(90)
木槿拟茎点霉 <i>Phomopsis hibisci</i> Dearn.	(90)
竹芋科 (Marantaceae)	(91)
竹芋拟茎点霉 <i>Phomopsis marantacearum</i> L. Zeng, Z. D. Jiang et P. K. Chi	(91)
楝科 (Meliaceae)	(92)
楝拟茎点霉 <i>Phomopsis abdita</i> (Sacc.) Traverso	(92)
坚木拟茎点霉 <i>Phomopsis dysorxylis</i> (Sacc.) Petrak	(93)
含羞草科 (Mimosaceae)	(95)
金合欢生拟茎点霉 <i>Phomopsis acaciicola</i> M. M. Xiang, Z. D. Jiang et P. K. Chi ...	(95)
多疣拟茎点霉 <i>Phomopsis mendax</i> (Sacc.) Traverso	(95)
桑科 (Moraceae)	(96)
灰色拟茎点霉 <i>Phomopsis cinerascens</i> (Sacc.) Bubák	(96)
胶榕拟茎点霉 <i>Phomopsis elastica</i> Petrak	(97)
榕拟茎点霉 <i>Phomopsis ficus</i> (Castagne) Traverso et Spessa	(98)
芭蕉科 (Musaceae)	(99)
香蕉生拟茎点霉 <i>Phomopsis musicola</i> (Stevens et Young) Punith.	(99)