


Yi-Bing Shen · Zhongmin Shen

# Introduction to Modern Finsler Geometry

现代芬斯勒几何初步



高等教育出版社

# Introduction to Modern Finsler Geometry

现代芬斯勒几何初步

Yi-Bing Shen

Zhejiang University, China

Zhongmin Shen

Indiana University-Purdue University Indianapolis, USA

高等教育出版社·北京

**Yi-Bing Shen**  
Zhejiang University  
China  
Email: yibingshen@zju.edu.cn

**Zhongmin Shen**  
Indiana University-Purdue University Indianapolis  
USA  
Email: zshen@iupui.edu

### 图书在版编目(CIP)数据

现代芬斯勒几何初步=Introduction to Modern Finsler  
Geometry: 英文 / 沈一兵, 沈忠民编著. —北京: 高等教育  
出版社, 2016.1

ISBN 978-7-04-044424-7

I. ①现… II. ①沈… ②沈… III. ①芬斯拉空间—  
英文 IV. ①O186.14

中国版本图书馆CIP数据核字(2015)第295718号

策划编辑 赵天夫      责任编辑 赵天夫      封面设计 张楠  
版式设计 马云      责任印制 韩刚

---

出版发行	高等教育出版社	网 址	<a href="http://www.hep.edu.cn">http://www.hep.edu.cn</a>
社 址	北京市西城区德外大街4号		<a href="http://www.hep.com.cn">http://www.hep.com.cn</a>
邮政编码	100120	网上订购	<a href="http://www.hepmall.com.cn">http://www.hepmall.com.cn</a>
印 刷	涿州市星河印刷有限公司		<a href="http://www.hepmall.com">http://www.hepmall.com</a>
开 本	787×960 1/16		<a href="http://www.hepmall.cn">http://www.hepmall.cn</a>
印 张	25.5	版 次	2016年1月第1版
字 数	530千字	印 次	2016年1月第1次印刷
购书热线	010-58581118	定 价	98.00元
咨询电话	400-810-0598		

---

本书如有缺页、倒页、脱页等质量问题, 请到所购图书销售部门联系调换  
版权所有 侵权必究  
物料号 44424-00

Introduction to  
Modern Finsler  
Geometry

现代芬斯勒几何初步

# Preface

“Finsler geometry is just Riemannian geometry without quadratic restriction”, commented by the past famous geometer, S. S. Chern. In fact, early in 1854, B. Riemann had introduced the concept of Finsler geometry in his ground-breaking Habilitationsvortrag. He had seen the difference between metrics of quadratic type (i.e. Riemannian metrics) and those in the general case. No essential development was made until 1918 when P. Finsler studied the geometry of curves and surfaces in the general case. Therefore, more precisely, we should call this subject Riemann-Finsler geometry.

Since more than twenty years ago, substantial progress has been made in Finsler geometry, especially in global Finsler geometry, so that we have seen a completely new outlook. Informally speaking, Riemannian geometry studies spaces with only black and white colors, while Finsler geometry studies a colorful world. The methods and ideas used in Finsler geometry not only are closely related to other mathematical branches such as differential equations, Lie groups, algebra, topology, nonlinear analysis, etc., but also have more and more applications to mathematical physics, theoretical physics, mathematical biology, control theory, informatics, etc. Therefore, not just in theory but also in application, Finsler geometry has shown its strong vitality and great value.

In order to meet the need of education for senior undergraduate and graduate students, under the influence of books [10, 103], we wrote this textbook, based on many years of teaching experience. The whole book is divided into 11 chapters: in the first five chapters, we discuss differential manifolds, Finsler metrics, the Chern connection, Riemannian and non-Riemannian quantities. The rest is written for further studies. This second part covers projective transformations, comparison theorems, fundamental group, minimal immersions, harmonic maps, Einstein metrics, conformal

transformations and conformal vector fields, the Finsler Laplacian and its first eigenvalue, etc. At the end of every chapter there are some exercises, which are important complements for the contents. The final Appendix is to provide Maple programs on the computations of geometric quantities in Finsler geometry.

With the main tool of tensor analysis, we systematically introduce the basic concepts and methods in Finsler geometry, and we do our best to include the classical theory as well as the newest developments, so that readers can do research independently after studying this book. This book may be used as a selective textbook for senior undergraduate students and a regular textbook for graduate students. It can also be used as a reference book for mathematical physics, theoretical physics, control theory, etc. We believe that this book is of positive significance as an addition and improvement to current textbooks in colleges.

We would like to take this opportunity to thank the National Natural Science Foundation of China (No. 11171297), the Center of Mathematical Sciences and the Department of Mathematics at Zhejiang University, and our many students who contributed to this book.

In conclusion we acknowledge the Higher Education Press in China and World Scientific Publishing Company in Singapore for their great support.

Yi-Bing Shen  
Zhongmin Shen  
March 2015

# Contents

<i>Preface</i>	v
<b>Foundations</b>	<b>1</b>
1. Differentiable Manifolds	3
1.1 Differentiable manifolds . . . . .	3
1.1.1 Differentiable manifolds . . . . .	3
1.1.2 Examples of differentiable manifolds . . . . .	3
1.2 Vector fields and tensor fields . . . . .	6
1.2.1 Vector bundles . . . . .	6
1.2.2 Tensor fields . . . . .	7
1.3 Exterior forms and exterior differentials . . . . .	9
1.3.1 Exterior differential operators . . . . .	9
1.3.2 de Rham theorem . . . . .	10
1.4 Vector bundles and connections . . . . .	10
1.4.1 Connection of the vector bundle . . . . .	10
1.4.2 Curvature of a connection . . . . .	12
Exercises . . . . .	13
2. Finsler Metrics	15
2.1 Finsler metrics . . . . .	15
2.1.1 Finsler metrics . . . . .	15
2.1.2 Examples of Finsler metrics . . . . .	17
2.2 Cartan torsion . . . . .	22
2.2.1 Cartan torsion . . . . .	22
2.2.2 Deicke theorem . . . . .	24

2.3	Hilbert form and sprays . . . . .	28
2.3.1	Hilbert form . . . . .	28
2.3.2	Sprays . . . . .	30
2.4	Geodesics . . . . .	31
2.4.1	Geodesics . . . . .	31
2.4.2	Geodesic coefficients . . . . .	32
2.4.3	Geodesic completeness . . . . .	33
	Exercises . . . . .	36
3.	Connections and Curvatures . . . . .	39
3.1	Connections . . . . .	39
3.1.1	Chern connection . . . . .	39
3.1.2	Berwald metrics and Landsberg metrics . . . . .	42
3.2	Curvatures . . . . .	44
3.2.1	Curvature form of the Chern connection . . . . .	44
3.2.2	Flag curvature and Ricci curvature . . . . .	45
3.3	Bianchi identities . . . . .	47
3.3.1	Covariant differentiation . . . . .	47
3.3.2	Bianchi identities . . . . .	51
3.3.3	Other formulas . . . . .	53
3.4	Legendre transformation . . . . .	54
3.4.1	The dual norm in the dual space . . . . .	54
3.4.2	Legendre transformation . . . . .	55
3.4.3	Example . . . . .	56
	Exercises . . . . .	57
4.	<b>S</b> -Curvature . . . . .	59
4.1	Volume measures . . . . .	59
4.1.1	Busemann-Hausdorff volume element . . . . .	59
4.1.2	The volume element induced from $SM$ . . . . .	62
4.2	<b>S</b> -curvature . . . . .	65
4.2.1	Distortion . . . . .	65
4.2.2	<b>S</b> -curvature and <b>E</b> -curvature . . . . .	65
4.3	Isotropic <b>S</b> -curvature . . . . .	68
4.3.1	Isotropic <b>S</b> -curvature and isotropic <b>E</b> -curvature . . . . .	68
4.3.2	Randers metrics of isotropic <b>S</b> -curvature . . . . .	69
4.3.3	Geodesic flow . . . . .	72
	Exercises . . . . .	75

5.	Riemann Curvature	77
5.1	The second variation of arc length . . . . .	77
5.1.1	The second variation of length . . . . .	77
5.1.2	Elements of curvature and topology . . . . .	80
5.2	Scalar flag curvature . . . . .	83
5.2.1	Schur theorem . . . . .	83
5.2.2	Constant flag curvature . . . . .	85
5.3	Global rigidity results . . . . .	89
5.3.1	Flag curvature with special conditions . . . . .	89
5.3.2	Manifolds with non-positive flag curvature . . . . .	91
5.4	Navigation . . . . .	94
5.4.1	Navigation problem . . . . .	94
5.4.2	Randers metrics and navigation . . . . .	99
5.4.3	Ricci curvature and Einstein metrics . . . . .	104
	Exercises . . . . .	112

**Further Studies** **115**

6.	Projective Changes	117
6.1	The projective equivalence . . . . .	117
6.1.1	Projective equivalence . . . . .	117
6.1.2	Projective invariants . . . . .	119
6.2	Projectively flat metrics . . . . .	123
6.2.1	Projectively flat metrics . . . . .	123
6.2.2	Projectively flat metrics with constant flag curvature	126
6.3	Projectively flat metrics with almost isotropic <b>S</b> -curvature	136
6.3.1	Randers metrics with almost isotropic <b>S</b> -curvature	136
6.3.2	Projectively flat metrics with almost isotropic <b>S</b> -curvature . . . . .	141
6.4	Some special projectively equivalent Finsler metrics . . . . .	144
6.4.1	Projectively equivalent Randers metrics . . . . .	144
6.4.2	The projective equivalence of $(\alpha, \beta)$ -metrics . . . . .	148
6.4.3	The projective equivalence of quadratic $(\alpha, \beta)$ -metrics . . . . .	152
	Exercises . . . . .	155

7.	Comparison Theorems	157
7.1	Volume comparison theorems for Finsler manifolds . . . . .	157
7.1.1	The Jacobian of the exponential map . . . . .	157
7.1.2	Distance function and comparison theorems . . . . .	162
7.1.3	Volume comparison theorems . . . . .	165
7.2	Berger-Kazdan comparison theorems . . . . .	174
7.2.1	The Kazdan inequality . . . . .	174
7.2.2	The rigidity of reversible Finsler manifolds . . . . .	180
7.2.3	The Berger-Kazdan comparison theorem . . . . .	185
	Exercises . . . . .	190
8.	Fundamental Groups of Finsler Manifolds	191
8.1	Fundamental groups of Finsler manifolds . . . . .	191
8.1.1	Fundamental groups and covering spaces . . . . .	191
8.1.2	Algebraic norms and geometric norms . . . . .	194
8.1.3	Growth of fundamental groups . . . . .	199
8.2	Entropy and finiteness of fundamental group . . . . .	205
8.2.1	Entropy of fundamental group . . . . .	205
8.2.2	The first Betti number . . . . .	209
8.2.3	Finiteness of fundamental group . . . . .	213
8.3	Gromov pre-compactness theorems . . . . .	218
8.3.1	General metric spaces . . . . .	218
8.3.2	$\delta$ -Gromov-Hausdorff convergence . . . . .	224
8.3.3	Pre-compactness of Finsler manifolds . . . . .	231
8.3.4	On the Gauss-Bonnet-Chern theorem . . . . .	233
	Exercises . . . . .	235
9.	Minimal Immersions and Harmonic Maps	237
9.1	Isometric immersions . . . . .	237
9.1.1	Finsler submanifolds . . . . .	237
9.1.2	The variation of the volume . . . . .	241
9.1.3	Non-existence of compact minimal submanifolds . . . . .	245
9.2	Rigidity of minimal submanifolds . . . . .	246
9.2.1	Minimal surfaces in Minkowski spaces . . . . .	246
9.2.2	Minimal surfaces in $(\alpha, \beta)$ -spaces . . . . .	249
9.2.3	Minimal surfaces in special Minkowskian $(\alpha, \beta)$ - spaces . . . . .	253
9.3	Harmonic maps . . . . .	259

9.3.1	A divergence formula . . . . .	259
9.3.2	Harmonic maps . . . . .	262
9.3.3	Composition maps . . . . .	267
9.4	Second variation of harmonic maps . . . . .	270
9.4.1	The second variation . . . . .	270
9.4.2	Stress-energy tensor . . . . .	273
9.5	Harmonic maps between complex Finsler manifolds . . . . .	276
9.5.1	Complex Finsler manifolds . . . . .	276
9.5.2	Harmonic maps between complex Finsler manifolds . . . . .	280
9.5.3	Holomorphic maps . . . . .	287
	Exercises . . . . .	289
10.	Einstein Metrics . . . . .	293
10.1	Projective rigidity and $m$ -th root metrics . . . . .	293
10.1.1	Projective rigidity of Einstein metrics . . . . .	293
10.1.2	$m$ -th root Einstein metrics . . . . .	296
10.2	The Ricci rigidity and Douglas-Einstein metrics . . . . .	298
10.2.1	The Ricci rigidity . . . . .	298
10.2.2	Douglas $(\alpha, \beta)$ -metrics . . . . .	300
10.3	Einstein $(\alpha, \beta)$ -metrics . . . . .	305
10.3.1	Polynomial $(\alpha, \beta)$ -metrics . . . . .	305
10.3.2	Kropina metrics . . . . .	307
	Exercises . . . . .	317
11.	Miscellaneous Topics . . . . .	319
11.1	Conformal changes . . . . .	319
11.1.1	Conformal changes . . . . .	319
11.1.2	Conformally flat metrics . . . . .	322
11.1.3	Conformally flat $(\alpha, \beta)$ -metrics . . . . .	328
11.2	Conformal vector fields . . . . .	330
11.2.1	Conformal vector fields . . . . .	330
11.2.2	Conformal vector fields on a Randers manifold . . . . .	338
11.3	A class of critical Finsler metrics . . . . .	344
11.3.1	The Einstein-Hilbert functional . . . . .	344
11.3.2	Some special $\mathcal{E}$ -critical metrics . . . . .	349
11.4	The first eigenvalue of Finsler Laplacian and the generalized maximal principle . . . . .	351
11.4.1	Finsler Laplacian and weighted Ricci curvature . . . . .	351

11.4.2	Lichnerowicz-Obata estimates . . . . .	354
11.4.3	Li-Yau-Zhong-Yang type estimates . . . . .	360
11.4.4	Mckean type estimates . . . . .	364
	Exercises . . . . .	365
Appendix A	Maple Program	367
A.1	Spray coefficients of two-dimensional Finsler metrics . . .	367
A.2	Gauss curvature . . . . .	372
A.3	Spray coefficients of $(\alpha, \beta)$ -metrics . . . . .	374
	<i>Bibliography</i>	379
	<i>Index</i>	389

PART I  
**Foundations**



## Chapter 1

# Differentiable Manifolds

### 1.1 Differentiable manifolds

#### 1.1.1 Differentiable manifolds

**Definition 1.1.** An  $n$ -dimensional *differentiable manifold* is a Hausdorff topological space  $M$  and a family of diffeomorphisms  $\phi_\alpha : U_\alpha \subset M \rightarrow \phi_\alpha(U_\alpha) \subset \mathbb{R}^n$  such that the following conditions are satisfied:

- (i)  $\{U_\alpha\}$  is an open covering of  $M$ , i.e.,  $U_\alpha$  is open and  $\bigcup_\alpha U_\alpha = M$ ;
- (ii) If  $U_\alpha \cap U_\beta = V \neq \emptyset$ , then  $\phi_\alpha(V)$  and  $\phi_\beta(V)$  are open sets in  $\mathbb{R}^n$ , and  $\phi_\alpha \circ \phi_\beta^{-1}|_{\phi_\beta(V)}$  is a diffeomorphism;
- (iii) The family  $\{(U_\alpha, \phi_\alpha)\}$  is maximal relative to (i) and (ii).

For a point  $p \in U_\alpha$ ,  $(U_\alpha, \phi_\alpha)$  is called a *coordinate neighborhood* of  $p$ , and the coordinate of  $\phi_\alpha(q) \in \phi_\alpha(U_\alpha)$ ,  $\forall q \in U_\alpha$ , in  $\mathbb{R}^n$  can be viewed as the coordinate of  $q \in U_\alpha \subset M$ . A family  $\{(U_\alpha, \phi_\alpha)\}$  satisfying (i) and (ii) is called a *differentiable structure* on  $M$ . The condition (iii) is included for purely technical reasons.

#### 1.1.2 Examples of differentiable manifolds

**Example 1.1.** The Euclidean space  $\mathbb{R}^n$ .

**Example 1.2.** The unit sphere in  $\mathbb{R}^{n+1}$

$$\mathbb{S}^n = \left\{ (y^1, \dots, y^{n+1}) \in \mathbb{R}^{n+1} \mid \sum_{\alpha=1}^{n+1} (y^\alpha)^2 = 1 \right\} \subset \mathbb{R}^{n+1}.$$

Take the topology of  $\mathbb{S}^n$  as the sub-topology in  $\mathbb{R}^{n+1}$ , i.e.,  $U \subset \mathbb{S}^n$  is open if and only if there is an open  $\tilde{U} \subset \mathbb{R}^{n+1}$  such that  $U = \tilde{U} \cap \mathbb{S}^n$ . Thus,  $\mathbb{S}^n$  is a Hausdorff topological space. We are going to introduce a differentiable structure on  $\mathbb{S}^n$ .

For all  $1 \leq \alpha \leq n+1$ , let

$$\tilde{U}_\alpha^+ = \{(y^1, \dots, y^{n+1}) | y^\alpha > 0\}, \quad \tilde{U}_\alpha^- = \{(y^1, \dots, y^{n+1}) | y^\alpha < 0\}.$$

$\tilde{U}_\alpha^\pm$  are two open sets in  $\mathbb{R}^{n+1}$  separated by the hyperplane  $y^\alpha = 0$ . Then the family  $\{U_\alpha^\pm = \tilde{U}_\alpha^\pm \cap \mathbb{S}^n\}$  covers  $\mathbb{S}^n$ . Take the orthogonal projection  $\phi_\alpha^\pm : U_\alpha^\pm \rightarrow \mathbb{R}^n$

$$\phi_\alpha^\pm(y^1, \dots, y^{n+1}) = (y^1, \dots, \widehat{y^\alpha}, \dots, y^{n+1}),$$

where  $\widehat{y^\alpha}$  means the corresponding term is omitted. It is easy to see that  $\phi_\alpha^\pm$  are diffeomorphisms from  $U_\alpha^\pm$  to the open set  $W_\alpha = \{(y^1, \dots, \widehat{y^\alpha}, \dots, y^{n+1}) \in \mathbb{R}^n | \sum_{\beta \neq \alpha} (y^\beta)^2 < 1\}$ . Moreover, coordinate transformations are smooth. In fact, for example, the transformation  $\phi_2^- \circ (\phi_1^+)^{-1}$  on  $U_2^- \cap U_1^+$  is

$$\begin{aligned} (y^2, \dots, y^{n+1}) &\xrightarrow{(\phi_1^+)^{-1}} \left( \sqrt{1 - \sum_{\alpha=2}^{n+1} (y^\alpha)^2}, y^2, \dots, y^{n+1} \right) \\ &\xrightarrow{\phi_2^-} \left( \sqrt{1 - \sum_{\alpha=2}^{n+1} (y^\alpha)^2}, y^3, \dots, y^{n+1} \right). \end{aligned}$$

Instead of  $(y^2, \dots, y^{n+1})$  and  $(y^1, y^3, \dots, y^{n+1})$  on  $U_1^+$  and  $U_2^-$ , we use  $(x^1, x^2, \dots, x^n)$  and  $(\bar{x}^1, \bar{x}^2, \dots, \bar{x}^n)$ , respectively. Then we have

$$\bar{x}^1 = \sqrt{1 - \sum_{i=1}^n (x^i)^2}, \quad \bar{x}^j = x^j, \quad j = 2, \dots, n.$$

Hence,  $\mathbb{S}^n$  is an  $n$ -dimensional smooth manifold.

**Example 1.3.** The real projective space  $\mathbb{R}P^n$ .

This is the set of lines of  $\mathbb{R}^{n+1}$  that pass through the origin  $O = (0, \dots, 0)$ . Thus,  $\mathbb{R}P^n$  can be viewed as the quotient space of the unit sphere  $\mathbb{S}^n = \{p \in \mathbb{R}^{n+1} | |p| = 1\}$  by the equivalence relation  $A$  that identifies  $p \in \mathbb{S}^n$  with its antipodal point  $A(p) = -p$ . Indeed, each line that passes through the origin determines two antipodal points and  $\mathbb{R}P^n = \mathbb{S}^n/A$ .

Let  $\pi : \mathbb{S}^n \rightarrow \mathbb{R}P^n$  be the canonical projection, i.e.,  $\pi(p) = \{p, -p\}$ . By using the differentiable structure of  $\mathbb{S}^n$  as in Example 1.2, one can see that  $\pi(U_\alpha^+) = \pi(U_\alpha^-)$  and  $\{\pi(U_\alpha^\pm)\}$  covers  $\mathbb{R}P^n$ . Thus,  $\phi_\alpha = \phi_\alpha^\pm \circ \pi^{-1}$  is a diffeomorphism from  $\pi(U_\alpha^\pm)$  to  $W_\alpha \subset \mathbb{R}^n$ . This gives a differentiable structure on  $\mathbb{R}P^n$ . Hence,  $\mathbb{R}P^n$  is a smooth manifold, and  $\mathbb{S}^n$  can be viewed as the two-fold covering of  $\mathbb{R}P^n$ .

Let  $(x^1, \dots, x^{n+1})$  be the coordinates of  $\mathbb{R}^{n+1}$ . Define an equivalent relation  $\sim$  in  $\mathbb{R}^{n+1} \setminus \{0\}$  as follows: For any two points  $\bar{x} = (\bar{x}^1, \dots, \bar{x}^{n+1})$  and  $x = (x^1, \dots, x^{n+1})$ , if there is a real number  $\lambda \neq 0$  such that  $\bar{x} = \lambda x$ , then  $\bar{x} \sim x$ . Thus, the real projective space  $\mathbb{R}P^n$  is the quotient space  $(\mathbb{R}^{n+1} \setminus \{0\}) / \sim$ , and  $(x^1, \dots, x^{n+1}) \neq (0, \dots, 0)$  is called the *homogeneous coordinate* of  $\mathbb{R}P^n$ .

**Example 1.4.** The *Grassmann manifold*

$G_{k,n} = \{k\text{-dimensional linear subspaces in } \mathbb{R}^n \text{ passing through the origin}\}.$

Fix an orthonormal base  $\{e_1, \dots, e_n\}$  in  $\mathbb{R}^n$ . The orthogonal group  $O(n)$  acts on  $\mathbb{R}^n$  via the matrix multiplication. On the other hand, any non-singular linear transformation  $\sigma \in O(n)$  maps a  $k$ -dimensional linear subspace  $P$  to a  $k$ -dimensional linear subspace  $Q = \mathcal{A}(\sigma, P)$ , where

$$\mathcal{A} : O(n) \times G_{k,n} \rightarrow G_{k,n}.$$

Let  $P_0$  denote the  $k$ -dimensional linear subspace spanned by  $e_1, \dots, e_k$ . Let  $H \subset O(n)$  be the subgroup preserving  $P_0$  invariant. We have

$$H = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \in O(n) \mid \alpha \in O(k), \beta \in O(n-k) \right\}.$$

Thus,  $H$  is the closed subgroup of  $O(n)$  which is isomorphic to  $O(k) \times O(n-k)$ . The map  $\sigma(O(k) \times O(n-k)) \mapsto \mathcal{A}(\sigma, P_0)$  from the quotient space  $O(n)/O(k) \times O(n-k)$  to  $G_{k,n}$  is bijective. We make the above map diffeomorphic, so that  $G_{k,n}$  is a  $k(n-k)$ -dimensional differentiable manifold. It is easy to see that the manifold structure of  $G_{k,n}$  is independent of a particular choice of the base in  $\mathbb{R}^n$ . In particular,  $G_{1,n+1} = \mathbb{R}P^n$ .

**Example 1.5.** The *product manifold*  $M \times N$ .

Let  $\{(U_\alpha, \phi_\alpha)\}$  and  $\{(V_\beta, \psi_\beta)\}$  be differentiable structures of  $M$  and  $N$ , respectively. Clearly, we have  $\bigcup_{\alpha, \beta} U_\alpha \times V_\beta = M \times N$ . Define the diffeomorphism  $\phi_\alpha \times \psi_\beta : U_\alpha \times V_\beta \rightarrow \phi_\alpha(U_\alpha) \times \psi_\beta(V_\beta) \subset \mathbb{R}^m \times \mathbb{R}^n = \mathbb{R}^{m+n}$  as follows:

$$(\phi_\alpha \times \psi_\beta)(p, q) = (\phi_\alpha(p), \psi_\beta(q)), \quad \forall (p, q) \in U_\alpha \times V_\beta.$$

Then  $\{(U_\alpha \times V_\beta, \phi_\alpha \times \psi_\beta)\}$  is a differentiable structure of  $M \times N$ .

An important example of product manifolds is the  $n$ -torus  $T^n = \mathbb{S}^1 \times \dots \times \mathbb{S}^1$ , which is the  $n$ -multiple Descartes product of circles.

Let  $f : M \rightarrow N$  be a map from a differentiable manifold  $M$  to another  $N$ . For  $p \in M$ , if there are a coordinate neighborhood  $(U, \phi)$  of  $p$  and a