

Phillip Griffiths, Joseph Harris

Principles of Algebraic Geometry

代数几何原理

WILEY



世界图书出版公司
www.wpcbj.com.cn

PRINCIPLES OF ALGEBRAIC GEOMETRY

PHILLIP GRIFFITHS and JOSEPH HARRIS

Harvard University

Wiley Classics Library Edition Published 1994



A WILEY-INTERSCIENCE PUBLICATION

JOHN WILEY & SONS, INC.

New York • Chichester • Brisbane • Toronto • Singapore

This text printed on acid-free paper.

Copyright © 1978 by John Wiley & Sons, Inc.

Wiley Classics Library edition published 1994.

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008.

Library of Congress Cataloging in Publication Data

Griffiths, Phillip, 1938–

Principles of algebraic geometry.

(Pure and applied mathematics)

“A Wiley-Interscience publication.”

Includes bibliographical references.

1. Geometry, Algebraic. I. Harris, Joseph,

1951– joint author. II. Title.

QA564.G64

516'.35

78-6993

ISBN 0-471-05059-8

All Rights Reserved. AUTHORIZED REPRINT OF THE EDITION PUBLISHED BY JOHN WILEY & SONS, INC. No part of this book may be reproduced in and form without the written permission of John Wiley & Sons, Inc.
reprint run: 1000 copies List Price: 149.00(RMB)

PREFACE

Algebraic geometry is among the oldest and most highly developed subjects in mathematics. It is intimately connected with projective geometry, complex analysis, topology, number theory, and many other areas of current mathematical activity. Moreover, in recent years algebraic geometry has undergone vast changes in style and language. For these reasons there has arisen about the subject a reputation of inaccessibility. This book gives a presentation of some of the main general results of the theory accompanied by—and indeed with special emphasis on—the applications to the study of interesting examples and the development of computational tools.

A number of principles guided the preparation of the book. One was to develop only that general machinery necessary to study the concrete geometric questions and special classes of algebraic varieties around which the presentation was centered.

A second was that there should be an alternation between the general theory and study of examples, as illustrated by the table of contents. The subject of algebraic geometry is especially notable for the balance provided on the one hand by the intricacy of its examples and on the other by the symmetry of its general patterns; we have tried to reflect this relationship in our choice of topics and order of presentation.

A third general principle was that this volume should be self-contained. In particular any “hard” result that would be utilized should be fully proved. A difficulty a student often faces in a subject as diverse as algebraic geometry is the profusion of cross-references, and this is one reason for attempting to be self-contained. Similarly, we have attempted to avoid allusions to, or statements without proofs of, related results. This book is in no way meant to be a survey of algebraic geometry, but rather is designed to develop a working facility with specific geometric questions. Our approach to the subject is initially analytic: Chapters 0 and 1 treat the basic techniques and results of complex manifold theory, with some emphasis on results applicable to projective varieties. Beginning in Chapter 2 with the theory of Riemann surfaces and algebraic curves, and continuing in Chapters 4 and 6 on algebraic surfaces and the quadric line

complex, our treatment becomes increasingly geometric along classical lines. Chapters 3 and 5 continue the analytic approach, progressing to more special topics in complex manifolds.

Several important topics have been entirely omitted. The most glaring are the arithmetic theory of algebraic varieties, moduli questions, and singularities. In these cases the necessary techniques are not fully developed here. Other topics, such as uniformization and automorphic forms or monodromy and mixed Hodge structures have been omitted, although the necessary techniques are for the most part available.

We would like to thank Giuseppe Canuto, S. S. Chern, Maurizio Cornalba, Ran Donagi, Robin Hartshorne, Bill Hoffman, David Morrison, David Mumford, Arthur Ogus, Ted Shifrin, and Loring Tu for many fruitful discussions; Ruth Suzuki for her wonderful typing; and the staff of John Wiley, especially Beatrice Shube, for enormous patience and skill in converting a very rough manuscript into book form.

PHILLIP GRIFFITHS
JOSEPH HARRIS

May 1978
Cambridge, Massachusetts

PRINCIPLES OF
ALGEBRAIC GEOMETRY

CONTENTS

CHAPTER 0 FOUNDATIONAL MATERIAL	1
1. Rudiments of Several Complex Variables	
Cauchy's Formula and Applications	2
Several Variables	6
Weierstrass Theorems and Corollaries	7
Analytic Varieties	12
2. Complex Manifolds	
Complex Manifolds	14
Submanifolds and Subvarieties	18
De Rham and Dolbeault Cohomology	23
Calculus on Complex Manifolds	27
3. Sheaves and Cohomology	
Origins: The Mittag-Leffler Problem	34
Sheaves	35
Cohomology of Sheaves	38
The de Rham Theorem	43
The Dolbeault Theorem	45
4. Topology of Manifolds	
Intersection of Cycles	49
Poincaré Duality	53
Intersection of Analytic Cycles	60
5. Vector Bundles, Connections, and Curvature	
Complex and Holomorphic Vector Bundles	66
Metrics, Connections, and Curvature	71
6. Harmonic Theory on Compact Complex Manifolds	
The Hodge Theorem	80
Proof of the Hodge Theorem I: Local Theory	84
	vii

Proof of the Hodge Theorem II: Global Theory	92
Applications of the Hodge Theorem	100
7. Kähler Manifolds	
The Kähler Condition	106
The Hodge Identities and the Hodge Decomposition	111
The Lefschetz Decomposition	118
CHAPTER 1 COMPLEX ALGEBRAIC VARIETIES	128
1. Divisors and Line Bundles	
Divisors	129
Line Bundles	132
Chern Classes of Line Bundles	139
2. Some Vanishing Theorems and Corollaries	
The Kodaira Vanishing Theorem	148
The Lefschetz Theorem on Hyperplane Sections	156
Theorem B	159
The Lefschetz Theorem on $(1, 1)$ -classes	161
3. Algebraic Varieties	
Analytic and Algebraic Varieties	164
Degree of a Variety	171
Tangent Spaces to Algebraic Varieties	175
4. The Kodaira Embedding Theorem	
Line Bundles and Maps to Projective Space	176
Blowing Up	182
Proof of the Kodaira Theorem	189
5. Grassmannians	
Definitions	193
The Cell Decomposition	194
The Schubert Calculus	197
Universal Bundles	207
The Plücker Embedding	209
CHAPTER 2 RIEMANN SURFACES AND ALGEBRAIC CURVES	212
1. Preliminaries	
Embedding Riemann Surfaces	213
The Riemann-Hurwitz Formula	216

CONTENTS

ix

The Genus Formula	219
Cases $g = 0, 1$	222
2. Abel's Theorem	
Abel's Theorem—First Version	224
The First Reciprocity Law and Corollaries	229
Abel's Theorem—Second Version	232
Jacobi Inversion	235
3. Linear Systems on Curves	
Reciprocity Law II	240
The Riemann-Roch Formula	243
Canonical Curves	246
Special Linear Systems I	249
Hyperelliptic Curves and Riemann's Count	253
Special Linear Systems II	259
4. Plücker Formulas	
Associated Curves	263
Ramification	264
The General Plücker Formulas I	268
The General Plücker Formulas II	271
Weierstrass Points	273
Plücker Formulas for Plane Curves	277
5. Correspondences	
Definitions and Formulas	282
Geometry of Space Curves	290
Special Linear Systems III	298
6. Complex Tori and Abelian Varieties	
The Riemann Conditions	300
Line Bundles on Complex Tori	307
Theta-Functions	317
The Group Structure on an Abelian Variety	324
Intrinsic Formulations	326
7. Curves and Their Jacobians	
Preliminaries	333
Riemann's Theorem	338
Riemann's Singularity Theorem	341
Special Linear Systems IV	349
Torelli's Theorem	359

CHAPTER 3 FURTHER TECHNIQUES	364
1. Distributions and Currents	
Definitions; Residue Formulas	366
Smoothing and Regularity	373
Cohomology of Currents	382
2. Applications of Currents to Complex Analysis	
Currents Associated to Analytic Varieties	385
Intersection Numbers of Analytic Varieties	392
The Levi Extension and Proper Mapping Theorems	395
3. Chern Classes	
Definitions	400
The Gauss Bonnet Formulas	409
Some Remarks—Not Indispensable—Concerning Chern Classes of Holomorphic Vector Bundles	416
4. Fixed-Point and Residue Formulas	
The Lefschetz Fixed-Point Formula	419
The Holomorphic Lefschetz Fixed-Point Formula	422
The Bott Residue Formula	426
The General Hirzebruch-Riemann-Roch Formula	435
5. Spectral Sequences and Applications	
Spectral Sequences of Filtered and Bigraded Complexes	438
Hypercohomology	445
Differentials of the Second Kind	454
The Leray Spectral Sequence	462
CHAPTER 4 SURFACES	469
1. Preliminaries	
Intersection Numbers, the Adjunction Formula, and Riemann-Roch	470
Blowing Up and Down	473
The Quadric Surface	478
The Cubic Surface	480
2. Rational Maps	
Rational and Birational Maps	489
Curves on an Algebraic Surface	498
The Structure of Birational Maps Between Surfaces	510
3. Rational Surfaces I	
Noether's Lemma	513
Rational Ruled Surfaces	514

CONTENTS

xi

The General Rational Surface	520
Surfaces of Minimal Degree	522
Curves of Maximal Genus	527
Steiner Constructions	528
The Enriques-Petri Theorem	533
4. Rational Surfaces II	
The Castelnuovo-Enriques Theorem	536
The Enriques Surface	541
Cubic Surfaces Revisited	545
The Intersection of Two Quadrics in \mathbb{P}^4	550
5. Some Irrational Surfaces	
The Albanese Map	552
Irrational Ruled Surfaces	553
A Brief Introduction to Elliptic Surfaces	564
Kodaira Number and the Classification Theorem I	572
The Classification Theorem II	582
K-3 Surfaces	590
Enriques Surfaces	594
6. Noether's Formula	
Noether's Formula for Smooth Hypersurfaces	600
Blowing Up Submanifolds	602
Ordinary Singularities of Surfaces	611
Noether's Formula for General Surfaces	618
Some Examples	628
Isolated Singularities of Surfaces	636
CHAPTER 5 RESIDUES	647
1. Elementary Properties of Residues	
Definition and Cohomological Interpretation	649
The Global Residue Theorem	655
The Transformation Law and Local Duality	656
2. Applications of Residues	
Intersection Numbers	662
Finite Holomorphic Mappings	667
Applications to Plane Projective Geometry	670
3. Rudiments of Commutative and Homological Algebra with Applications	
Commutative Algebra	678
Homological Algebra	682
The Koszul Complex and Applications	687
A Brief Tour Through Coherent Sheaves	695

4. Global Duality	
Global Ext	705
Explanation of the General Global Duality Theorem	707
Global Ext and Vector Fields with Isolated Zeros	708
Global Duality and Superabundance of Points on a Surface	712
Extensions of Modules	722
Points on a Surface and Rank-Two Vector Bundles	726
Residues and Vector Bundles	729
CHAPTER 6 THE QUADRIC LINE COMPLEX	733
1. Preliminaries: Quadrics	
Rank of a Quadric	734
Linear Spaces on Quadrics	735
Linear Systems of Quadrics	741
Lines on Linear Systems of Quadrics	746
The Problem of Five Conics	749
2. The Quadric Line Complex: Introduction	
Geometry of the Grassmannian $G(2,4)$	756
Line Complexes	759
The Quadric Line Complex and Associated Kummer Surface I	762
Singular Lines of the Quadric Line Complex	767
Two Configurations	773
3. Lines on the Quadric Line Complex	
The Variety of Lines on the Quadric Line Complex	778
Curves on the Variety of Lines	780
Two Configurations Revisited	784
The Group Law	787
4. The Quadric Line Complex: Reprise	
The Quadric Line Complex and Associated Kummer Surface II	791
Rationality of the Quadric Line Complex	796
INDEX	805

0

FOUNDATIONAL MATERIAL

In this chapter we sketch the foundational material from several complex variables, complex manifold theory, topology, and differential geometry that will be used in our study of algebraic geometry. While our treatment is for the most part self-contained, it is tacitly assumed that the reader has some familiarity with the basic objects discussed. The primary purpose of this chapter is to establish our viewpoint and to present those results needed in the form in which they will be used later on. There are, broadly speaking, four main points:

1. *The Weierstrass theorems and corollaries*, discussed in Sections 1 and 2. These give us our basic picture of the local character of analytic varieties. The theorems themselves will not be quoted directly later, but the picture—for example, the local representation of an analytic variety as a branched covering of a polydisc—is fundamental. The foundations of local analytic geometry are further discussed in Chapter 5.

2. *Sheaf theory*, discussed in Section 3, is an important tool for relating the analytic, topological, and geometric aspects of an algebraic variety. A good example is the *exponential sheaf sequence*, whose individual terms \mathbb{Z} , \mathcal{O} , and \mathcal{O}^* reflect the topological, analytic, and geometric structures of the underlying variety, respectively.

3. *Intersection theory*, discussed in Section 4, is a cornerstone of classical algebraic geometry. It allows us to treat the incidence properties of algebraic varieties, a priori a geometric question, in topological terms.

4. *Hodge theory*, discussed in Sections 6 and 7. By far the most sophisticated technique introduced in this chapter, Hodge theory has, in the present context, two principal applications: first, it gives us the *Hodge decomposition* of the cohomology of a Kähler manifold; then, together with the formalism introduced in Section 5, it gives the vanishing theorems of the next chapter.

1. RUDIMENTS OF SEVERAL COMPLEX VARIABLES

Cauchy's Formula and Applications

NOTATION. We will write $z = (z_1, \dots, z_n)$ for a point in \mathbb{C}^n , with

$$z_i = x_i + \sqrt{-1} y_i;$$

$$\|z\|^2 = (z, z) = \sum_{i=1}^n |z_i|^2.$$

For U an open set in \mathbb{C}^n , write $C^\infty(U)$ for the set of C^∞ functions defined on U ; $C^\infty(\bar{U})$ for the set of C^∞ functions defined in some neighborhood of the closure \bar{U} of U .

The cotangent space to a point in $\mathbb{C}^n \cong \mathbb{R}^{2n}$ is spanned by $\{dx_i, dy_i\}$; it will often be more convenient, however, to work with the complex basis

$$dz_i = dx_i + \sqrt{-1} dy_i, \quad d\bar{z}_i = dx_i - \sqrt{-1} dy_i$$

and the dual basis in the tangent space

$$\frac{\partial}{\partial z_i} = \frac{1}{2} \left(\frac{\partial}{\partial x_i} - \sqrt{-1} \frac{\partial}{\partial y_i} \right), \quad \frac{\partial}{\partial \bar{z}_i} = \frac{1}{2} \left(\frac{\partial}{\partial x_i} + \sqrt{-1} \frac{\partial}{\partial y_i} \right).$$

With this notation, the formula for the total differential is

$$df = \sum_i \frac{\partial f}{\partial z_i} dz_i + \sum_j \frac{\partial f}{\partial \bar{z}_j} d\bar{z}_j.$$

In one variable, we say a C^∞ function f on an open set $U \subset \mathbb{C}$ is *holomorphic* if f satisfies the Cauchy-Riemann equations $\partial f / \partial \bar{z} = 0$. Writing $f(z) = u(z) + \sqrt{-1} v(z)$, this amounts to

$$\operatorname{Re} \left(\frac{\partial f}{\partial \bar{z}} \right) = \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} = 0,$$

$$\operatorname{Im} \left(\frac{\partial f}{\partial \bar{z}} \right) = \frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} = 0.$$

We say f is *analytic* if, for all $z_0 \in U$, f has a local series expansion in $z - z_0$, i.e.,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

in some disc $\Delta(z_0, \varepsilon) = \{z : |z - z_0| < \varepsilon\}$, where the sum converges absolutely and uniformly. The first result is that f is analytic if and only if it is holomorphic; to show this, we use the

Cauchy Integral Formula. For Δ a disc in \mathbb{C} , $f \in C^\infty(\bar{\Delta})$, $z \in \Delta$,

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w) dw}{w - z} + \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w - z},$$

where the line integrals are taken in the counterclockwise direction (the fact that the last integral is defined will come out in the proof).

Proof. The proof is based on Stokes' formula for a differential form with singularities, a method which will be formalized in Chapter 3. Consider the differential form

$$\eta = \frac{1}{2\pi\sqrt{-1}} \frac{f(w) dw}{w-z};$$

we have for $z \neq w$

$$\frac{\partial}{\partial \bar{w}} \left(\frac{1}{w-z} \right) = 0$$

and so

$$d\eta = -\frac{1}{2\pi\sqrt{-1}} \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z}.$$

Let $\Delta_\epsilon = \Delta(z, \epsilon)$ be the disc of radius ϵ around z . The form η is C^∞ in $\Delta - \Delta_\epsilon$, and applying Stokes' theorem we obtain

$$\begin{aligned} \frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta_\epsilon} \frac{f(w) dw}{w-z} &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta} \frac{f(w) dw}{w-z} \\ &+ \frac{1}{2\pi\sqrt{-1}} \int_{\Delta - \Delta_\epsilon} \frac{\partial f}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z}. \end{aligned}$$

Setting $w - z = re^{i\theta}$,

$$\frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta_\epsilon} \frac{f(w) dw}{w-z} = \frac{1}{2\pi} \int_0^{2\pi} f(z + \epsilon e^{i\theta}) d\theta,$$

which tends to $f(z)$ as $\epsilon \rightarrow 0$; moreover,

$$dw \wedge d\bar{w} = -2\sqrt{-1} dx \wedge dy = -2\sqrt{-1} r dr \wedge d\theta$$

so

$$\left| \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z} \right| = 2 \left| \frac{\partial f}{\partial \bar{w}} dr \wedge d\theta \right| \leq c |dr \wedge d\theta|.$$

Thus $(\partial f / \partial \bar{w})(dw \wedge d\bar{w}) / (w - z)$ is absolutely integrable over Δ , and

$$\int_{\Delta_\epsilon} \frac{\partial f}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z} \rightarrow 0$$

as $\epsilon \rightarrow 0$; the result follows. Q.E.D.

Now we can prove the

Proposition. For U an open set in \mathbb{C} and $f \in C^\infty(U)$, f is holomorphic if and only if f is analytic.

Proof. Suppose first that $\partial f/\partial \bar{z} = 0$. Then for $z_0 \in U$, ε sufficiently small, and z in the disc $\Delta = \Delta(z_0, \varepsilon)$ of radius ε around z_0 ,

$$\begin{aligned} f(z) &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{w-z} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0) - (z-z_0)} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)\left(1 - \frac{z-z_0}{w-z_0}\right)} \\ &= \sum_{n=0}^{\infty} \left(\frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)^{n+1}} \right) (z-z_0)^n; \end{aligned}$$

so, setting

$$a_n = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)^{n+1}},$$

we have

$$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n$$

for $z \in \Delta$, where the sum converges absolutely and uniformly in any smaller disc.

Suppose conversely that $f(z)$ has a power series expansion

$$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n$$

for $z \in \Delta = \Delta(z_0, \varepsilon)$. Since $(\partial/\partial \bar{z})(z-z_0)^n = 0$, the partial sums of the expansion satisfy Cauchy's formula without the area integral, and by the uniform convergence of the sum in a neighborhood of z_0 the same is true of f , i.e.,

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{w-z}.$$

We can then differentiate under the integral sign to obtain

$$\frac{\partial}{\partial \bar{z}} f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{\partial}{\partial \bar{z}} \left(\frac{f(w)}{w-z} \right) dw = 0,$$

since for $z \neq w$

$$\frac{\partial}{\partial \bar{z}} \left(\frac{1}{w-z} \right) = 0.$$

Q.E.D.

We prove a final result in one variable, that given a C^∞ function g on a disc Δ the equation

$$\frac{\partial f}{\partial \bar{z}} = g$$

can always be solved on a slightly smaller disc; this is the

$\bar{\partial}$ -Poincaré Lemma in One Variable. Given $g(z) \in C^\infty(\bar{\Delta})$, the function

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{g(w)}{w-z} dw \wedge d\bar{w}$$

is defined and C^∞ in Δ and satisfies

$$\frac{\partial f}{\partial \bar{z}} = g.$$

Proof. For $z_0 \in \Delta$ choose ε such that the disc $\Delta(z_0, 2\varepsilon) \subset \Delta$ and write

$$g(z) = g_1(z) + g_2(z),$$

where $g_1(z)$ vanishes outside $\Delta(z_0, 2\varepsilon)$ and $g_2(z)$ vanishes inside $\Delta(z_0, \varepsilon)$. The integral

$$f_2(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} g_2(w) \frac{dw \wedge d\bar{w}}{w-z}$$

is well-defined and C^∞ for $z \in \Delta(z_0, \varepsilon)$; there we have

$$\frac{\partial}{\partial \bar{z}} f_2(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial}{\partial \bar{z}} \left(\frac{g_2(w)}{w-z} \right) dw \wedge d\bar{w} = 0.$$

Since $g_1(z)$ has compact support, we can write

$$\begin{aligned} \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} g_1(w) \frac{dw \wedge d\bar{w}}{w-z} &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} g_1(w) \frac{dw \wedge d\bar{w}}{w-z} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} g_1(u+z) \frac{du \wedge d\bar{u}}{u}, \end{aligned}$$

where $u = w - z$. Changing to polar coordinates $u = re^{i\theta}$ this integral becomes

$$f_1(z) = -\frac{1}{\pi} \int_{\mathbb{C}} g_1(z + re^{i\theta}) e^{-i\theta} dr \wedge d\theta,$$

which is clearly defined and C^∞ in z . Then

$$\begin{aligned} \frac{\partial f_1(z)}{\partial \bar{z}} &= -\frac{1}{\pi} \int_{\mathbb{C}} \frac{\partial g_1}{\partial \bar{z}}(z + re^{i\theta}) e^{-i\theta} dr \wedge d\theta \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial g_1}{\partial \bar{w}}(w) \frac{dw \wedge d\bar{w}}{w-z}; \end{aligned}$$