

GALATHEA REPORT

Scientific Results of the Danish Deep-Sea Expedition Round the World 1950-52

- Volume 1, 1957-1959. 260 pp., 8 pls. Price: 90 Danish Kr.
Volume 2, 1956. 253 pp., 4 pls. Price: 75 Danish Kr.
Volume 3, 1959. 87 pp., 60 pls. Price: 60 Danish Kr.
Volume 4, 1961. 226 pp., 14 pls. Price: 95 Danish Kr.
Volume 5, 1961. 209 pp., 10 pls. Price: 90 Danish Kr.
Volume 6, 1962. 320 pp., 19 pls. Price: 120 Danish Kr.
Volume 7, 1963 (in print)

ATLANTIDE-REPORT

Scientific Results of the Danish Expedition to the Coasts of Tropical West Africa 1945-46

- No. 1, 1950. 247 pp., 16 pls. Price: 35 Danish Kr.
No. 2, 1951. 303 pp., 2 pls. Price: 40 Danish Kr.
No. 3, 1955. 324 pp., 15 pls. Price: 50 Danish Kr.
No. 4, 1956. 212 pp., 5 pls. Price: 40 Danish Kr.
No. 5, 1959. 332 pp. Price: 60 Danish Kr.
No. 6, 1961. 262 pp., 9 pls. Price: 55 Danish Kr.
No. 7, 1963 (in print)

THE NATURAL HISTORY OF RENNELL ISLAND, BRITISH SOLOMON ISLANDS

Scientific Results of the Danish Rennell Expedition, 1951
and the British Museum (Nat. Hist.) Expedition, 1953

- Volume 1: Vertebrates, 1958. 228 pp., 9 pls. Price: 45 Danish Kr.
Volume 2: Invertebrates, 1959. 234 pp., 35 pls. Price: 50 Danish Kr.
Volume 3: Botany and Geology, 1960. 132 pp., 10 pls. Price: 30 Danish Kr.
Volume 4: Invertebrates, 1962. 120 pp., 18 pls. Price: 35 Danish Kr.

LIST OF DANISH VERTEBRATES

ANTON FR. BRUUN and J. R. PFAFF: Fishes

J. R. PFAFF: Amphibians and Reptiles

BERNT LØPPENTHIN: Birds

MAGNUS DEGERBØL: Mammals

180 pp., one chart. Price: 10.50 Danish Kr.

Rate of exchange: 7 Danish Kroner equal to about 7 British sh. or one U. S. A. dollar

DANISH SCIENCE PRESS, LTD.

(Dansk Videnskabs Forlag, A/S)

Copenhagen

GALATHEA REPORTS

VOLUME 1

- BRUUN, ANTON F.: General Introduction to the Reports and List of Deep-Sea Stations. 42 pp., 11 text-figs, 4 pls.
- NIELSEN, E. STEEMANN & E. AABYE JENSEN: Primary Oceanic Production. The Autotrophic Production of Organic Matter in the Oceans. 88 pp., 41 text-figs.
- BRAARUD, TRYGVE: A Red Water Organism from Walvis Bay. 2 pp., 2 text-figs.
- ZOBELL, CLAUDE E. & RICHARD Y. MORITA: Deep-Sea Bacteria. 16 pp., 4 text-figs.
- KILLERICH, A.: Bathymetric Features of the Philippine Trench. 17 pp., 12 text-figs., 4 pls.
- KRAMP, P.L.: *Stephanoscyphus* (Scyphozoa). 15 pp., 12 text-figs., 1 pl.
- MILLAR, R.H.: Ascidiacea. 21 pp., 20 text-figs., 1 pl.
- DAHL, ERIK: Amphipoda from Depths Exceeding 6000 Meters. 31 pp., 20 text-figs.
- PICKFORD, GRACE E.: Vampyromorpha. 11 pp., 1 text-fig.
- ROFEN, ROBERT R.: The Whale-Fishes: Families Cetomimidae, Barbourisiidae and Rondelctiidae (Order Cetunculi). 6 pp., 4 text-figs.

VOLUME 2

- CARLGRÉN, OSCAR: Actiniaria from Depths Exceeding 6000 Meters. 7 pp., 7 text-figs., 1 pl.
- KRAMP, P.L.: Hydroids from Depths Exceeding 6000 Meters. 4 pp., 6 text-figs.
- MADSEN, F.JENSENIUS: *Primnoella krampi* n.sp. A New Deep-Sea Octocoral. 2 pp., 1 text-fig.
- MADSEN, F.JENSENIUS: Echinoidea, Asteroidea, and Ophiuroidea from Depths Exceeding 6000 Meters. 11 pp., 5 text-figs., 1 pl.
- HANSEN, BENT: Holothurioidea from Depths Exceeding 6000 Meters. 22 pp., 25 text-figs.
- HANSEN, BENT & F.JENSENIUS MADSEN: On two Bathypelagic Holothurians from the South China Sea. *Galatheathuria* n.g. *aspera* (Théel) and *Enypniastes globosa* n.sp. 5 pp., 2 text-figs., 1 pl.
- GISLÉN, TORSTEN: Crinoids from Depths Exceeding 6000 Meters. 2 pp., 1 pl.
- KIRKEGAARD, J.B.: Benthic Polychæta from Depths Exceeding 6000 Meters. 16 pp., 13 text-figs.
- KIRKEGAARD, J.B.: Pogonophora. *Galathealinum bruuni* n. gen. n.sp., a New Representative of the Class. 5 pp., 2 text-figs.
- WOLFF, TORBEN: Isopoda from Depths Exceeding 6000 Meters. 73 pp., 56 text-figs.

- FAGE, LOUIS: Les Pycnogonides du genre *Nymphon*. 7 pp., 10 text-figs.
- FAGE, LOUIS: Les Pycnogonides (excl. le genre *Nymphon*). 16 pp., 22 text-figs.
- KIRKEGAARD, J.B.: Pogonophora. First Records from the Eastern Pacific. 4 pp., 4 text-figs.
- WOLFF, TORBEN: Crustacea Tanaidacea from Depths Exceeding 6000 Meters. 55 pp., 54 text-figs.
- WIESER, WOLFGANG: Some Free Living Marine Nematodes. 11 pp., 4 text-figs.

VOLUME 3

- LEMICHE, HENNING and KARL GEORG WINGSTRAND: The Anatomy of *Neopilina galathea* Lemche, 1957 (Mollusca Tryblidiacea). 63 pp., 1 text-fig., 56 pls.
- SCHMIDT, W.J.: Bemerkungen zur Schalenstruktur von *Neopilina galathea*. 5 pp., 2 pls.
- MUNK, O.: The Eyes of *Ipnots murrayi* Günther, 1878. 9 pp., 3 text-figs., 2 pls.

VOLUME 4

- KIRKEGAARD, J.B.: Pogonophora III. The Genus *Lamellisabella*. 4 pp., 3 text-figs.
- WOLFF, TORBEN: Description of a Remarkable Deep-Sea Hermit Crab, with Notes on the Evolution of the Paguridea. 22 pp., 11 text-figs.
- MADSEN, F.JENSENIUS: The Porcellanasteridae. A Monographic Revision of an Abyssal Group of Sea-Stars. 142 pp., 43 text-figs., 13 pls.
- MADSEN, F.JENSENIUS: On the Zoogeography and Origin of the Abyssal Fauna, in View of the Knowledge of the Porcellanasteridae. 42 pp., 2 text-figs.
- NIELSEN, JØRGEN: Heterosomata (Pisces). 8 pp., 3 text-figs., 1 pl.

VOLUME 5

- ÅKESSON, BERTIL: Some Observations on *Pelagosphaera* Larvae (Sipunculoidea). 11 pp., 1 text-fig., 6 pls.
- COHEN, DANIEL M.: On the Identity of the Species of the Fish Genus *Argentina* in the Indian Ocean. 3 pp., 1 text-fig.
- BARNARD, J.LAURENS: Gammaridean Amphipoda from Depths of 400 to 6000 Meters. 106 pp., 83 text-figs.
- WOLFF, TORBEN: Animal Life from a Single Abyssal Trawling. 34 pp., 26 text-figs., 4 pls.
- KNUDSEN, JØRGEN: The Bathyal and Abyssal *Xylophaga* (Pholadidae, Bivalvia). 47 pp., 41 text-figs.

LIST OF PAPERS RESULTING IN WHOLE OR IN PART FROM
THE GALATHEA DEEP-SEA EXPEDITION
ROUND THE WORLD 1950-52

(Supplementary to the list given in the Introduction to vol. 1, 1959, pp. 18-19).

- ABBOTT, R. TUCKER, 1958: Marine Mollusca of Rennell Island, Solomon Islands. - The Natural History of Rennell Island, British Solomon Islands, 2: 203-206, 1 pl.
- BALFOUR-BROWNE, J., 1958: Coleoptera (Pars). - Ibid.: 127-133.
- 1962: Coleoptera from Rennell and Bellona Islands, Supplement. - Ibid. 4: 85-91.
- BAUCHOT, M. L. & M. BLANC, 1961: Poissons marins de l'Est Atlantique tropical. I. Labroidei (Téléostéens Perciformes). - Atlantide Rep., 6: 43-64, 6 text-figs.
- 1961: Poissons marins de l'Est Atlantique tropical. II. Percoides (Téléostéens Perciformes). 1ère partie. - Ibid.: 65-100, 4 text-figs., 1 pl.
- BRUUN, A. F., 1958: On the Restricted Distribution of two Deep-Sea Fishes, *Borophryne apogon* and *Stomias colubrinus*. - Sears Found.: Journ. Mar. Res., 17: 103-112, 2 text-figs.
- 1960: Exploring the World's Deepest Trenches. - The Courier, UNESCO, Nos. 7-8: 20-22, 5 text-figs.
- & TORBEN WOLFF, 1961: Abyssal Benthic Organisms: Nature, Origin, Distribution, and Influence on Sedimentation. - In: Oceanography (M. Sears ed.), Publ. No. 67 A. A. A. S., Washington: 391-397, 1 text-fig.
- BUCK, F. D., 1958: Cistelidae and Tenebrionidae (Coleoptera) from Rennell Island. - The Natural History of Rennell Island, British Solomon Islands, 2: 115-119, 1 pl.
- CHAMBERLIN, RALPH V., 1958: Millipeds and Centipeds from Rennell and Guadalcanal Islands. - Ibid.: 207-212, 4 text-figs.
- CLAY, THERESA, 1958: Mallophaga (Insecta). Part I. - Ibid.: 143-149.
- CLENCH, WILLIAM J., 1958: The Land and Freshwater Mollusca of Rennell Island, Solomon Islands. - Ibid.: 155-202, 3 text-figs., 4 pls.
- CLIFFORD, M. CARLETON & DANIEL SONENSHINE, 1962: A New Bat-Infesting *Ornithodoros* (Ixodoidea, Argasidae) of the Subgenus *Reticulninus* Schulze, 1941 from Rennell Island. - Ibid. 4: 25-29, 1 text-fig.
- FOREST, JACQUES, 1961: Pagurides de l'Afrique occidentale. - Atlantide Rep., 6: 203-250, 19 text-figs.
- GAUD, J., 1962: Sarcopitiformes plumicoles (Analgesoidea) parasites d'oiseaux de l'île Rennell. - The Natural History of Rennell Island, British Solomon Islands, 4: 31-50, 11 text-figs.
- HAIG, J., 1959: Porcellanid Crabs from West Africa. - Atlantide Rep. 5: 327-332.
- HANSEN, L., 1960: Some Macromycetes from Rennell and Alcester Islands. - The Natural History of Rennell Island, British Solomon Islands, 3: 127-132.
- HARDING, J. P., 1962: *Mungava munda* and Four Other New Species of Ostracod Crustaceans from Fish Stomachs. - Ibid. 4: 51-62, 53 text-figs.
- HJORTAA, H., 1961: "Galatheas" Wunderfisch - *Galatheathauma axell*. - Der Präparator, Bonn, 7, 1: 169-171, 2 text-figs.
- HOWARTH, T. G., 1962: The Rhopalocera of Rennell and Bellona Islands. - The Natural History of Rennell Island, British Solomon Islands, 4: 63-83, 10 pls.
- KIRKEGAARD, J. B., 1959: The Polychaeta of West Africa. Part I. Sedentary Species. - Atlantide Rep., 5: 7-117, 25 text-figs.
- KRAMP, P. L., 1959: Some New and Little-known Indo-Pacific Medusae. - Vidensk. Medd. Dansk Naturh. Foren., 121: 223-259, 19 text-figs.
- LAIRD, MARSHALL & ELISABETH LAIRD, 1959: Culicidae and Haematozoa from Bellona and Rennell. - The Natural History of Rennell Island, British Solomon Islands, 2: 213-234, 2 pls.
- LEMICHE, H., 1959: Molluscan phylogeny in the light of *Neopilina*. - Proc. XV. Intern. Congr. Zool. London: 380-381.
- LEMICHE, H., 1959: Protostomian interrelationship in the light of *Neopilina*. - Ibid.: 381-389.
- 1959: *Neopilina*. (Un molusco actual, con caracteres de tipo fosil, y su significado). - Revista Univ. Madrid, 8: 411-442, 12 text-figs.
- 1960: A Possible Central Place for *Stenothecoides* Resser, 1939 and *Cambridium* Horny, 1957 (Mollusca Monoplacophora) in Invertebrate Phylogeny. - Intern. Geol. Congr., XXI Session Norden, Part XXII: 92-101, 3 text-figs.
- 1962: The primitive mollusc that emerged from our oceans. - Scientia 56, 6: 1-8, 2 text-figs.
- & K. G. WINGSTRAND, 1959: The comparative anatomy of *Neopilina galathea* Lemche, 1957 (Mollusca Monoplacophora). - Proc. XV. Intern. Congr. Zool. London: 378-380.
- 1960: Classe des Monoplacophores. Monoplacophora (Odhner, 1940). - In: Traité de Zoologie, Anatomie, Systematique, Biologie, 5, 2: 1787-1821, 23 text-figs.
- LEVRING, T., 1960: A List of Marine Algae from Rennell Island. - The Natural History of Rennell Island, British Solomon Islands, 3: 121-125.
- MARSHALL, G. A. K., 1958: On Curculionidae (Coleoptera) from Rennell Island, Solomon Islands. - Ibid. 2: 121-126, 2 pls.
- NIELSEN, JØRGEN, 1961: Psittodoidea and Pleuronectoidea (Pisces, Heterosomata). - Atlantide Rep., 6: 101-127, 8 text-figs., 1 pl.
- NØRVANG, AKSEL, 1961: Schizaminidae, a New Family of Foraminifera. - Ibid.: 169-201, 2 text-figs., 4 pls.
- POULSEN, ERIK M., 1962: Ostracoda-Myodocopa. Part I. Cypridiniformes-Cypridinidae. - Dana Rep. No. 57: 1-414, 181 text-figs.
- POWELL, A. W. B., 1958: New Zealand Molluscan Systematics with Descriptions of New Species, Part 3. - Rec. Auck. Inst. Mus., 5, 1-2: 87-91, 4 text-figs., 1 pl.
- SCUDDER, G. G. E., 1958: Lygaeidae (Hemiptera) of Rennell and Bellona Islands. - The Natural History of Rennell Island, British Solomon Islands, 2: 135-142, 1 text-fig.
- STUBBINGS, H. G., 1961: Cirripedia Thoracica from Tropical West Africa. - Atlantide Rep., 6, 7-41, 11 text-figs.
- TANDAN, B. K., 1958: Mallophaga (Insecta). Part II. - A New Species of *Ibidoecus* Cummings, 1916. - The Natural History of Rennell Island, British Solomon Islands, 2, 151-154, 6 text-figs.
- TATTERSALL, OLIVE S., 1961: Mysidacea from the Coasts of Tropical West Africa. - Atlantide Rep., 6: 143-159, 3 text-figs.
- VIETTE, P. E. L., 1954: Une nouvelle espèce de Lépidoptère brachyptère de l'île Campbell. - Ent. Medd., København, 27, 1: 19-22, 1 text-fig.
- WILSON, EDWARD O., 1962: The Ants of Rennell and Bellona Islands. - The Natural History of Rennell Island, British Solomon Islands, 4: 13-23.
- WOLFF, TORBEN, 1959: La faune hadale ou faune des profondeurs supérieures à 6000-7000 metres. - Terre et Vie, 106, 2-3: 244-266, 4 text-figs., 1 pl.
- 1959: L'exploration des mers profondes. - La Nature, No. 3294, Oct. 1959: 425-431, 9 text-figs.
- 1960: The Hadal Community, an Introduction. - Deep-Sea Res., 6, 2: 95-124.
- 1960: Die tödliche Laterne. Was noch keines Menschen Auge sah: Lebewesen aus Tiefen bis 10.000 m. - Österreichischer Tierschutzkalender 1961, Wien: 55-58, 13 text-figs.
- 1961: The Deepest Recorded Fishes. - Nature, 190: 283.
- 1961: A Bathyal-abyssal Hermit Crab with a Calcified Asymmetrical Abdomen. - Ibid.: 931-932.
- YOSHIMOTO, C. M., J. L. GRESSITT & TORBEN WOLFF: Air-borne insects from the Galathea Expedition. - Pacific Insects, 4, 2: 269-291, 4 text-figs.

50.51531
G146
:6

GALATHEA REPORT

Volume 6

Scientific Results
of The Danish Deep-Sea Expedition
Round the World 1950-52



ISSUED BY THE GALATHEA COMMITTEE

PRESIDENT: H. R. H. PRINCE AXEL OF DENMARK

EDITORIAL COMMITTEE:

R. Spärck, D. Sc.

Professor of Zoology, Vice-President of the Committee

Sv. Greve, Captain, R. D. N.

Commander of the Galathea

F. Jensenius Madsen, D. Sc.

EDITOR:

Torben Wolff, Ph. D.

Deputy Leader of the Expedition

DANISH SCIENCE PRESS, LTD.

COPENHAGEN

Published by means of a grant from
STATENS ALMINDELIGE VIDENSKABSFOND
(The Danish State Research Foundation)

Printed by Nordlundes Bogtrykkeri
Copenhagen

Price: 120 Danish Kroner

GALATHEA REPORT

Volume 6



To the memory of

THE SCIENTIFIC LEADER
OF THE GALATHEA EXPEDITION

DR. ANTON FREDERIK BRUUN

LECTURER OF OCEANOLOGY
AT THE UNIVERSITY OF COPENHAGEN

*1901 †1961

THE SYSTEMATICS AND BIOLOGY OF BATHYAL AND ABYSSAL ISOPODA ASELLOTA

By TORBEN WOLFF
Zoological Museum, Copenhagen

CONTENTS

(* indicates species collected by the *Galathea* Expedition)

I. Preface	9	Key to the species	42
II. Introduction		<i>Janira maculosa</i> Leach and <i>J. hanseni</i> Menzies	42
Historical review	10	* <i>Janira operculata</i> n. sp.	44
Material and methods	13	<i>Janthura</i> n. gen.	45
List by station of <i>Galathea</i> Asellota (bathyal, abyssal, and hadal)	14	* <i>Janthura abyssicola</i> n. sp.	46
III. Terminology		Echinothambematidae	49
Morphological terms	15	Thambematidae	49
Oceanographical terms	16	Haploniscidae	49
IV. Systematic Part		Key to the genera	50
Classification of the Asellota	17	<i>Haploniscus</i> Richardson	50
Key to the subtribes	18	* <i>Haploniscus helgei</i> n. sp.	51
The significance of various characters in the taxonomy	18	* <i>Haploniscus kermadecensis</i> n. sp.	55
Stenetrioidea	21	<i>Haploniscus ingolfi</i> n. sp.	56
Key to the species of <i>Stenetrium</i>	22	<i>Haploniscus bicuspis tepidus</i> n. subsp.	58
<i>Stenetrium antillense</i> Hansen	25	<i>Haploniscus spinifer</i> Hansen	59
<i>Stenetrium siamense</i> Hansen	25	Munnidae	59
<i>Stenetrium occidentale</i> Hansen	25	"Pleurogoniidae"	60
<i>Stenetrium serratum</i> Hansen	25	<i>Astrurus</i> Beddard	61
* <i>Stenetrium abyssale</i> n. sp.	25	<i>Caecimunna</i> Richardson	61
Paraselloidea		<i>Notoxenoides</i> Menzies	62
Classification	29	Key to the genera	62
Key to the families	31	The <i>Dendrotion-Antias-Abyssianira</i> complex.	63
Janiridae	33	<i>Xostylus</i> Menzies	63
"Janirellidae"	33	<i>Urias</i> Richardson	64
<i>Iolella vilhelminae</i> Stephensen	33	<i>Haplomunna</i> Richardson	61, 64
<i>Iolella spinosissima</i> Stephensen	33	<i>Mesosignum</i> Menzies	64
<i>Rhacura pulchra</i> Richardson	34	Dendrotionidae	64
<i>Ianthopsis nasicornis</i> Vanhöffen	34	Key to the genera	65
<i>Acanthaspidia typhlops</i> (G. O. Sars)	34	<i>Munella</i> Bonnier	64
<i>Janirella</i> Bonnier	34	<i>Pleurocope</i> Walker	64
"Acanthaspidiidae"	35	<i>Dendrotion</i> G. O. Sars	65
"Microparasellidae"	35	Key to the species	65
<i>Ectias turqueti</i> Richardson	37	<i>Acanthomunna</i> Beddard	61, 65
Comments on separate genera of Janiridae	39	<i>Acanthomunna hystrix</i> (Hansen)	66
List of the genera of Janiridae	39	<i>Dendromunna</i> Menzies	66
<i>Jaera</i> Leach	40	Key to the species	66
<i>Jaera nordmanni</i> (Rathke)	40	* <i>Dendromunna mirabile</i> n. sp.	66
<i>Janira</i> Leach	41	Antiasidae	69
		Key to the genera	70
		<i>Abyssianira</i> Menzies	70
		<i>Kuphomunna</i> Barnard	61, 70
		<i>Antias</i> Richardson	70
		Key to the species	70

Ischnomesidae	71	<i>Storthyngura</i> Vanhöffen	118
Key to the genera	73	Key to the species	120
<i>Ischnomesus</i> Richardson	73	<i>Storthyngura novae-zelandiae</i> (Beddard)	122
Key to the species	73	<i>Storthyngura atlantica</i> (Beddard)	125
<i>Ischnomesus wolffi</i> Menzies	75	<i>Storthyngura fragilis</i> (Beddard)	126
* <i>Ischnomesus birsteini</i> n. sp.	75	<i>Storthyngura challengeri</i> n. sp.	128
* <i>Ischnomesus</i> sp.	77	<i>Storthyngura gordonae</i> n. sp.	130
* <i>Ischnomesus anacanthus</i> n. sp.	78	<i>Storthyngura</i> sp.	132
* <i>Ischnomesus bruuni</i> Wolff	79	<i>Storthyngura</i> sp. juv.	132
* <i>Ischnomesus roseus</i> n. sp.	80	<i>Storthyngura chelata</i> Birstein	133
* <i>Ischnomesus planus</i> n. sp.	82	<i>Storthyngura bicornis</i> Birstein	133
<i>Bactromesus</i> n. gen.	83	<i>Storthyngura pulchra</i> (Hansen)	134
Key to the species	83	* <i>Storthyngura p. pulchra</i> (Hansen)	138
<i>Stylomesus</i> Wolff	83	<i>Storthyngura p. caribbea</i> (Benedict)	138
Key to the species	83	* <i>Storthyngura p. kermadecensis</i> n. subsp.	139
<i>Stylomesus inermis</i> (Vanhöffen)	84	* <i>Storthyngura furcata</i> Wolff	139
<i>Stylomesus pacificus</i> Birstein	84	* <i>Storthyngura abyssalis</i> n. sp.	139
<i>Gomphomesus</i> n. gen.	84	* <i>Storthyngura serrata</i> n. sp.	141
<i>Helomesus</i> n. gen.	84	<i>Eurycope</i> G. O. Sars	143
Key to the species	84	Key to the species	144
<i>Heteromesus</i> Richardson	84	<i>Eurycope sarsi</i> Beddard	147
Key to the species	85	* <i>Eurycope complanata</i> Bonnier	147
<i>Haplomesus</i> Richardson	86	* <i>Eurycope gibberifrons</i> n. sp.	147
Key to the species	86	<i>Munnopsurus</i> Richardson	149
<i>Haplomesus thomsoni</i> (Beddard)	87	Key to the species	150
<i>Haplomesus modestus</i> Hansen	87	<i>Munnopsurus longipes</i> (Tattersall)	151
<i>Mixomesus</i> n. gen.	88	<i>Munnopsurus mimus</i> Barnard	153
* <i>Mixomesus pellucidus</i> n. sp.	89	<i>Munnopsurus australis</i> (Vanhöffen)	153
Macrostylidae	90	<i>Munnopsurus laevis</i> (Richardson)	154
<i>Macrostylis</i> G. O. Sars	91	<i>Munnopsurus giganteus ochotensis</i> Gurjanova	154
<i>Macrostylis subinermis</i> Hansen	91	<i>Munnopsurus minutus</i> Gurjanova	154
<i>Macrostylis magnifica</i> n. sp.	91	<i>Munneurycope</i> Stephensen	154
* <i>Macrostylis hadalis</i> Wolff	92	Key to the species	156
<i>Macrostylis spinifera</i> Hansen	92	<i>Munneurycope murrayi</i> (Walker)	157
Pseudomesidae	93	<i>Munneurycope incisa</i> (Gurjanova)	161
Ilyarachnidae	93	* <i>Munneurycope harrietae</i> n. sp.	161
Key to the genera	93	<i>Munneurycope nodifrons</i> (Hansen)	164
<i>Ilyarachna</i> G. O. Sars	94	* <i>Munneurycope menziesi</i> n. sp.	164
Key to the species	94	<i>Munneurycope elongata</i> n. sp.	167
<i>Ilyarachna multispinosa</i> Menzies	97	Bathyopsurinae nov. subfam.	170
<i>Ilyarachna acarina</i> Menzies & Barnard	97	Key to the genera and species	171
<i>Ilyarachna longicornis</i> (G. O. Sars)	97	<i>Bathyopsurus</i> Nordenstam	171
<i>Ilyarachna thori</i> n. sp.	97	* <i>Bathyopsurus nybelini</i> Nordenstam	171
<i>Ilyarachna dubia</i> Hansen	100	<i>Bathyopsurus abyssicolus</i> (Beddard)	176
<i>Ilyarachna bicornis</i> Hansen	101	<i>Paropsurus</i> n. gen.	176
<i>Ilyarachna antarctica</i> Vanhöffen	102	* <i>Paropsurus giganteus</i> n. sp.	177
<i>Ilyarachna nordenstami</i> n. sp.	102	<i>Paropsurus pellucidus</i> (Beddard)	181
* <i>Ilyarachna kermadecensis</i> n. sp.	102	Munnopsidae	183
* <i>Ilyarachna aspidophora</i> n. sp.	106	Key to the genera and species	186
Eurycopidae	108	<i>Paramunnopsis</i> Hansen	187
Key to the subfamilies	109	<i>Munnopsis</i> M. Sars	187
Acanthocopinae nov. subfam.	109	<i>Munnopsis beddardi</i> (Tattersall)	188
<i>Acanthocope</i> Beddard	110	<i>Munnopsis eximius</i> (Hansen)	188
Key to the species	110	<i>Munnopsis typica</i> M. Sars	188
<i>Acanthocope spinicauda</i> Beddard	111	<i>Munnopsis australis</i> Beddard	190
<i>Acanthocope acutispina</i> Beddard	112	<i>Munnopsis latifrons</i> Beddard	191
* <i>Acanthocope galathea</i> n. sp.	113	* <i>Munnopsis longiremis</i> Richardson	195
Syneurycopinae nov. subfam.	116	<i>Munnopsis gracilis</i> Beddard	199
Key to the species	117	* <i>Munnopsis bathyalis</i> n. sp.	200
Eurycopinae nov. subfam.	117	* <i>Munnopsis mandibularis</i> n. sp.	203
Key to the genera	117	List of generic synonyms of Stenetrioidea and	
<i>Lipomera</i> Tattersall	118	Paraselloidea	206

V. General Part	
A. Postmarsupial development	206
<i>Jaera albifrons albifrons</i>	206
<i>Pseudasellus nichollsi</i>	207
Other species	
<i>Haploniscus bicuspis bicuspis</i>	208
<i>Haploniscus spinifer</i>	210
<i>Haploniscus antarcticus</i>	211
<i>Haploniscus helgei</i>	212
<i>Haploniscus tridens</i> and <i>percavix</i>	212
<i>Janirella vema</i> , <i>bifida</i> and <i>magnifrons</i>	212
<i>Echinothambema ophiuroides</i>	212
<i>Ilyarachna multispinosa</i>	212
Conclusions	212
B. Reproduction	214
Number of preparatory stages	214
One preparatory stage before sexual maturity	214
Two preparatory stages before sexual maturity	215
The developing oostegites	
Length	218
Number	219
The incubatory stage	219
Possible reasons for deficit of ovigerous females	219
The marsupium	221
Eggs and embryos	221
Males – sex ratio	222
C. Hermaphroditism in Isopoda	223
a. Previous studies	223
1. Flabellifera	223
2. Epicaridea	223
3. Oniscoidea	224
4. Asellota	225
b. Own investigations	226
<i>Haploniscus bicuspis bicuspis</i>	226
<i>Haploniscus helgei</i>	226
c. Conclusions	228
D. Some characteristics of deep-sea Asellota	229
Correlation between size and depth	229
Previous evidence	229
The genera of the Asellota	229
Elongation of legs	238
Eyes	238
E. The food of some deep-sea Asellota	238
a. Previous studies	238
b. Own investigations	239
c. Discussion	243
F. Habitat	245
1. Benthic occurrence	245
Substratum	245
2. Pelagic occurrence	247
G. Distribution of the marine Asellota	248
1. Bathymetrical distribution	270
a. Species	
Depth	270
Temperature	272
Depth-temperature	273
b. Genera	
Depth	275
Temperature	280
Depth-temperature	281
c. Families	282
d. Vertical zonation in the abyssal zone	283
2. Regional distribution	286
a. Species	287
b. Genera	290
c. Families	294
d. Conclusions	296
e. Remarks on the regional distribution of the abyssal fauna	300
H. Summary	303
J. References	307
K. Index	315

I. PREFACE

The present work arises from a study of bathyal and abyssal¹ Isopoda Asellota, collected mainly during the Danish *Galathea* Deep-Sea Expedition Round the World 1950-52 at depths between about 200 and 6000 m. Since the Asellota is by far the most abundant suborder of Isopoda in the deep-sea, the present investigation was confined to this group. The hadal¹ isopods from the *Galathea* have been dealt with previously (WOLFF 1956 a), and it is intended to publish at a later date in the *Galathea Report* results obtained from an examination of the remaining suborders from bathyal and abyssal depths.

Much of the systematic literature on deep-sea isopods – both old and new – is inadequate, owing to descriptions being too short and incomplete, and lack of, or far too few illustrations. In order to

1. Definitions of these terms are given on p. 16.

evaluate systematic relationships and to arrive at the greatest possible degree of certainty when dealing with e.g. the zoogeography of the species of a genus, I have deemed it necessary, in many cases to consult previously examined material from other sources. In this respect the Copenhagen Museum houses many rich collections from earlier Danish expeditions, principally, that of the *Ingolf*. In addition, I have borrowed material from several other museums and have on three occasions visited the British Museum (Natural History), to study types from the *Challenger* Expedition. It was found necessary to re-describe and revise much of this borrowed material, and even to establish a number of new species or subspecies.

The study is essentially morphological and taxonomical. For obvious reasons, deep-sea collec-

tions are generally restricted in number of specimens. However, as far as was possible with the material available, other aspects have also been considered.

Since very little is known on the general biology and ecology of asellotes (those from the deep-sea in particular) the *Galathea* and additional material was utilized for investigations on development, propagation, hermaphroditism, food content, etc.

Up to the present time the Asellota have not been used for purposes of monographic treatment¹. A general survey of the distribution of the many species of the suborder – regional as well as bathymetrical – was therefore not available. With regard to the zoogeography of material dealt with in this paper, I thus considered it appropriate to gather data from all available literature on distribution, depth records, etc.; this providing at the same time, a basis for comparison between the distribution of littoral, sublittoral, bathyal, abyssal and hadal asellotes. Unfortunately, our knowledge of the bathyal and abyssal isopods is still so fragmentary that their contribution to a general conception of the zoogeography of the deep-sea is very limited.

Acknowledgments

In presenting this work it is my privilege to thank all those who have been of assistance to me. In the first place, I wish to express my gratitude to colleagues from the *Galathea*, in particular the leader, the late Dr. ANTON F. BRUUN, for friendly and fruitful discussions both on board and during subsequent treatment of the material.

1. NIERSTRASZ's compilation (1941) deals with Indo-Pacific species only. In addition, several species are not mentioned and the paper contains many errors in records of distribution and depth.

My thanks are also due to Dr. HENNING LEMCHE (the Zoological Museum), for advice on problems of nomenclature, to Professor K. G. WINGSTRAND (the Anatomical Institute) for valuable help in microscopical sectioning and interpretation of the gonads of *Haploniscus*, to Mr. TYGE CHRISTENSEN (the Botanical Museum), for advice on questions of terminology, and on determination of plant material in intestine contents, and to Mr. JENS SMED (the Hydrographic Office), for assistance with references to temperature records.

A considerable number of colleagues in foreign museums and other institutions have been very helpful in placing material at my disposal. The aid of the following is particularly appreciated: Dr. CHARLOTTE HOLMQUIST (Stockholm), Professor O. NYBELIN (Gothenburg), Mr. N. KNABEN (Oslo), Dr. ISABELLA GORDON and Mr. R. W. INGLE (London), Dr. H.-E. GRUNER (Berlin), Dr. A. CAPART (Brussels), Professor J. A. BIRSTEIN (Moscow), Dr. ELIZABETH DEICHMANN (Harvard) and Drs. FENNER A. CHACE and T. E. BOWMAN (Washington). My thanks are also due to Dr. BOWMAN, Dr. GORDON and Mr. INGLE for providing sketches of a few type specimens which could not be borrowed, and to the latter two for their assistance during my three visits to the British Museum (Natural History) in London. I am also grateful to Dr. R. J. MENZIES (Los Angeles) for permission to go through the proof of his paper on the systematics of the rich collections of the *Vema* Expeditions in the Atlantic Ocean, to Professor CLAUDE LÉVI (Strasbourg) who identified sponge material in gut contents of various isopods, and to Professor ELLSWORTH C. DOUGHERTY (Berkeley) for suggestions regarding discussions on terminology.

Finally, I thank Mrs. M. GOODFELLOW for careful revision of the English text and my wife for invaluable help with proof-reading, etc.

II. INTRODUCTION

HISTORICAL REVIEW

The main contributions to our knowledge of the bathyal, abyssal and (recently) hadal asellote isopods come from the reports of the great oceanographical expeditions. But important work has also been published as a result of dredgings undertaken on the continental slope, particularly in the vicinity of universities and marine institutes in Western and Northern Europe.

The first reports on asellotes from outside the shelf area date back almost a hundred years, when the great Norwegian carcinologist G. O. SARS described and recorded several new or previously described species from the upper slope area in the outer part of Oslo (Christiania) and Hardanger Fjord and off Lofoten (SARS 1869, 1870, 1872, and 1879). In 1885 G. O. SARS published his large paper on the Crustacea from the Norwegian North Atlantic Expedition 1876-1878 and described the first

known, truly bathyal asellotes. The following year came BEDDARD's important work on the Isopoda of the British *Challenger* Expedition 1872-1876, which dealt with 34 asellotes, 32 of which were new to science (two species from less than 200 m, fourteen from 200-2000 m and sixteen from 2000 to 5011 m, the latter being for the next seventy years the greatest known depth for any asellote!). BEDDARD gave also a survey of the zoogeography of deep-sea isopods known at that time and discussed their special peculiarities.

For obvious reasons, the German Plankton-Expedition obtained only a few Asellota, the great majority being benthic. In connexion with the description of these species HANSEN (1895) pointed out the difference between *Asellus*, *Stenetrium* and the remaining asellotes, and supported CLAUS' division between Isopoda and Tanaidacea. BONNIER (1896) reported on the Cumacea, Isopoda and Amphipoda collected by the *Caudan* in the Gulf of Gascony and described eight species of asellotes, all from 950 m depth.

In 1899 appeared G. O. SARS' volume on the isopods in his outstanding "Crustacea of Norway", in which the diagnoses of all hitherto described and several new genera and species from Northern Europe are given together with observations on swimming methods, etc. The monograph includes 21 species from the shelf and as many again from the lower sublittoral and upper bathyal zones. OHLIN (1901) described a single abyssal species from the Arctic Ocean and supplied temperature records for this and the other, sublittoral species dealt with in his report. TATTERSALL (1905) recorded nineteen bathyal species from off Ireland, seven of which were new, taken by the Irish fishing vessel, *Helga*.

The bathyal and abyssal isopods collected by the American *Albatross* Expeditions, mainly in North American waters, were described by several authors (HANSEN 1897, BENEDICT (in RICHARDSON 1901), and mainly by RICHARDSON (1905, 1908a, 1908b, 1909, 1911b, and 1912b)); out of twenty asellotes, nine bathyal and ten abyssal species were new. RICHARDSON (1911a) also gave descriptions (but no figures) of three new bathyal and two new abyssal species from the East Atlantic, taken by the French *Travailleur* and *Talisman* (1880-1883), STEBBING (1913) and STEPHENSEN (1915) contributed with descriptions of a few abyssal or bathyal asellotes from the expeditions of the British *Porcupine* (1869-1870) and the Danish *Thor* (1904-1910).

In 1914 VANHÖFFEN published his large paper on

the isopods of the German South Polar Expedition of the *Gauss* (1901-1903). This is the first report on deep-sea isopods in the Antarctic area (apart from a few from the *Challenger*) and contains descriptions of 63 asellote species. Of the twenty shelf species twelve were previously known, although this applied to only five of thirty slope and one of ten abyssal species. Unfortunately, the entirety of the descriptions and the accuracy of the figures are not always satisfactory in this important work; it also contained a survey of the zoogeography of the Antarctic isopods. In 1914 came the first of K. H. BARNARD's papers on South African crustaceans in which bathyal asellotes are also mentioned; this paper and its successor (1920), contain descriptions of fourteen new species from the continental slope.

Probably the most important paper on deep-sea isopods is HANSEN's report (1916) on the collections from the Danish *Ingolf* and *Thor* Expeditions to the North Atlantic (in 1895-1896 and 1904-1910, respectively). His revision of the classification of the Asellota is that generally accepted today. A total of 105 species belonging to this group, are excellently described, their relationship discussed, and the bottom temperatures recorded. Seventeen of them are from the shelf, 36 from bathyal depths, 27 from abyssal depths and 5 ranged from bathyal into abyssal depths. There are 10 new genera and 61 new species, in spite of the fact that the Eastern North Atlantic was then - and still is - the most thoroughly investigated area. Finally, the paper includes a survey of the regional and bathymetrical distribution.

MONOD (1926) recorded eight slope species from the Antarctic. Although NORDENSTAM's large paper (1933) on the isopods from the Swedish Antarctic Expedition in 1901-1903 deals with only a few species from depths exceeding 200 m, it is important insofar as it revises the systematics of isopods from this area. GURJANOVA gave (1933b) a useful survey of all marine isopods known from Arctic and Subarctic waters, based partly on the results of a number of (mostly Russian) expeditions or collectors, partly on the literature. The paper includes short diagnoses of the genera, keys to genera and species, and a chapter on the zoogeography of the Arctic Sea and adjacent areas. It should be noted that the majority of calculations of the original records from fathoms to metres are incorrect.¹ In

1. This is probably in part due to the fact that a Danish fathom (as e. g. used by HANSEN and SARS) is notably longer than an English fathom, thus causing considerable error where depths of several thousand metres are concerned.

1933a GURJANOVA described four bathyal asellotes from the Bering Sea, the Sea of Ochotsk and the Sea of Japan, and in 1936 she reviewed the zoogeography of these and all other species from that area. STEPHENSEN (1936) mentioned a few asellotes from bathyal depths from the Danish *Godthaab* Expedition to West Greenland in 1928.

The only contribution to our knowledge of the ecology of soft bottom isopods (almost exclusively asellotes) is HULT's very important paper (1941). A total of 28 asellotes was secured in the Skagerrak and adjacent deep fjords, 23 from depths exceeding 200 m. Besides the regional and bathymetrical distribution of the species, their relation to temperature, salinity, oxygen, type of bottom, etc. is discussed and the dredging methods evaluated.

NIERSTRASZ (1941) worked up some of the isopods from the Dutch *Siboga* Expedition and gave fairly complete lists of all Indo-Pacific marine isopods (excl. of Flabellifera and Epicaridea), including distribution and depth records; unfortunately, the deeper depth records have all been calculated erroneously from fathoms to metres. Probably due to the untimely death of Professor NIERSTRASZ 1937, the *Siboga* material of Asellota was, unfortunately, not treated.

GURJANOVA (1946) recorded two shelf species, six slope species and two abyssal species, collected in the Arctic Ocean by the *Sadko* in 1935 and 1937-38 and by the *Sedov* in 1937-38. Gorbunov (1946) listed 26 previously described asellotes from the slope and abyssal depths, procured by the same vessels, and discussed the composition of the fauna.

In 1955 NORDENSTAM described the first isopod known from depths exceeding 5000 m, this being a new genus and species of an asellote dredged by the Swedish Deep Sea Expedition in 1948 in and close to the Puerto Rico Trench at depths between 5500 and 7900 m. NORDENSTAM also included a somewhat incomplete list of isopods recorded from depths greater than 3000 m. MENZIES (1956a, 1956b) described two new asellotes from 1250 m and five from 5100 m in the West Indies, all collected by the American *Vema* Expedition; in both papers MENZIES also supplied information on the nutrition of the species.

WOLFF (1956a) dealt with the hadal representatives of the collections of the Danish *Galathea* Expedition 1950-1952, recording eleven asellotes from depths exceeding 6000 m and down to 9790 m¹ in

1. This depth was later revised to 9820-10.000 m.

the Philippine Trench, the greatest depth so far from where isopods have been described¹. In addition, eleven other deep-sea species were revised. The regional and vertical distribution of the genera with hadal species was recorded in detail, and the relationship, possible origin, special peculiarities, and nutrition of the hadal species discussed; the first evidence of possible gigantism in deep-sea isopods was given.

In 1957 BIRSTEIN published the first descriptions of asellotes from the Russian *Vitjaz* Expeditions in the N.W. Pacific, including seven new species and subspecies of the genus *Storothyngura*, from depths between 5670 and 8430 m. He also discussed the distribution of this genus, the origin of the abyssal and hadal faunas, and the gigantism of the species, ascribing this phenomenon to the effect of hydrostatic pressure on the metabolism. Later (1960), BIRSTEIN treated the Ischnomesidae, recording eleven new and one previously described species and subspecies from the deeper abyssal and the hadal zones. Moreover, the distribution of the genera was evaluated and the bipolar and amphiboreal distribution of the deep-sea fauna discussed. A new genus and species of an abyssal isopod was described and its systematic position discussed in detail by BIRSTEIN (1961).

Finally, MENZIES & TINKER (1960) described a single abyssal species of *Haploniscus* from the East Pacific and a short time ago MENZIES (1962b)² published his large work on the systematics of the abyssal isopod fauna of the Atlantic Ocean. The paper was based upon the remarkable results of 84 trawlings, carried out primarily by the *Vema*, and initiated by the Lamont Geological Observatory, New York. A total of 176, mainly abyssal species from the North and South Atlantic, including the Arctic Ocean and Caribbean Sea, are diagnosed; the majority, 159 species, are Asellota. Fifty-nine of the latter have been previously described; besides a short diagnosis, one or more of the original figures have

1. BELJAEV *et al.* (1958) recorded "isopods" from 10.415-10.687 m in the Tonga Trench and WOLFF (1960) "isopods" from 9995-10.002 m in the Kermadec Trench and 8980-9043 m in the New Britain Trench, in all three cases collected by the Soviet research vessel "*Vitjaz*".

2. Since the publication of this monograph was already anticipated by the Columbia University Press by June 1961, Dr. MENZIES kindly allowed me to go through the final proof on receiving it. Unfortunately, the publication was considerably postponed with the result that I have been unable to make full use of the important results brought about by that paper.

been traced. No less than 100 asellote species and 7 genera are new (1 sublittoral, 10 bathyal, and 89 abyssal species). This more than doubles the number of deep-sea species from the Atlantic. Keys are given to most of the species and genera dealt with. It is intended to publish the monograph in two parts, the second dealing with the zoogeographical, ecological, and phylogenetical relationships of the collections.

MATERIAL AND METHODS

The *Galathea* material of isopods was collected with various kinds of trawls, described by BRUUN (1959). The types of gear are recorded in the following paragraph. Both when using large otter trawls and sledge trawls, a canvas bag was attached to the cod end of the net in order to retain samples of ooze and the more minute animals. The bottom material was subsequently washed through sieves with a mesh diameter of 1 mm (occasionally $\frac{1}{2}$ mm). However, when making a comparison between the average size of isopods obtained by the *Galathea* and the *Vema* (MENZIES 1962 b), it seems that the efficacy of this method may be questionable; the number of specimens measuring less than 2-3 mm being much smaller in the *Galathea* than in the *Vema* material. The latter was obtained by sorting the entire sample¹, which usually consisted of a quart of sediment and animals, under a binocular microscope. The samples were collected by a small trawl or dredge; the orifice was 1 m wide and 10 cm high and the bag was 3 m long and had a mesh diameter of only 0.5 mm. See WOLFF (1961, p. 155) on a discussion of the very variable results obtained with different types of gear within a limited area of the abyssal zone.

A total of 23 species of bathyal and abyssal asellotes (15 different genera) were collected by the *Galathea*. There were 83 specimens in all. No less than 15 species were represented by a single specimen only.

The isopods were sorted onboard the ship and preserved in 70 % alcohol as soon as I had taken notes of colour, etc.

Dissection during the working up was done under microscope while the specimens were immersed in ethylene glycol, which has a high viscosity. Small specimens were dissected on a microscope slide. As far as was possible, dissected mouthparts etc. were

1. This method was highly recommended by HULT (1941, p. 6).

preserved in minute vials of alcohol instead of being mounted on slides, as the latter method prohibits a thorough study (especially of the mandibles), at a later date. Only the dissected parts of very small species were mounted (in Faure-Berlese medium, with a slight red staining of Lignin Pink to make the transparent parts visible).

Measurements were made with an ocular micrometer. Body lengths were measured from the anterior end of the frontal area (or projection, if present), to the posterior tip of the pleotelson, thus excluding the uropods. If spines reached beyond the posterior end of the pleotelson, the length was measured to the tips of the spines; in doubtful cases it is stated exactly what the measurement covers.

The descriptions are rather extensive, since I personally found detailed descriptions to be of great help during the study – and this not only for purely taxonomic reasons. Reasonably detailed descriptions are especially important in groups such as the asellotes, in which a satisfactory systematic division has not yet been reached. Diagnoses are applied in all cases where they were considered adequate. Attention has been paid to avoid repetition in inferior taxons, of characters which have already been covered in the diagnoses of the superior taxon(s).

Keys had been prepared to all genera dealt with prior to receiving the proof of Dr. MENZIES' paper (cf. above), which includes his keys to the majority of the same genera. The many new species he described have been included in my keys. These are worked out in such a way that each couplet, whenever practicable, is based on more than one character (preferably three) to make the use of the key more reliable. In addition, the use of characters such as legs, uropods, etc. have been avoided to the greatest possible extent, since these are very often lost in deep-sea material. Characters requiring dissection of mouthparts have only been applied when absolutely necessary. In several cases insufficient descriptions and illustrations, or the fragmentary condition of the specimen described, made it very difficult to incorporate such species in the key.

As in the paper on the hadal species (WOLFF 1956a), lectotypes have been selected in several of those species studied from the *Ingolf*, the *Challenger* and the American *Albatross* Expeditions.¹

1. Throughout the *Vema* paper MENZIES (1962 b) has selected type localities not only for the new species but also for all previously described Atlantic abyssal species mentioned. However, none of the old carcinologists (SARS, RICHARDSON, HANSEN, etc.) selected holo- and allotypes, and today

The illustrations consist of drawings and photographs. The former were made in pencil with the aid of a camera lucida and I later inked them in. A few of the drawings have been "shadowed" by the artist, Mr. POUL H. WINTHER. All drawings are based on the holotype when not otherwise stated. The photographs were included in cases where they were considered the best means of illustration and where it was found unnecessary to prepare time-consuming drawings of certain species. They were almost all taken by Mr. H. V. CHRISTENSEN.

Methods used for examining contents of intestines are explained on p. 239.

Records of bottom temperatures were obtained in the same way as for hadal isopods and tanaids (WOLFF 1956a, p. 86; 1956b, p. 188).

**LIST BY STATION
OF GALATHEA ASELOTA**
(Bathyal, abyssal, and hadal¹)

Abbreviation of gear used (cf. BRUUN 1959, p. 22): HOT: herring otter trawl; SOT: shrimp otter trawl; ST 300 and ST 600: sledge (Agassiz, Sigsbee) trawl, 3 m and 6 m wide; PGI 0.2: Petersen grab (bottom sampler) covering 0.2 sq. m.

Stations

52. Bay of Guinea (1°42'N, 7°51'E), 2550 m, c. 3.0°C., 30. XI. 1950, SOT
Janthura abyssicola n. gen., n. sp.; 1 ♂
241. Off Kenya (4°00'S, 41°27'E), 1510 m, c. 3.2°C., 15. III. 1951, HOT
Munnopsis mandibularis n. sp.; 1 ♀
435. Philippine Trench (10°20'N, 126°41'E), 9820-10,000 m, 2.6°C., 7. VIII. 1951, ST 300
Macrostylis galathea Wolff; 2 ♀, 2 ♂
490. Bali Sea (5°25'S, 117°03'E), 545-570 m, c. 6.5°C., 14. IX. 1951, ST 300
Munnopsis bathyalis n. sp.; 2 ♀, 4 ♂

it is impossible to designate a type locality, except when the material on which the description was based originates from one locality only – or when a lectotype is selected from the original material. Since MENZIES has not seen any of the original material, he has been unable to select lectotypes. Moreover, while in the "Copenhagen Decisions" (HEMMING 1953) it was recommended (p. 27) that it should "be open to that taxonomist, acting as first reviser, to specify a restricted portion of the region or area cited by the original author to be the locality for the nominal species concerned" (provided that a holo- or lectotype was available), this paragraph was not entered in the Rules (1961 – as accepted by the Congress in London 1958) and is, therefore, now invalid.

1. The hadal species were treated in WOLFF 1956a.

496. Banda Trench (5°36'S, 131°06'E), 7270 m, 3.6°C., 23. IX. 1951, PGI 0.2
Macrostylis hadalis Wolff; 1 ♀, 1 ♂
554. Great Australian Bight (37°28'S, 138°55'E), 1320-1340 m, c. 3.5°C., 5. XII. 1951, ST 300
Haploniscus helgei n. sp.; 9 ♀, 5 ♂
575. Tasman Sea (40°11'S, 163°35'E), 3710 m, c. 1.1°C., 19. XII. 1951, SOT
Ischnomesus anacanthus n. sp.; 1 ♀
Ischnomesus sp.; 1 ♀
601. Tasman Sea (45°51'S, 164°32'E), 4400 m, c. 1.1°C., 14. I. 1952, HOT
Storhyngura abyssalis n. sp.; 2 ♀
Munneurycope harrietae n. sp.; 2 ♂
Bathyopsurus nybelini Nordenstam; 1 ♀
Paropsurus giganteus n. gen., n. sp.; 1 ♀
602. Tasman Sea (43°58'S, 165°24'E), 4510 m, c. 1.1°C., 15. I. 1952, ST 300
Stenetrium abyssale n. sp.; 1 ♀
626. Tasman Sea (42°10'S, 170°10'E), 610 m, c. 7.6°C., 20. I. 1952, HOT
Mixomesus pellucidus n. gen., n. sp.; 1 ♀
639. Off E. New Zealand (37°31'S, 177°08'E), 213 m, c. 14.7°C., 26. I. 1952, PGI 0.2
Ilyarachna aspidophora n. sp.; 1 ♀
Eurycope gibberifrons n. sp.; 1 ♂
650. Kermadec Trench (32°20'S, 176°54'W), 6620-6730 m, 1.3°C., 15. II. 1952, ST 600
Storhyngura benti Wolff; 1 ♀
Storhyngura furcata Wolff; 1 ♀
Storhyngura pulchra kermadecensis n. subsp., 5 ♀, 3 ♂
651. Kermadec Trench (32°10'S, 177°14'W), 6960-7000 m, 1.3°C., 16. II. 1952, HOT
Ischnomesus bruuni Wolff; 1 ♂
Ischnomesus spärcki Wolff; 1 ♂
Ilyarachna kermadecensis n. sp.; 3 ♀, 2 ♂
Storhyngura benti Wolff; 18-21 ♀, 2-4 ♂
Eurycope galathea Wolff; 1 ♂, 1 spm.
Eurycope madseni Wolff; 1 ♂
Munneurycope menziesi n. sp.; 1 ♀
654. Kermadec Trench (32°10'S, 175°54'W), 5850-5900 m, 1.2°C., 18. II. 1952, HOT
Storhyngura benti Wolff; 1 ♀
Storhyngura furcata Wolff; 3 ♀
Bathyopsurus nybelini Nordenstam, 1 ♂
658. Kermadec Trench (35°51'S, 178°31'W), 6660-6770 m, 1.3°C., 20. II. 1952, ST 600
Ischnomesus spärcki Wolff; 1 ♂, 1 spm.
Ilyarachna kermadecensis n. sp.; 2 ♂
Storhyngura benti Wolff; 2 ♀, 3 ♂
Storhyngura furcata Wolff; 2 ♀, 1 ♂