
偏微分方程导论

（第2版）

[美] 瓦尔特·A. 施特劳斯 著



WILEY

世界图书出版公司

需要全本请访问 www.ertongbook.com

PARTIAL DIFFERENTIAL EQUATIONS

AN INTRODUCTION

WALTER A. STRAUSS
Brown University

BICENTENNIAL

John Wiley & Sons, Ltd

图书在版编目（CIP）数据

偏微分方程导论：第2版 = Partial Differential Equations: An Introduction, 2nd Edition：英文 /（美）W. A. 施特劳斯（Walter A. Strauss）著 . — 影印本 . —北京：世界图书出版有限公司北京分公司，2019.6

ISBN 978-7-5192-6375-1

Ⅰ . ①偏… Ⅱ . ① W… Ⅲ . ①偏微分方程—英文 Ⅳ . ① O175.2

中国版本图书馆 CIP 数据核字（2019）第 113065 号

中文书名 偏微分方程导论（第2版）
英文书名 Partial Differential Equations: An Introduction *Second Edition*
著　　者 [美] 瓦尔特·A. 施特劳斯（Walter A. Strauss）
责任编辑 刘　慧　高　蓉

出版发行 世界图书出版有限公司北京分公司
地　　址 北京市东城区朝内大街 137 号
邮　　编 100010
电　　话 010-64038355（发行）　64033507（总编室）
网　　址 http://www.wpcbj.com.cn
邮　　箱 wpcbjst@vip.163.com
销　　售 新华书店
印　　刷 北京建宏印
开　　本 710 mm × 1(
印　　张 28.75
字　　数 458 千字
版　　次 2019 年 7 月
印　　次 2019 年 7 月
版权登记 01-2019-1
国际书号 ISBN 978-7
定　　价 99.00 元

版权所有　翻印必究

（如发现印装质量问题，请与所购图书销售部门联系调换）

PARTIAL DIFFERENTIAL EQUATIONS

AN INTRODUCTION



THE WILEY BICENTENNIAL–KNOWLEDGE FOR GENERATIONS

Each generation has its unique needs and aspirations. When Charles Wiley first opened his small printing shop in lower Manhattan in 1807, it was a generation of boundless potential searching for an identity. And we were there, helping to define a new American literary tradition. Over half a century later, in the midst of the Second Industrial Revolution, it was a generation focused on building the future. Once again, we were there, supplying the critical scientific, technical, and engineering knowledge that helped frame the world. Throughout the 20th Century, and into the new millennium, nations began to reach out beyond their own borders and a new international community was born. Wiley was there, expanding its operations around the world to enable a global exchange of ideas, opinions, and know-how.

For 200 years, Wiley has been an integral part of each generation's journey, enabling the flow of information and understanding necessary to meet their needs and fulfill their aspirations. Today, bold new technologies are changing the way we live and learn. Wiley will be there, providing you the must-have knowledge you need to imagine new worlds, new possibilities, and new opportunities.

Generations come and go, but you can always count on Wiley to provide you the knowledge you need, when and where you need it!

William J. Pesce

Peter Booth Wiley

WILLIAM J. PESCE
PRESIDENT AND CHIEF EXECUTIVE OFFICER

PETER BOOTH WILEY
CHAIRMAN OF THE BOARD

需要全本请访问 www.ertongbook.com

PREFACE

Our understanding of the fundamental processes of the natural world is based to a large extent on partial differential equations. Examples are the vibrations of solids, the flow of fluids, the diffusion of chemicals, the spread of heat, the structure of molecules, the interactions of photons and electrons, and the radiation of electromagnetic waves. Partial differential equations also play a central role in modern mathematics, especially in geometry and analysis. The availability of powerful computers is gradually shifting the emphasis in partial differential equations away from the analytical computation of solutions and toward both their numerical analysis and the qualitative theory.

This book provides an introduction to the basic properties of partial differential equations (PDEs) and to the techniques that have proved useful in analyzing them. My purpose is to provide for the student a broad perspective on the subject, to illustrate the rich variety of phenomena encompassed by it, and to impart a working knowledge of the most important techniques of analysis of the solutions of the equations.

One of the most important techniques is the method of separation of variables. Many textbooks heavily emphasize this technique to the point of excluding other points of view. The problem with that approach is that only certain kinds of partial differential equations can be solved by it, whereas others cannot. In this book it plays a very important but not an overriding role. Other texts, which bring in relatively advanced theoretical ideas, require too much mathematical knowledge for the typical undergraduate student. I have tried to minimize the advanced concepts and the mathematical jargon in this book. However, because partial differential equations is a subject at the forefront of research in modern science, I have not hesitated to mention advanced ideas as further topics for the ambitious student to pursue.

This is an undergraduate textbook. It is designed for juniors and seniors who are science, engineering, or mathematics majors. Graduate students, especially in the sciences, could surely learn from it, but it is in no way conceived of as a graduate text.

The main prerequisite is a solid knowledge of calculus, especially multivariate. The other prerequisites are small amounts of ordinary differential

equations and of linear algebra, each much less than a semester's worth. However, since the subject of partial differential equations is by its very nature not an easy one, I have recommended to my own students that they should already have taken full courses in these two subjects.

The presentation is based on the following principles. Motivate with physics but then do mathematics. Focus on the three classical equations: All the important ideas can be understood in terms of them. Do one spatial dimension before going on to two and three dimensions with their more complicated geometries. Do problems without boundaries before bringing in boundary conditions. (By the end of Chapter 2, the student will already have an intuitive and analytical understanding of simple wave and diffusion phenomena.) Do not hesitate to present some facts without proofs, but provide the most critical proofs. Provide introductions to a variety of important advanced topics.

There is plenty of material in this book for a year-long course. A quarter course, or a fairly relaxed semester course, would cover the starred sections of Chapters 1 to 6. A more ambitious semester course could supplement the basic starred sections in various ways. The unstarred sections in Chapters 1 to 6 could be covered as desired. A computational emphasis following the starred sections would be provided by the numerical analysis of Chapter 8. To resume separation of variables after Chapter 6, one would take up Chapter 10. For physics majors one could do some combination of Chapters 9, 12, 13, and 14. A traditional course on boundary value problems would cover Chapters 1, 4, 5, 6, and 10.

Each chapter is divided into sections, denoted A.B. An equation numbered (A.B.C) refers to equation (C) in section A.B. A reference to equation (C) refers to the equation in the same section. A similar system is used for numbering theorems and exercises. The references are indicated by brackets, like [AS].

The help of my colleagues is gratefully acknowledged. I especially thank Yue Liu and Brian Loe for their extensive help with the exercises, as well as Costas Dafermos, Bob Glassey, Jerry Goldstein, Manos Grillakis, Yan Guo, Chris Jones, Keith Lewis, Gustavo Perla Menzala, and Bob Seeley for their suggestions and corrections.

Walter A. Strauss

PREFACE TO SECOND EDITION

In the years since the first edition came out, partial differential equations has become yet more prominent, both as a model for scientific theories and within mathematics itself. In this second edition I have added 30 new exercises. Furthermore, this edition is accompanied by a solutions manual that has answers to about half of the exercises worked out in detail. I have added a new section on water waves as well as new material and explanatory comments in many places. Corrections have been made wherever necessary.

I would like to take this opportunity to thank all the people who have pointed out errors in the first edition or made useful suggestions, including Andrew Bernoff, Rustum Choksi, Adrian Constantin, Leonid Dickey, Julio Dix, Craig Evans, A. M. Fink, Robert Glassey, Jerome Goldstein, Leon Greenberg, Chris Hunter, Eva Kallin, Jim Kelliher, Jeng-Eng Lin, Howard Liu, Jeff Nunemacher, Vassilis Papanicolaou, Mary Pugh, Stan Richardson, Stuart Rogers, Paul Sacks, Naoki Saito, Stephen Simons, Catherine Sulem, David Wagner, David Weinberg, and Nick Zakrasek. My warmest thanks go to Julie and Steve Levandosky who, besides being my co-authors on the solutions manual, provided many suggestions and much insight regarding the text itself.

目 录

（标星章节构成本书核心内容）

第1章 偏微分方程的由来

1.1* 什么是偏微分方程? …… 1
1.2* 一阶线性方程 …… 6
1.3* 流、振动与扩散 …… 10
1.4* 初始条件与边界条件 …… 20
1.5 适定性问题 …… 25
1.6 二阶方程的类型 …… 28

第2章 波与扩散

2.1* 波动方程 …… 33
2.2* 因果律与能量 …… 39
2.3* 扩散方程 …… 42
2.4* 全直线上的扩散方程 …… 46
2.5* 波与扩散的比较 …… 54

第3章 反射与源

3.1 半直线上的扩散方程 …… 57
3.2 波的反射 …… 61
3.3 具有源的扩散 …… 67
3.4 具有源的波动 …… 71
3.5 再论扩散 …… 80

第4章 边界值问题

4.1* 分离变量法与Dirichlet条件 …… 84

4.2* Neumann条件 …… 89
4.3* Robin条件 …… 92

第5章 Fourier级数
5.1* Fourier系数 …… 104
5.2* 偶函数、奇函数、周期函数与复函数 …… 113
5.3* 正交性及广义Fourier级数 …… 118
5.4* 完备性 …… 124
5.5 完备性与Gibbs现象 …… 136
5.6 非齐次边界条件 …… 147

第6章 调和函数
6.1* Laplace方程 …… 152
6.2* 矩形域与立方体 …… 161
6.3* Poisson公式 …… 165
6.4 圆形域、楔形域与圆环域 …… 172

（以下四章可以按任何顺序学习）

第7章 Green公式与Green函数
7.1 Green第一公式 …… 178
7.2 Green第二公式 …… 185
7.3 Green函数 …… 188
7.4 半空间与球面 …… 191

第8章 解的计算
8.1 机遇与风险 …… 199
8.2 扩散方程的逼近 …… 203
8.3 波动方程的逼近 …… 211
8.4 Laplace方程的逼近 …… 218
8.5 有限元方法 …… 222

第9章 全空间中的波
9.1 能量与因果律 …… 228

9.2 时空中的波动方程 …… 234
9.3 射线、奇异性及源 …… 242
9.4 扩散方程与Schrödinger方程 …… 248
9.5 氢原子 …… 254

第10章 平面与空间中的边界值问题
10.1 再论Fourier方法 …… 258
10.2 鼓面的振动 …… 264
10.3 球内的固态振动 …… 270
10.4 结点 …… 278
10.5 Bessel函数 …… 282
10.6 Legendre函数 …… 289
10.7 量子力学中的角动量 …… 294

第11章 广义本征值问题
11.1 本征值是势能函数的极小值 …… 299
11.2 本征值的计算 …… 304
11.3 完备性 …… 310
11.4 对称微分算子 …… 314
11.5 完备性与分离变量 …… 318
11.6 本征值的渐近值 …… 322

第12章 分布与变换
12.1 分布 …… 331
12.2 再论Green函数 …… 338
12.3 Fourier变换 …… 343
12.4 源函数 …… 349
12.5 Laplace变换 …… 353

第13章 物理学中的偏微分方程问题
13.1 电磁学 …… 358
13.2 流体力学与声学 …… 361
13.3 散射 …… 366

需要全本请访问 www.ertongbook.com

13.4 连续谱 …… 370
13.5 基本粒子方程 …… 373

第14章 非线性偏微分方程
14.1 激波 …… 380
14.2 孤立子 …… 390
14.3 变分法 …… 397
14.4 分叉理论 …… 401
14.5 水波 …… 406

附录
A.1 连续函数与可微函数 …… 414
A.2 函数项无穷级数 …… 418
A.3 微分与积分 …… 420
A.4 微分方程 …… 423
A.5 Gamma函数 …… 425

参考书目 …… 427
部分习题答案与提示 …… 431

1

WHERE PDEs COME FROM

After thinking about the meaning of a partial differential equation, we will flex our mathematical muscles by solving a few of them. Then we will see how naturally they arise in the physical sciences. The physics will motivate the formulation of boundary conditions and initial conditions.

1.1 WHAT IS A PARTIAL DIFFERENTIAL EQUATION?

The key defining property of a partial differential equation (PDE) is that there is more than one independent variable $x, y, \ldots$. There is a dependent variable that is an unknown function of these variables $u(x, y, \ldots)$. We will often denote its derivatives by subscripts; thus $\partial u/\partial x = u_x$, and so on. A PDE is an identity that relates the independent variables, the dependent variable u, and the partial derivatives of u. It can be written as

$$F(x, y, u(x, y), u_x(x, y), u_y(x, y)) = F(x, y, u, u_x, u_y) = 0. \tag{1}$$

This is the most general PDE in two independent variables of *first* order. The *order* of an equation is the highest derivative that appears. The most general *second*-order PDE in two independent variables is

$$F(x, y, u, u_x, u_y, u_{xx}, u_{xy}, u_{yy}) = 0. \tag{2}$$

A *solution* of a PDE is a function $u(x, y, \ldots)$ that satisfies the equation identically, at least in some region of the $x, y, \ldots$ variables.

When solving an ordinary differential equation (ODE), one sometimes reverses the roles of the independent and the dependent variables—for instance, for the separable ODE $\dfrac{du}{dx} = u^3$. For PDEs, the distinction between the independent variables and the dependent variable (the unknown) is always maintained.

Some examples of PDEs (all of which occur in physical theory) are:

1. $u_x + u_y = 0$ (transport)
2. $u_x + yu_y = 0$ (transport)
3. $u_x + uu_y = 0$ (shock wave)
4. $u_{xx} + u_{yy} = 0$ (Laplace's equation)
5. $u_{tt} - u_{xx} + u^3 = 0$ (wave with interaction)
6. $u_t + uu_x + u_{xxx} = 0$ (dispersive wave)
7. $u_{tt} + u_{xxxx} = 0$ (vibrating bar)
8. $u_t - iu_{xx} = 0$ $(i = \sqrt{-1})$ (quantum mechanics)

Each of these has two independent variables, written either as x and y or as x and t. Examples 1 to 3 have order one; 4, 5, and 8 have order two; 6 has order three; and 7 has order four. Examples 3, 5, and 6 are distinguished from the others in that they are not "linear." We shall now explain this concept.

Linearity means the following. Write the equation in the form $\mathcal{L}u = 0$, where $\mathcal{L}$ is an *operator*. That is, if v is any function, $\mathcal{L}v$ is a new function. For instance, $\mathcal{L} = \partial/\partial x$ is the operator that takes v into its partial derivative v_x. In Example 2, the operator $\mathcal{L}$ is $\mathcal{L} = \partial/\partial x + y\partial/\partial y$. ($\mathcal{L}u = u_x + yu_y$.) The definition we want for linearity is

$$\mathcal{L}(u + v) = \mathcal{L}u + \mathcal{L}v \qquad \mathcal{L}(cu) = c\mathcal{L}u \tag{3}$$

for any functions u, v and any constant c. Whenever (3) holds (for all choices of u, v, and c), $\mathcal{L}$ is called *linear operator*. The equation

$$\mathcal{L}u = 0 \tag{4}$$

is called *linear* if $\mathcal{L}$ is a linear operator. Equation (4) is called a *homogeneous linear equation*. The equation

$$\mathcal{L}u = g, \tag{5}$$

where $g \neq 0$ is a given function of the independent variables, is called an *inhomogeneous linear equation*. For instance, the equation

$$(\cos xy^2)u_x - y^2u_y = \tan(x^2 + y^2) \tag{6}$$

is an inhomogeneous linear equation.

As you can easily verify, five of the eight equations above are linear as well as homogeneous. Example 5, on the other hand, is not linear because although $(u + v)_{xx} = u_{xx} + v_{xx}$ and $(u + v)_{tt} = u_{tt} + v_{tt}$ satisfy property (3), the cubic term does not:

$$(u + v)^3 = u^3 + 3u^2v + 3uv^2 + v^3 \neq u^3 + v^3.$$

The advantage of linearity for the equation $\mathcal{L}u = 0$ is that if u and v are both solutions, so is $(u + v)$. If $u_1, \ldots, u_n$ are all solutions, so is any linear combination

$$c_1u_1(x) + \cdots + c_nu_n(x) = \sum_{j=1}^{n} c_ju_j(x) \qquad (c_j = \text{constants}).$$

(This is sometimes called the superposition principle.) Another consequence of linearity is that if you add a homogeneous solution [a solution of (4)] to an inhomogeneous solution [a solution of (5)], you get an inhomogeneous solution. (Why?) The mathematical structure that deals with linear combinations and linear operators is the vector space. Exercises 5–10 are review problems on vector spaces.

We'll study, almost exclusively, linear systems with constant coefficients. Recall that for ODEs you get linear combinations. The coefficients are the arbitrary constants. *For an ODE of order m, you get m arbitrary constants.*

Let's look at some PDEs.

Example 1.

Find all $u(x, y)$ satisfying the equation $u_{xx} = 0$. Well, we can integrate once to get $u_x =$ constant. But that's not really right since there's another variable y. What we really get is $u_x(x, y) = f(y)$, where $f(y)$ is arbitrary. Do it again to get $u(x, y) = f(y)x + g(y)$. This is the solution formula. Note that *there are two arbitrary* functions *in the solution.* We see this as well in the next two examples. □

Example 2.

Solve the PDE $u_{xx} + u = 0$. Again, it's really an ODE with an extra variable y. We know how to solve the ODE, so the solution is

$$u = f(y)\cos x + g(y)\sin x,$$

where again $f(y)$ and $g(y)$ are two arbitrary functions of y. You can easily check this formula by differentiating twice to verify that $u_{xx} = -u$. □

Example 3.

Solve the PDE $u_{xy} = 0$. This isn't too hard either. First let's integrate in x, regarding y as fixed. So we get

$$u_y(x, y) = f(y).$$

Next let's integrate in y regarding x as fixed. We get the solution

$$u(x, y) = F(y) + G(x),$$

where $F' = f$. □

Moral *A PDE has arbitrary functions in its solution.* In these examples the arbitrary functions are functions of one variable that combine to produce a function $u(x, y)$ of two variables which is only partly arbitrary.

A function of two variables contains *immensely* more information than a function of only one variable. Geometrically, it is obvious that a surface $\{u = f(x, y)\}$, the graph of a function of two variables, is a much more complicated object than a curve $\{u = f(x)\}$, the graph of a function of one variable.

To illustrate this, we can ask how a computer would record a function $u = f(x)$. Suppose that we choose 100 points to describe it using equally spaced values of x: $x_1, x_2, x_3, \ldots, x_{100}$. We could write them down in a column, and next to each x_j we could write the corresponding value $u_j = f(x_j)$. Now how about a function $u = f(x, y)$? Suppose that we choose 100 equally spaced values of x and also of y: $x_1, x_2, x_3, \ldots, x_{100}$ and $y_1, y_2, y_3, \ldots, y_{100}$. Each *pair* x_i, y_j provides a value $u_{ij} = f(x_i, y_j)$, so there will be $100^2 = 10{,}000$ lines of the form

$$x_i \qquad y_j \qquad u_{ij}$$

required to describe the function! (If we had a prearranged system, we would need to record only the values u_{ij}.) A function of three variables described discretely by 100 values in each variable would require a million numbers!

To understand this book what do you have to know from calculus? Certainly all the basic facts about partial derivatives and multiple integrals. For a brief discussion of such topics, see the Appendix. Here are a few things to keep in mind, some of which may be new to you.

1. Derivatives are *local*. For instance, to calculate the derivative $(\partial u/\partial x)(x_0, t_0)$ at a particular point, you need to know just the values of $u(x, t_0)$ for x *near* x_0, since the derivative is the limit as $x \to x_0$.
2. Mixed derivatives are equal: $u_{xy} = u_{yx}$. (We assume throughout this book, unless stated otherwise, that all derivatives exist and are continuous.)
3. The chain rule is used frequently in PDEs; for instance,
$$\frac{\partial}{\partial x}[f(g(x, t))] = f'(g(x, t)) \cdot \frac{\partial g}{\partial x}(x, t).$$
4. For the integrals of derivatives, the reader should learn or review Green's theorem and the divergence theorem. (See the end of Section A.3 in the Appendix.)
5. Derivatives of integrals like $I(t) = \int_{a(t)}^{b(t)} f(x, t)\, dx$ (see Section A.3).
6. Jacobians (change of variable in a double integral) (see Section A.1).
7. Infinite series of functions and their differentiation (see Section A.2).
8. Directional derivatives (see Section A.1).
9. We'll often reduce PDEs to ODEs, so we must know how to solve simple ODEs. But we won't need to know anything about tricky ODEs.

EXERCISES

1. Verify the linearity and nonlinearity of the eight examples of PDEs given in the text, by checking whether or not equations (3) are valid.
2. Which of the following operators are linear?
   (a) $\mathcal{L}u = u_x + xu_y$
   (b) $\mathcal{L}u = u_x + uu_y$
   (c) $\mathcal{L}u = u_x + u_y^2$
   (d) $\mathcal{L}u = u_x + u_y + 1$
   (e) $\mathcal{L}u = \sqrt{1+x^2}\,(\cos y)u_x + u_{yxy} - [\arctan(x/y)]u$
3. For each of the following equations, state the order and whether it is nonlinear, linear inhomogeneous, or linear homogeneous; provide reasons.
   (a) $u_t - u_{xx} + 1 = 0$
   (b) $u_t - u_{xx} + xu = 0$
   (c) $u_t - u_{xxt} + uu_x = 0$
   (d) $u_{tt} - u_{xx} + x^2 = 0$
   (e) $iu_t - u_{xx} + u/x = 0$
   (f) $u_x(1+u_x^2)^{-1/2} + u_y(1+u_y^2)^{-1/2} = 0$
   (g) $u_x + e^y u_y = 0$
   (h) $u_t + u_{xxxx} + \sqrt{1+u} = 0$
4. Show that the difference of two solutions of an inhomogeneous linear equation $\mathcal{L}u = g$ with the same g is a solution of the homogeneous equation $\mathcal{L}u = 0$.
5. Which of the following collections of 3-vectors $[a, b, c]$ are vector spaces? Provide reasons.
   (a) The vectors with $b = 0$.
   (b) The vectors with $b = 1$.
   (c) The vectors with $ab = 0$.
   (d) All the linear combinations of the two vectors $[1, 1, 0]$ and $[2, 0, 1]$.
   (e) All the vectors such that $c - a = 2b$.
6. Are the three vectors $[1, 2, 3]$, $[-2, 0, 1]$, and $[1, 10, 17]$ linearly dependent or independent? Do they span all vectors or not?
7. Are the functions $1 + x$, $1 - x$, and $1 + x + x^2$ linearly dependent or independent? Why?
8. Find a vector that, together with the vectors $[1, 1, 1]$ and $[1, 2, 1]$, forms a basis of $\mathbb{R}^3$.
9. Show that the functions $(c_1 + c_2 \sin^2 x + c_3 \cos^2 x)$ form a vector space. Find a basis of it. What is its dimension?
10. Show that the solutions of the differential equation $u''' - 3u'' + 4u = 0$ form a vector space. Find a basis of it.
11. Verify that $u(x, y) = f(x)g(y)$ is a solution of the PDE $uu_{xy} = u_x u_y$ for all pairs of (differentiable) functions f and g of one variable.

需要全本请访问 www.ertongbook.com